

## Medical Policy



### Title: Magnetic Resonance Imaging (MRI) of the Breast

<b>Professional / Institutional</b>
Original Effective Date: October 1, 1997 / July 1, 2005
Latest Review Date: October 24, 2023
Current Effective Date: November 22, 2022

**State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).**

**The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.**

**The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.**

**If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.**

Populations	Interventions	Comparators	Outcomes
<b>Individuals:</b> <ul style="list-style-type: none"> <li>Who are asymptomatic with high risk of breast cancer</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging as an adjunct to screen for breast cancer</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Mammography</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>Who are asymptomatic with average risk of breast cancer</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging as an adjunct to screen for breast cancer</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Mammography</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> </ul>

Populations	Interventions	Comparators	Outcomes
			<ul style="list-style-type: none"> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With characteristics limiting accuracy of mammography (eg, dense breasts)</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging as an adjunct to screen for breast cancer</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Mammography</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With suspected occult breast primary tumor with axillary nodal adenocarcinoma with negative mammography</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging as an adjunct to detect breast cancer eligible for breast-conserving therapy</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Preemptive mastectomy</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With breast cancer</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Adjunctive magnetic resonance imaging of the contralateral breast</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Mammography and clinical assessment alone</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With low-suspicion findings on conventional mammography</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging as an adjunct to detect breast cancer</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Standard care with short-interval mammographic follow-up</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With suspicious breast lesions</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging as an adjunct to further characterize lesions</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Biopsy based on mammography and clinical assessment</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>

Populations	Interventions	Comparators	Outcomes
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With clinically localized breast cancer</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging for preoperative mapping to identify multicentric disease</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Standard workup without magnetic resonance imaging</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With locally advanced breast cancer undergoing neoadjuvant chemotherapy</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging to guide surgical decisions after neoadjuvant chemotherapy</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Mammography</li> <li>Clinical assessment</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With posteriorly located breast tumors</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging to diagnose chest wall involvement</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Mammography</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With a suspicious breast lesion recommended for biopsy but not localizable by mammography or ultrasonography</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging to evaluate and localize the lesion prior to biopsy</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Waiting until lesion becomes palpable or visible on mammography or ultrasonography</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With locally advanced breast cancer undergoing neoadjuvant chemotherapy</li> </ul>	<b>Interventions of interest are:</b> <ul style="list-style-type: none"> <li>Magnetic resonance imaging to evaluate response to chemotherapy</li> </ul>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Clinical assessment</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>
<b>Individuals:</b> <ul style="list-style-type: none"> <li>With positive surgical margins after</li> </ul>	<b>Interventions of interest are:</b>	<b>Comparators of interest are:</b> <ul style="list-style-type: none"> <li>Pathologic inspection</li> </ul>	<b>Relevant outcomes include:</b> <ul style="list-style-type: none"> <li>Overall survival</li> </ul>

Populations	Interventions	Comparators	Outcomes
lumpectomy or breast conservation surgery	<ul style="list-style-type: none"> <li>Magnetic resonance imaging to evaluate residual tumor</li> </ul>		<ul style="list-style-type: none"> <li>Disease-specific survival</li> <li>Test accuracy</li> <li>Test validity</li> <li>Resource utilization</li> </ul>

## DESCRIPTION

Magnetic resonance imaging (MRI) of the breast is performed using scanners and intravenous imaging contrast agents in combination with specialized breast coils. This evidence review only addresses the use of breast MRI for clinical indications related to the detection or diagnosis of breast cancer.

## OBJECTIVE

The objective of this evidence review is to determine whether magnetic resonance imaging of the breast improves the net health outcome for individuals undergoing breast cancer screening, breast cancer detection, and/or evaluation for breast cancer before and/or after treatment.

## BACKGROUND

### Health Disparities in Breast Cancer

Based on data from 2014 through 2018, age-adjusted breast cancer mortality is approximately 40% higher among Black women compared to non-Hispanic White women in the United States (27.7 vs. 20.0 deaths per 100,000 women), despite a lower overall incidence of breast cancer among Black women (125.8 vs. 139.2 cases per 100,000 women).<sup>1</sup> Experts postulate that this divergence in mortality may be related to access issues; Black women are more likely than White women to lack health insurance limiting access to screening and appropriate therapies. Socioeconomic status is also a driver in health and health outcome disparities related to breast cancer.<sup>2</sup> Women with low incomes have significantly lower rates of breast cancer screening, a higher probability of late-stage diagnosis, and are less likely to receive high quality care, resulting in higher mortality from breast cancer.

### Magnetic Resonance Imaging

Magnetic resonance imaging (MRI) of the breast can be used to screen, detect, and/or diagnose breast cancer. An MRI can be used as a replacement for mammography screening, or as an additional imaging test alone, or in combination with other imaging modalities. Each potential use is described below.

## REGULATORY STATUS

### Health Disparities in Breast Cancer

Based on data from 2014 through 2018, age-adjusted breast cancer mortality is approximately 40% higher among Black women compared to non-Hispanic White women in the United States

(27.7 vs. 20.0 deaths per 100,000 women), despite a lower overall incidence of breast cancer among Black women (125.8 vs. 139.2 cases per 100,000 women).<sup>1</sup> Experts postulate that this divergence in mortality may be related to access issues; Black women are more likely than White women to lack health insurance limiting access to screening and appropriate therapies. Socioeconomic status is also a driver in health and health outcome disparities related to breast cancer.<sup>2</sup> Women with low incomes have significantly lower rates of breast cancer screening, a higher probability of late-stage diagnosis, and are less likely to receive high quality care, resulting in higher mortality from breast cancer.

**Magnetic Resonance Imaging**

Magnetic resonance imaging (MRI) of the breast can be used to screen, detect, and/or diagnose breast cancer. An MRI can be used as a replacement for mammography screening, or as an additional imaging test alone, or in combination with other imaging modalities. Each potential use is described below.

**POLICY**

- A. MRI of the breast may be considered **medically necessary** for screening for breast cancer in individuals with high risk of breast cancer including but not limited to the following: (see Policy Guidelines section.)
1. With a known *BRCA1* or *BRCA2* variant; or
  2. At high risk of *BRCA1* or *BRCA2* variant due to a known presence of the variant in relatives; or
  3. Who have Li-Fraumeni syndrome or Cowden syndrome or Bannayan-Riley-Ruvalcaba syndrome or who have a first-degree relative with one of these syndromes; or
  4. At high risk (lifetime risk about 20% to 25% or greater) of developing breast cancer as identified by models that are largely defined by family history; or
  5. Who received radiation therapy to the chest between 10 and 30 years of age.
- B. MRI of the breast may be considered **medically necessary** for the following:
1. For detection of a suspected occult breast primary tumor in individuals with axillary nodal adenocarcinoma (i.e., negative mammography and physical exam).
  2. To confirm the clinical diagnosis of rupture of silicone breast implant.
  3. For presurgical planning in individuals with locally advanced breast cancer before and after completion of neoadjuvant chemotherapy to permit tumor localization and characterization.
  4. To determine the presence of pectoralis major muscle/chest wall invasion in individuals with posteriorly located tumors.
  5. In those with a new diagnosis of breast cancer.
  6. For preoperative tumor mapping of the involved (ipsilateral) breast to evaluate the presence of multicentric disease in individuals with clinically localized breast cancer who are candidates for breast-conservation therapy (see Policy Guidelines).
  7. To evaluate a documented abnormality of the breast prior to obtaining an MRI-guided biopsy when there is documentation that other methods, such as palpation or ultrasound, are not able to localize the lesion for biopsy.
  8. Further evaluation of suspicious clinical findings or imaging results, which remain indeterminate after complete mammographic and sonographic evaluation, combined with a thorough physical examination.

9. To detect the extent of residual cancer in the recently postoperative breast with positive pathological margins after incomplete lumpectomy when the individual still desires breast conservation and local re-excision is planned.
- C. MRI of the breast is considered **experimental / investigational** for the following:
1. As a screening technique in average-risk individuals.
  2. As a screening technique for the detection of breast cancer when the sensitivity of mammography (i.e., mammography using low-dose x-rays for imaging) is limited (i.e., dense breasts, breast implants, scarring after treatment for breast cancer).
  3. For diagnosis of low-suspicion findings on conventional testing not indicated for immediate biopsy and referred for short-interval follow-up.
  4. For diagnosis of a suspicious breast lesion in order to avoid biopsy.
  5. To determine response during neoadjuvant chemotherapy in individuals with locally advanced breast cancer.
  6. To monitor the integrity of silicone gel-filled breast implants when there are no signs or symptoms of rupture.

**Note:** All of the policy statements above refer to performing MRI of the breast with a breast coil and the use of contrast. MRI of the breast without the use of a breast coil, regardless of the clinical indication, is considered **experimental / investigational**.

### **POLICY GUIDELINES**

- A. Families at high risk for harboring a *BRCA1* or *BRCA2* variant are those in which the incidence of breast or ovarian cancer in first-degree (i.e., parent, sibling, offspring) or second-degree (i.e., grandparent, grandchild, uncle, aunt, nephew, niece, half-sibling) relatives suggests an autosomal dominant inheritance, i.e., about half the family members are affected.
- B. A number of risk assessment tools based mainly on family history can assist practitioners in estimating breast cancer risk and include the Claus,<sup>70</sup> modified Gail,<sup>71</sup> Tyrer-Cuzick,<sup>72</sup> and BRCAPRO<sup>73</sup> models.
- C. Breast MRI exams should be performed and interpreted by an expert breast imaging team working together with the multidisciplinary oncology treatment team.
- D. As noted, breast MRI exams require a dedicated breast coil and the use of contrast by radiologists familiar with the optimal timing sequences and other technical aspects of image interpretation. The breast MRI center should also have the ability to perform MRI-guided biopsy and/or wire localization of findings detected by MRI.
  1. Preoperative MRI in individuals with localized disease apparently results in higher rates of mastectomy and lower rates of breast-conserving therapy (BCT). There is uncertainty from the available evidence on whether outcomes are improved by changing to a more extensive operation. If biopsies are performed on all MRI-identified lesions, and if shared individual decision making is used for altering the surgical approach, then the probability of improved outcomes is increased

**Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.**

## **RATIONALE**

This evidence review has been updated regularly with searches of the PubMed database. The most recent literature update was performed through July 21, 2023.

Evidence reviews assess whether a medical test is clinically useful. A useful test provides information to make a clinical management decision that improves the net health outcome. That is, the balance of benefits and harms is better when the test is used to manage the condition than when another test or no test is used to manage the condition.

The first step in assessing a medical test is to formulate the clinical context and purpose of the test. The test must be technically reliable, clinically valid, and clinically useful for that purpose. Evidence reviews assess the evidence on whether a test is clinically valid and clinically useful. Technical reliability is outside the scope of these reviews, and credible information on technical reliability is available from other sources.

Promotion of greater diversity and inclusion in clinical research of historically marginalized groups (e.g., People of Color [African-American, Asian, Black, Latino and Native American]; LGBTQIA (Lesbian, Gay, Bisexual, Transgender, Queer, Intersex, Asexual); Women; and People with Disabilities [Physical and Invisible]) allows policy populations to be more reflective of and findings more applicable to our diverse members. While we also strive to use inclusive language related to these groups in our policies, use of gender-specific nouns (e.g., women, men, sisters, etc.) will continue when reflective of language used in publications describing study populations.

## **SCREENING USES**

### **SCREENING INDIVIDUALS AT HIGH-RISK OF BREAST CANCER**

#### **Clinical Context and Test Purpose**

Screening uses include screening for breast cancer in patients who are at high genetic risk for breast cancer. Magnetic resonance imaging (MRI) of the breast has been investigated as a screening tool in specific higher-risk subgroups of patients. First, it has been studied in patients considered to be at high genetic risk of breast cancer, such as women with known *BRCA1* or *BRCA2* genetic variants or with a family history consistent with a hereditary pattern of breast cancer. Screening for breast cancer often begins at an earlier age in these patients, and mammography is considered less sensitive in younger patients due to the prevalence of dense breast tissue.

The questions addressed in this portion of the evidence review:

- Does the use of MRI as an adjunct to screening for breast cancer improve the net health outcome of patients at high-risk of breast cancer compared with standard mammographic techniques?
- Is this degree of increased accuracy likely to improve net health outcomes via earlier diagnosis and treatment?



The following PICO was used to select literature to inform this review.

**Populations**

The population of interest is patients at high-risk of developing breast cancer.

**Interventions**

The intervention of interest is MRI as an adjunct to screening with mammography.

**Comparators**

The following test is currently being used to make decisions about managing breast cancer: mammography alone.

**Outcomes**

The outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility are overall mortality and breast cancer-specific mortality. Another outcome of interest for clinical utility is resource utilization (eg, need for additional testing or procedures).

Breast MRI is performed as an adjunct to routine screening; timing can be guided by national guidelines on breast cancer screening (see Supplemental Information section).

**Study Selection Criteria**

This evidence review focuses on systematic reviews. For the evaluation of the clinical validity of MRI as an adjunct to screening with mammography, we sought systematic reviews that focused on studies meeting the following eligibility criteria:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard
- Patient/sample clinical characteristics were described
- Patient/sample selection criteria were described.

**Clinically Valid**

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

**REVIEW OF EVIDENCE****Systematic Reviews**

Three systematic reviews identified have included women at high-risk of developing breast cancer. Warner et al (2008) reviewed 11 studies published through 2008.<sup>4</sup> Two reviews by Phi et al (2015, 2017) reported 2 individual patient data meta-analyses from the same 6 studies published between 2010 and 2013.<sup>5,6</sup> Phi et al (2015) included women with *BRCA1* or *BRCA2* variants and Phi et al (2017) included women with a strong family history of breast cancer without a known variant. Ding et al (2023) included women with *BRCA1* or *BRCA2* variants, personal or family history of breast or ovarian cancer, or history of prior chest irradiation.<sup>7</sup> Characteristics of the systematic reviews are shown in Table 1.

**Table 1. Characteristics of Systematic Reviews Assessing Magnetic Resonance Imaging Screening in High-Risk Women**

Study	Dates	Studies	Participants	N (Range)	Design	Reference Standard
Ding et al (2023) <sup>7</sup> ,	2000-2021	18	Women with <i>BRCA1</i> or <i>BRCA2</i> variants, family or personal history of breast or ovarian cancer, history of chest irradiation	1799 (NR)	Prospective and retrospective	Pathological examination
Phi et al (2017) <sup>6</sup> ,	2010-2013	6	Women with a family history of breast cancer without a known genetic variant	2226	Prospective	Biopsy-confirmed cancer for positive; at least 1 y follow-up for negative
Phi et al (2015) <sup>5</sup> ,	2010-2013	6	Women with <i>BRCA1</i> or <i>BRCA2</i> variants	2033	Prospective	Biopsy-confirmed cancer for positive; at least 1 y follow-up for negative
Warner et al (2008) <sup>4</sup> ,	1995-2008	11	Women at very high-risk of breast cancer ( <i>BRCA1</i> or <i>BRCA2</i> or other variants or family history consistent with hereditary breast cancer)	4983 (41 to 1909)	Prospective	Biopsy-confirmed cancer

NR: not reported

Results of the systematic reviews are shown in Table 2. The reviews concluded that screening breast MRI is more sensitive but less specific than mammography for the detection of invasive cancers in high-risk women. The sensitivity of combined MRI and mammography was approximately 93% or higher in the reviews while the sensitivity of mammography alone was between approximately 40% and 55%. The Warner et al (2008) review did not present a risk of bias or quality assessment of included studies. Phi et al (2015) assessed quality using the QUADAS-2 tool. All included studies were considered good quality.

**Table 2. Results of Systematic Reviews Assessing Magnetic Resonance Imaging Screening in High-Risk Women**

Study	MRI		Mammogram		MRI Plus Mammogram	
	<i>Sensitivity, %</i>	<i>Specificity, %</i>	<i>Sensitivity, %</i>	<i>Specificity, %</i>	<i>Sensitivity, %</i>	<i>Specificity, %</i>
Ding et al (2023) <sup>7</sup> ,						
Mean cancer detection rate	15.4	NR	7.0	NR	16.7	NR
Phi et al (2017) <sup>6</sup> ,						

Study	MRI		Mammogram		MRI Plus Mammogram	
Total N	2226	2226	2226	2226	2226	2226
PE (95% CI)	89 (76 to 96)	83 (77 to 88)	55 (41 to 69)	94 (90 to 96)	98 (86 to 100)	79 (73 to 84)
Phi et al (2015) <sup>5</sup> ,						
Total N	1951	1951	1951	1951	1951	1951
PE (95% CI)	85 (69 to 94)	85 (79 to 89)	40 (30 to 50)	94 (89 to 97)	93 (80 to 98)	80 (73 to 86)
Warner et al (2008) <sup>4</sup> ,						
Total N	15576	15576	15496	15496	6781	6781
PE (95% CI)	77 (70 to 84)	86 (81 to 92)	39 (37 to 41)	95 (93 to 97)	94 (90 to 97)	77 (75 to 80)

CI: confidence interval; MRI: magnetic resonance imaging; NR: not reported; PE: pooled estimate.

### Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

The clinical usefulness of MRI as an adjunct to mammography for screening individuals at high risk of breast cancer is supported by an indirect chain of evidence. The clinical validity of MRI for screening in high-risk women has been demonstrated in good quality studies. Breast MRI is more sensitive but less specific than mammography for detecting invasive cancers in high-risk women and the sensitivity of combined MRI and mammography is approximately 93% or higher. Given the high likelihood of malignancy among women at high-risk for breast cancer, the benefits of detecting cancer earlier with adjunctive MRI outweigh the disadvantages of incurring more unnecessary workups and biopsies due to false-positive results.

### Section Summary: Screening Individuals at High-Risk of Breast Cancer

Breast MRI is more sensitive than mammography in detecting malignancy during screening. Because of the high likelihood of malignancy among women at high-risk for breast cancer, the benefits of detecting cancer earlier with adjunctive MRI outweigh the disadvantages of incurring more unnecessary workups and biopsies due to false-positive results.

## SCREENING INDIVIDUALS AT AVERAGE-RISK OF BREAST CANCER

### Clinical Context and Test Purpose

The questions addressed in this portion of the evidence review:

- Does the use of MRI as an adjunct to screening for breast cancer improve the net health outcome of patients who are asymptomatic with average-risk of developing breast cancer?
- Is this degree of increased accuracy likely to improve net health outcomes via the earlier diagnosis and treatment?

The following PICO was used to select literature to inform this review.

***Populations***

The relevant population of interest is patients at average-risk of developing breast cancer.

***Interventions***

The intervention of interest is MRI as an adjunct to screening with mammography.

***Comparators***

The following test is currently being used to make decisions about managing breast cancer: mammography alone.

***Outcomes***

The outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility are overall mortality and breast cancer-specific mortality. Another outcome of interest for clinical utility is resource utilization (eg, need for additional testing or procedures).

Breast MRI is performed as an adjunct to routine screening; timing can be guided by national guidelines on breast cancer screening (see Supplemental Information section).

**Study Selection Criteria**

This evidence review focuses on systematic reviews. For the evaluation of the clinical validity of MRI as an adjunct to screening with mammography, we sought systematic reviews that focused on studies meeting the following eligibility criteria:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard
- Patient/sample clinical characteristics were described
- Patient/sample selection criteria were described.

**Clinically Valid**

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

**REVIEW OF EVIDENCE****Systematic Reviews**

In a systematic review of literature conducted by Nelson et al (2016) for the U.S. Preventive Services Task Force breast cancer screening recommendation update, no randomized controlled trials (RCTs) or nonrandomized observational studies identified evaluated adjunctive MRI for screening average-risk women for breast cancer.<sup>8</sup> Because the prevalence of breast cancer is extremely low in average-risk young women, screening with a test such as MRI that has lower specificity would result in a lower positive predictive value (PPV) and many more false-positive results. Compared with mammography, there would be greater numbers of workups and biopsies with increased anxiety and morbidity with adjunctive MRI screening applied to young, average-risk women.

Health Quality Ontario (2016) published a systematic review of MRI as an adjunct to mammography for women, not at high-risk of breast cancer.<sup>9</sup> Reviewers searched for studies evaluating screening breast MRI as an adjunct to mammography compared with mammography alone. Studies needed to use pathology results as a reference standard for positive tests and clinical follow-up as a reference standard for negative tests. In addition, studies needed to report one or more outcomes of interest, which included effectiveness outcomes (eg, mortality, health-related quality of life, screening-related harms), diagnostic outcomes (eg, sensitivity, specificity), and biopsy and recall rates. Reviewers did not find any studies that met eligibility criteria. They concluded that there was a lack of evidence to inform the questions of the diagnostic accuracy of MRI plus mammography versus MRI alone and the impact of adjunct screening MRI on health outcomes in patients at less than high-risk of breast cancer.

### **Section Summary: Screening of Individuals at Average-Risk of Breast Cancer**

The U.S. Preventative Services Task Force systematic review and guideline concluded that because the prevalence of breast cancer is low in average risk young women, screening with MRI, which has lower specificity, would result in a lower PPV and many more false positive results. A systematic review by Health Quality Ontario concluded that there was lack of evidence on the impact of MRI on health outcomes of individuals at less than high risk of breast cancer.

## **SCREENING WHEN BREAST CHARACTERISTICS LIMIT THE SENSITIVITY OF MAMMOGRAPHY**

### **Clinical Context and Test Purpose**

Screening MRI has been suggested for patients who may or may not be at increased risk but who have breast tissue characteristics that limit the sensitivity of mammographic screening (these characteristics are dense breast tissue, breast implants, or scarring after breast-conserving therapy [BCT]). Use of BCT consists of breast-conserving surgery (BCS) followed by radiotherapy.

The questions addressed in this portion of the evidence review:

- Does the use of MRI as an adjunct to screening for breast cancer improve the net health outcome of patients who are asymptomatic with breast characteristics that limit the sensitivity of mammography?
- Is this degree of increased accuracy likely to improve net health outcomes via the earlier diagnosis and treatment?

The following PICO was used to select literature to inform this review.

### ***Populations***

The population of interest is patients with breast characteristics that limit the sensitivity of mammography. For example, individuals who have dense breasts or prior BCT.

### ***Interventions***

The intervention of interest is MRI as an adjunct to screening with mammography.

### ***Comparators***

The following test is currently being used to make decisions about managing breast cancer: mammography alone.

### Outcomes

The outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility are overall mortality and breast cancer-specific mortality. Another outcome of interest for clinical utility is resource utilization (eg, need for additional testing or procedures).

Breast MRI is performed as an adjunct to routine screening; timing can be guided by national guidelines on breast cancer screening (see the Supplemental Information section).

### Study Selection Criteria

For the evaluation of the clinical validity of MRI as an adjunct to screening with mammography, studies that met the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard
- Patient/sample clinical characteristics were described
- Patient/sample selection criteria were described.

### Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

## REVIEW OF EVIDENCE

### Randomized Controlled and Single Arm Studies: Dense Breasts

One RCT and a prospective observational study were identified that evaluated the use of supplemental MRI in patients who received screening mammography and/or ultrasound. Characteristics of the studies are shown in Table 3.

**Table 3. Characteristics of Clinical Validity Studies Assessing Supplemental Breast Magnetic Resonance Imaging for Routine Screening in Women**

Study	Study Population	Design	Reference Standard	Identification of Positive MRI Test	Timing of Tests	Blinding of Assessors	Comment
Bakker et al (2019) <sup>10</sup>	Women aged 50 to 75 years in the Netherlands with extremely dense breast tissue with negative results on	Randomized controlled trial	<ul style="list-style-type: none"> <li>• Incidence of interval cancers (positive MRI result that was confirmed histologically)</li> </ul>	Assessed as BI-RADS category 4 or 5 by 1 radiologist with 5+ years of experience in breast MRI; Patients	Mammography screening with or without MRI every 2 years	NR	<ul style="list-style-type: none"> <li>• Funded by the University Medical Center Utrecht, the Netherlands Organization for Health</li> </ul>

Study	Study Population	Design	Reference Standard	Identification of Positive MRI Test	Timing of Tests	Blinding of Assessors	Comment
	screening mammography; socioeconomic status was recorded at baseline: 36.1% of women were in the highest status quartile (quartile 1), 23.6% in quartile 3, 22.7% in quartile 2, and 17.4% were in the lowest status quartile 4		during 2-year screening period	with BI-RADS category of 3 received follow-up MRI after 6 months			Research and Development, the Dutch Cancer Society, the Dutch Pink Ribbon-A Sister's Hope organization, Stichting Kankerpreventie Midden-West, Bayer Pharmaceuticals, and Volpara Health Technologies
Berg et al (2012) <sup>11</sup> ,	Women aged 25 years and older with heterogeneously dense or extremely dense breast tissue with at least 1 risk factor for breast cancer. Women had undergone 3 negative screenings	Prospective trial	<ul style="list-style-type: none"> <li>Most severe biopsy result within 365 days of mammographic screening and/or clinical follow-up at 1 year</li> </ul>	Assessed as BI-RADS score of 3, 4, or 5	MRI within 8 weeks of last screening mammography	Yes (interpretation was blinded to other test results)	<ul style="list-style-type: none"> <li>Funded by the Avon Foundation and National Institutes of Health grants</li> </ul>

Study	Study Population	Design	Reference Standard	Identification of Positive MRI Test	Timing of Tests	Blinding of Assessors	Comment
	of mammography and supplemental ultrasound. 93% of women in the study were White; the remainder of women were Hispanic or Latino, Black, Native Hawaiian or Pacific Islander, Asian, or American Indian or Alaskan Native.						

BI-RADS: Breast Imaging Reporting and Data System; MRI: magnetic resonance imaging; NR: not reported.

Results of the clinical validity studies are shown in Table 4. Bakker et al (2019) conducted a multicenter RCT (DENSE) with 40,373 women with extremely dense breast tissue and normal mammography results who were assigned to an optional supplemental MRI or mammography-only screening.<sup>10</sup> There were 8061 patients invited to undergo MRI (MRI-invitation group); however, 4783 patients participated in supplemental MRI screening and 3278 chose not to participate. There were 32,312 patients who only received mammography (mammography-only group). The interval-cancer rate was 2.5 per 1000 screenings in the MRI-invitation group compared to 5.0 per 1000 screenings in the mammography-only group (rate difference, 2.5; 95% confidence interval [CI], 1.0 to 3.7;  $p < .001$ ). Of note, among the 20 interval cancers diagnosed in the MRI-invitation group, 16 were diagnosed in patients who did not accept the supplemental MRI invitation (4.9 per 1000 screenings), while 4 were diagnosed in patients who underwent MRI screening (0.8 per 1000 screenings). In the 2012 ACRIN (American College of Radiology Imaging Network) 6666 trial, mammography alone was compared with mammography plus ultrasound in women 25 years or older with at least heterogeneously dense breast tissue and at least 1 other breast cancer risk factor.<sup>11</sup> Half (54%) of women had a personal history of breast cancer. In a MRI subanalysis, women who completed 3 rounds of screening and did not have contraindications or renal impairment were asked to undergo contrast-enhanced MRI within 8 weeks of the last screening mammography. Six hundred twenty-seven women consented and



were eligible for this subanalysis, and 612 (98%) completed the needed tests; 16 cancers were detected in these women. Sensitivity increased from 44% (95% CI, 20% to 70%) for mammography plus ultrasound to 100% (95% CI, 79% to 100%;  $p=.004$ ) when MRI was added. Specificity declined from 84% (95% CI, 81% to 87%) for mammography plus ultrasound to 65% (95% CI, 61% to 69%;  $p<.001$ ) for all 3 tests. Over the 3 year study period, another 9 cancers were identified between screening tests, and 2 additional cancers were identified off-study.

**Table 4. Results of Clinical Validity Studies Assessing Supplemental Breast Magnetic Resonance Imaging for Routine Screening in Women**

Study	Initial N	Final N	Excluded Images	Cancer Rate	Clinical Validity, % (95% CI)			
					Sensitivity	Specificity	PPV	NPV
Bakker et al (2019) <sup>10</sup> ,	40,373 (8061 were invited to undergo MRI screening )	40,373 (Of 8061 who were invited to undergo MRI screening, 4783 underwent screening)	11 died, 3 moved abroad	<i>Interval Cancer Rate</i>				
MRI invitation + mammography				2.5 per 1000 screenings (95% CI, 1.6 to 3.8)	95.2 (88.1 to 98.7)	92 (NR)	Recall for additional testing: 17.4 (14.2 to 21.2) Biopsy: 26.3 (21.7 to 31.6)	NR
Mammography alone				5.0 per 1000 screenings (95% CI, 4.3 to 5.8)	NR	NR	NR	NR
Berg et al (2012) <sup>11</sup> ,	627 women were screened for the MRI substudy	612 MRI participants	15 were excluded because there was no reference standard	<i>Cancer diagnosis</i>				

Study	Initial N	Final N	Excluded Images	Cancer Rate	Clinical Validity, % (95% CI)			
Supplemental MRI				16 (2.6%) participants	100 (79 to 100)	65 (61 to 69)	19 (11 to 29)	NR
Mammography and ultrasound				NA	44 (20 to 70)	84 (81 to 87)	18 (8 to 34)	NR

CI: confidence interval; NA: not applicable; MRI: magnetic resonance imaging; NPV: negative predictive value; NR: not reported; PPV: positive predictive value.

Tables 5 and 6 discuss relevant limitations of the studies.

**Table 5. Study Relevance Limitations of Clinical Validity Studies of Supplemental Breast Magnetic Resonance Imaging for Routine Screening in Women**

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Duration of Follow-Up <sup>e</sup>
Bakker et al (2019) <sup>10</sup> ,	4. Enrolled populations do not reflect relevant diversity			1. Health outcomes not reported	
Berg et al (2012) <sup>11</sup> ,	4. Enrolled populations not reflect relevant diversity			1. Health outcomes not reported	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

<sup>b</sup> Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest.

<sup>c</sup> Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

<sup>d</sup> Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity, and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

<sup>e</sup> Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true-positives, true-negatives, false-positives, false-negatives cannot be determined).

**Table 6. Study Design and Conduct Limitations of Clinical Validity Studies of Supplemental Breast Magnetic Resonance Imaging for Routine Screening in Women**

Study	Selection <sup>a</sup>	Blinding <sup>b</sup>	Delivery of Test <sup>c</sup>	Selective Reporting <sup>d</sup>	Data Completeness <sup>e</sup>	Statistical <sup>f</sup>
Bakker et al (2019) <sup>10</sup> ,		1. Not blinded to test groups				

Study	Selection <sup>a</sup>	Blinding <sup>b</sup>	Delivery of Test <sup>c</sup>	Selective Reporting <sup>d</sup>	Data Completeness <sup>e</sup>	Statistical <sup>f</sup>
Berg et al (2012) <sup>11</sup> ,			4. Expertise of evaluators not described.			2. Comparison with other tests not reported.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Selection key: 1. Selection not described; 2. Selection not random or consecutive (i.e., convenience).

<sup>b</sup> Blinding key: 1. Not blinded to results of reference or other comparator tests.

<sup>c</sup> Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

<sup>d</sup> Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>e</sup> Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

<sup>f</sup> Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison with other tests not reported.

### Observational Studies: Following Breast-Conserving Therapy

Two prospective studies have reported on the performance of surveillance breast MRI following BCT.<sup>12,13</sup> Study characteristics are shown in Table 7. Both studies were performed in Korea and it is unclear whether the populations overlapped.

**Table 7. Characteristics of Clinical Validity Studies Assessing Surveillance Breast Magnetic Resonance Imaging After Breast-Conserving Therapy**

Study	Study Population	Design	Reference Standard	Identification of Positive MRI Test	Timing of Tests	Blinding of Assessors	Comment
Kim et al (2017) <sup>13</sup> ,	Women in Korea undergoing surveillance breast MRI following BCT from 2014 to 2016	Prospective observational	<ul style="list-style-type: none"> <li>Pathology for positive results</li> <li>Cancer not confirmed at 1-year surveillance imaging for negative results</li> </ul>	Assessed as BI-RADS category 4 or 5 by 1 radiologist with 10+ years of experience in breast MRI	MRI within 4 wk of screening mammography and breast US	No (readers knew results of prior imaging studies)	<ul style="list-style-type: none"> <li>Funded by Bayer Korea</li> </ul>
Cho et al (2017) <sup>12</sup> ,	Women aged ≤50 years in Korea undergoing	Prospective observational	<ul style="list-style-type: none"> <li>Pathology for positive results</li> <li>Cancer not</li> </ul>	Assessed as BI-RADS category 3+ by 1 radiologist with 5+	MRI within 2 mo of screening mammography and breast US	Yes	<ul style="list-style-type: none"> <li>Funded by Bayer Korea</li> </ul>

Study	Study Population	Design	Reference Standard	Identification of Positive MRI Test	Timing of Tests	Blinding of Assessors	Comment
	surveillance breast MRI following BCT from 2010 to 2016		confirmed at 1-year surveillance imaging for negative results	years of experience in breast MRI			<ul style="list-style-type: none"> <li>Overlap with Kim (2017) unclear</li> </ul>

BCT: breast-conserving therapy; BI-RADS: Breast Imaging Reporting and Data System; MRI: magnetic resonance imaging; US: ultrasound.

Results of the clinical validity studies for surveillance of breast MRI following BCT are shown in Table 8. The sensitivity of MRI was higher than mammography and ultrasound with overlapping CIs in both studies. Specificity of MRI was lower than mammography and ultrasound. The combination of mammography and MRI was 100% sensitive and 87% specific. The review by Cho et al (2017) reported that the recall rate was significantly higher for mammography plus MRI (13.8%; 95% CI, 12.0% to 15.5%) compared with mammography alone (4.4%; 95% CI, 3.3% to 5.5%), as was the biopsy rate (2.7% [95% CI, 2.0% to 3.4%] vs. 0.5 [95% CI, 0.2% to 0.8%]). The yield per 1000 examinations was 8.2 (95% CI, 4.3 to 12.2) for mammography plus MRI versus 4.4 (95% CI, 1.5 to 7.2) for mammography.<sup>12,</sup>

**Table 8. Results of Clinical Validity Studies Assessing Surveillance Breast Magnetic Resonance Imaging After Breast-Conserving Therapy**

Study	Initial N	Final N	Excluded Images	Recurrence Rate, %	Clinical Validity (95% CI), %			
					Sensitivity	Specificity	PPV	NPV
Kim et al (2017) <sup>13,</sup>	421 women (429 breast MRIs)	414 women (422 breast MRIs)	Initial diagnosis of malignant phyllodes tumor, lobular carcinoma in situ (n=6), or developed supraclavicular lymph node metastasis within 12 mo (n=1)	2.6				
MRI					82 (48 to 98)	95 (92 to 97)	31 (15 to 51)	99 (98 to 100)
US					18 (2 to 52)	98 (96 to 99)	20 (3 to 56)	98 (96 to 99)

Study	Initial N	Final N	Excluded Images	Recurrence Rate, %	Clinical Validity (95% CI), %			
Mammography					18 (2 to 52)	99 (98 to 100)	40 (5 to 85)	98 (96 to 99)
Cho et al (2017) <sup>12</sup> ,	801	754	Withdrew consent (n=39) or had systemic metastasis (n=7); unclear (n=1)	2.3				
MRI					88 (66 to 97)	90 (88 to 91)	24 (14 to 37)	NR
US					65 (41 to 83)	90 (89 to 92)	35 (19 to 55)	NR
Mammography					53 (31 to 74)	96 (95 to 97)	73 (43 to 90)	NR
Mammography plus MRI					100 (82 to 100)	87 (85 to 89)	29 (18 to 42)	NR

CI: confidence interval; MRI: magnetic resonance imaging; NPV: negative predictive value; NR: not reported; PPV: positive predictive value; US: ultrasound.

Tables 9 and 10 display notable limitations identified in each study.

**Table 9. Study Relevance Limitations of Clinical Validity Studies of Surveillance Breast Magnetic Resonance Imaging After Breast-Conserving Therapy**

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Duration of Follow-Up <sup>e</sup>
Kim et al (2017) <sup>13</sup> ,				1. Health outcomes not reported	
Cho et al (2017) <sup>12</sup> ,				1. Health outcomes not reported	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

<sup>b</sup> Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest.

<sup>c</sup> Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

<sup>d</sup> Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity, and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

<sup>e</sup> Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true-positives, true-negatives, false-positives, false-negatives cannot be determined).

**Table 10. Study Design and Conduct Limitations of Clinical Validity Studies of Surveillance Breast Magnetic Resonance Imaging After Breast-Conserving Therapy**

Study	Selection <sup>a</sup>	Blinding <sup>b</sup>	Delivery of Test <sup>c</sup>	Selective Reporting <sup>d</sup>	Data Completeness <sup>e</sup>	Statistical <sup>f</sup>
Kim et al (2017) <sup>13</sup> ,		1. Not blinded to results of mammography, US, or PET/CT				
Cho et al (2017) <sup>12</sup> ,						

CT: computed tomography; PET: positron emission tomography; US: ultrasound.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Selection key: 1. Selection not described; 2. Selection not random or consecutive (i.e., convenience).

<sup>b</sup> Blinding key: 1. Not blinded to results of reference or other comparator tests.

<sup>c</sup> Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

<sup>d</sup> Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>e</sup> Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

<sup>f</sup> Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison with other tests not reported.

### Section Summary: Screening When Breast Characteristics Limit the Sensitivity of Mammography

The RCT from the Netherlands (Bakker 2019) found that among women with dense breasts, the use of MRI increased the cancer detection rate and decreased the interval cancer rate compared to mammography. However, the false positive rate was 79.8 per 1000 screenings. The trial is continuing in order to assess the effects over time of adjunctive screening with MRI. The prospective cohort trial by the American College of Radiology Imaging Network (ACRIN 6666; Berg 2012) found that the addition of MRI resulted in high cancer detection, but with increased false positive findings. The evidence is insufficient to show that the use of adjunctive MRI to screen average risk individuals who have dense breasts improves the net health outcome.

Two studies assessed the addition of MRI to mammography for surveillance of women who had been treated for cancer with BCT. The sensitivity of adjunct MRI was greater than mammography alone, but with overlapping confidence intervals. The companion study of women under 50 years showed higher cancer detection rates with adjunct MRI but lower specificity than mammography alone; the authors suggested that adjunctive mammography improves detection of early stage but biologically aggressive cancer in the population of younger women. However, to the extent that younger women may constitute a higher risk population, the delineation of MRI for screening high risk individuals is addressed in high risk screening section of this policy. The evidence is insufficient to demonstrate that adjunctive MRI for screening improves the net health outcome when breast characteristics limit the sensitivity of mammography.

## DETECTION USES

### DETECTING SUSPECTED OCCULT BREAST PRIMARY TUMOR WITH AXILLARY NODAL ADENOCARCINOMA WITH A NEGATIVE MAMMOGRAPHY AND PHYSICAL EXAM

**Clinical Context and Test Purpose**

Breast MRI has been advocated to help detect suspected occult primary breast cancer in patients with adenocarcinoma in the axillary lymph nodes after mammography and physical exam have failed to reveal a breast tumor. Localization of a primary breast tumor might permit BCT instead of presumptive mastectomy.

The questions addressed in this portion of the evidence review:

- Does the use of MRI as an adjunct to detect breast cancer eligible for BCT improve the net health outcome compared to standard techniques in individuals with suspected occult breast primary tumor with axillary nodal adenocarcinoma and negative mammography?
- Is this degree of increased accuracy likely to improve net health outcomes via the earlier diagnosis, better patient management decisions, and more appropriate treatment?

The following PICO was used to select literature to inform this review.

***Populations***

The population of interest is individuals with suspected occult breast primary tumor with axillary nodal adenocarcinoma and negative mammography.

***Interventions***

The intervention of interest is MRI examination as an adjunct to detect breast cancer eligible for BCT.

***Comparators***

The comparator of interest is a preemptive mastectomy.

***Outcomes***

The outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility are the avoidance of invasive procedures (eg, biopsy, mastectomy), the ability to detect cancer that would require additional or earlier treatment, and overall mortality and breast cancer-specific mortality rates.

Breast MRI is performed after a positive breast cancer screening or diagnostic examination.

**Study Selection Criteria**

For the evaluation of the clinical validity of MRI as an adjunct to detect breast cancer eligible for BCT, studies that met the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard
- Patient/sample clinical characteristics were described
- Patient/sample selection criteria were described.

**Clinically Valid**

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

**REVIEW OF EVIDENCE**

**Systematic Reviews**

De Besser et al (2010) evaluated 8 retrospective studies in a systematic review of studies on the use of MRI in patients (N=220) with mammographically occult breast cancer and an axillary metastasis.<sup>14</sup> In 7 studies, a potential primary lesion was detected in a mean of 72% of cases (range, 36% to 86%). Pooling individual patient data yielded a sensitivity of 90% (range, 85% to 100%) in detecting an actual malignant tumor. Specificity, however, was 31% (range, 22% to 50%).

**Clinically Useful**

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

**Direct Evidence**

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

Evidence on detection of suspected occult breast cancer is based on a TEC Assessment (2004)<sup>15</sup>, and a subsequent meta-analysis, which appear to be the only direct evidence available for this indication. The Assessment concluded that, in this small subgroup of patients, adjunctive use of breast MRI allowed a substantial portion of patients (25% to 61%) to avoid the morbidity of mastectomy; risk of the unnecessary biopsy was estimated to be 8%.

**Section Summary: Detecting Suspected Occult Breast Primary Tumor With Axillary Nodal Adenocarcinoma With a Negative Mammography and Physical Exam**

The use of MRI to guide BCS rather than presumptive mastectomy appears to offer the substantial benefit of breast conservation for those patients in whom MRI detects the primary tumor.

**DETECTING CONTRALATERAL BREAST CANCER AFTER ESTABLISHED BREAST CANCER****Clinical Context and Test Purpose**

Patients with a diagnosed breast cancer are at higher risk for a synchronous or subsequent breast cancer in the contralateral breast, and breast MRI has been suggested as a more sensitive screening test compared to mammography.

The questions addressed in this portion of the evidence review:

- Does the use of MRI as an adjunct to detect breast cancer in the contralateral breast improve the net health outcome compared to standard techniques in individuals with suspected occult breast primary tumor with axillary nodal adenocarcinoma and negative mammography?
- Is this degree of increased accuracy likely to improve net health outcomes via the earlier diagnosis, better patient management decisions, and more appropriate treatment?

The following PICO was used to select literature to inform this review.

**Populations**

The population of interest is individuals with breast cancer.



**Interventions**

The intervention of interest is MRI examination as an adjunct to detect breast cancer in the contralateral breast.

**Comparators**

The comparator of interest is mammography and clinical assessment alone.

**Outcomes**

The outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility are the avoidance of invasive procedures (eg, biopsy, mastectomy), the ability to detect cancer that would require additional or earlier treatment, and overall mortality and breast cancer-specific mortality rates.

Breast MRI is performed after a positive breast cancer screening or diagnostic examination.

**Study Selection Criteria**

For the evaluation of the clinical validity of MRI examination as an adjunct to detect breast cancer in the contralateral breast, studies that met the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard
- Patient/sample clinical characteristics were described
- Patient/sample selection criteria were described.

**Clinically Valid**

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

**REVIEW OF EVIDENCE****Single Arm Studies**

Lehman et al (2007) reported on the results of the ACRIN-A6667 trial.<sup>16</sup> They found that 30 (3%) of 969 women with a recent diagnosis of unilateral breast cancer had contralateral cancer at the time of initial diagnosis using MRI. Contralateral lesions were not detected by mammography or physical exam. Eighteen (60%) of the 30 cancers were invasive and 12 (40%) were ductal carcinoma in situ (DCIS). In this study, 121 (12.5%) patients had biopsies, with a positive biopsy rate of 24.8%. With 1-year follow-up, the sensitivity of MRI was 91% and specificity was 88%. Results of this trial in a diverse group of patients were similar to the findings of others.

Liberman et al (2003) reported on 212 women who had negative mammograms of the asymptomatic contralateral breast and found 12 cancers (prevalence, 5%) on MRI, including 6 DCIS and 6 infiltrating carcinomas.<sup>17</sup> However, the PPV of these findings was only 20%, with a specificity of 76%. Lehman et al (2005) found 4 contralateral cancers in 103 patients; in this study, 10 biopsies were done.<sup>18</sup>

**Clinically Useful**

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

**Direct Evidence**

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

No RCTs assessing diagnostic breast MRI in individuals with suspected contralateral breast cancer after established breast cancer were identified.

**Chain of Evidence**

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

A trial with nearly 1000 women found that MRI had high sensitivity and reasonably high specificity for identifying contralateral lesions not detected by mammography or physical examination. Although long-term outcomes of contralateral breast cancers are not fully known, important management changes will occur based on such findings, and these management changes should lead to improved outcomes.

**Section Summary: Detecting Contralateral Breast Cancer After Established Breast Cancer**

The available evidence suggests that adjunctive MRI can identify contralateral breast cancers in women with negative mammograms. A trial with nearly 1000 women found that MRI had high sensitivity and reasonably high specificity for identifying contralateral lesions not detected by mammography or physical examination. Although long-term outcomes of contralateral breast cancers are not fully known, important changes in management will occur as a result of the findings, and these management changes should lead to improved outcomes. That is, in addition to the presumed benefits of early detection, simultaneous treatment of synchronous cancers can occur rather than multiple treatments on separate occasions.

**DETECTING BREAST CANCER IN THE CASE OF LOW-SUSPICION FINDINGS ON CONVENTIONAL MAMMOGRAPHY****Clinical Context and Test Purpose**

Patients with abnormal findings on mammography are categorized according to the level of suspicion of the findings. Patients with low-suspicion findings are often recommended to undergo short-interval follow-up after 3 to 6 months (instead of immediate biopsy). This follow-up may continue for 2 years to demonstrate the stability of benign findings or to detect progression; progression would indicate the need for biopsy. Breast MRI has been investigated as a more sensitive technique to further characterize low-suspicion breast lesions, so that patients with MRI-negative lesions may be reassured and avoid prolonged follow-up and those with MRI-positive lesions may be referred for early biopsy, possibly leading to earlier diagnosis and treatment.

The question addressed in this portion of the evidence review is: In patients with a diagnosis of low-suspicion findings on conventional testing not indicated for immediate biopsy, are net health outcomes improved by the use of MRI as an adjunct compared to standard care with short-interval mammographic follow-up to detect breast cancer?

The following PICO was used to select literature to inform this review.

***Populations***

The population of interest is individuals with low-suspicion findings on conventional mammography.

***Interventions***

The intervention of interest is MRI examination as an adjunct to standard care with short-interval mammographic follow-up.

***Comparators***

The comparator of interest is standard care and short-interval mammographic follow-up.

***Outcomes***

The outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility are the avoidance of invasive procedures (eg, biopsy, mastectomy), the ability to detect cancer that would require additional or earlier treatment, and overall mortality and breast cancer-specific mortality rates.

Breast MRI is performed after a positive breast cancer screening or diagnostic examination.

**Study Selection Criteria**

For the evaluation of the clinical validity of MRI examination as an adjunct to standard care with short-interval mammographic follow-up, studies that met the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard
- Patient/sample clinical characteristics were described
- Patient/sample selection criteria were described.

**Clinically Valid**

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

See the Clinically Useful section for discussion.

**Clinically Useful**

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

**REVIEW OF EVIDENCE**

**Direct Evidence**

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs. Currently, there is a lack of direct evidence supporting use.

**Chain of Evidence**

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because the clinical validity of adjunctive MRI has not been established, a chain of evidence supporting the clinical utility of this modality cannot be constructed.

**Section Summary: Detecting Breast Cancer in the Case of Low-Suspicion Findings on Mammography**

Currently, there is a lack of direct evidence supporting use for this indication. Well-designed prospective confirmatory studies would be necessary to permit conclusions about the effect of this adjunctive use of breast MRI on health outcomes.

**DETECTING BREAST CANCER BY FURTHER CHARACTERIZING SUSPICIOUS BREAST LESIONS****Clinical Context and Test Purpose**

Breast lesions detected by clinical exam or mammography that are considered suspicious are frequently referred for biopsy; however, only a minority of such biopsies reveal breast cancer due to the relatively low specificity of clinical and radiologic exams. Breast MRI has been investigated as a technique to further characterize suspicious breast lesions so that patients with benign lesions may be spared a biopsy procedure. One infrequent situation (niche use) in which MRI of the breast may be helpful and improve health outcomes is in the management of patients who have a suspicious lesion that can only be seen on one mammographic view (i.e., the lesion cannot be seen in other views or on an ultrasound). Patients who fall under this category have a lesion that is not palpable, and therefore, percutaneous biopsy localization cannot be performed. Instead, MRI would be used to localize the suspicious lesion and permit biopsy (this technique would presumably lead to earlier diagnosis of breast cancer as opposed to waiting until the lesion was visible on 2 mammographic views or on ultrasound). The previously described scenario is an infrequent occurrence, so the evidence base addressing this use is mainly anecdotal, but the clinical rationale supporting this use is good.

The question addressed in this portion of the evidence review is: Does MRI as an adjunct to further characterize breast lesions lead to better net health outcomes than biopsy based on mammography and clinical assessment?

The following PICO was used to select literature to inform this review.

**Populations**

The population of interest is individuals with suspicious breast lesions.

**Interventions**

The intervention of interest is MRI examination as an adjunct to mammography and clinical assessment.

**Comparators**

The comparator of interest is biopsy based on mammography and clinical assessment.

**Outcomes**

The outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility are the avoidance of invasive procedures (eg, biopsy, mastectomy), the ability to detect cancer that would require additional or earlier treatment, and overall mortality and breast cancer-specific mortality rates.

Use of MRI is performed after a positive breast cancer screening or diagnostic examination.

**Study Selection Criteria**

For the evaluation of the clinical validity of MRI examination as an adjunct to mammography and clinical assessment, studies that met the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard
- Patient/sample clinical characteristics were described
- Patient/sample selection criteria were described.

**Clinically Valid**

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

**REVIEW OF EVIDENCE****Systematic Reviews**

A systematic review published by Medeiros et al (2011) analyzed 69 studies including 9298 women.<sup>19</sup> Pooled sensitivity was 90% (95% CI, 88% to 92%), and pooled specificity was 75% (95% CI, 70% to 79%). The pooled positive likelihood ratio of an abnormal MRI for malignancy was 3.6 (95% CI, 3.0 to 4.2) and the pooled negative likelihood ratio was 0.12 (95% CI, 0.09 to 0.15). For breast cancer or high-risk lesions versus benign lesions, the area under the curve for MRI was 0.91.

A systematic review published by Zhang et al (2022) included 29 studies with 2976 patients and 3365 suspicious breast lesions.<sup>20</sup> The sensitivity and specificity of MRI features in differentiating malignant from benign breast lesions ranged from 73.8% to 91.9% and from 33.9% to 85.4%, respectively. The enrolled studies showed high heterogeneity. For differentiating malignant from benign breast lesions, the area under the curve values of MRI features; irregular shape, noncircumscribed margin, mass enhancement, heterogeneous internal enhancement, and type II or III time intensity curve patterns were 0.79, 0.87, 0.63, 0.82, and 0.89, respectively.

### Single Arm Studies

Two single-institution, prospective cohort studies examined the diagnostic accuracy of breast MRI for lesions identified by mammography or ultrasound. Strobel et al (2015) in Germany included lesions characterized as Breast Imaging Reporting and Data System (BI-RADS) category 4 by conventional workup in 340 women.<sup>21</sup> Most women were postmenopausal (61%), had no previous breast biopsy (64%), or family history of breast cancer (62%), and underwent initial evaluation for routine screening (88%). Of 353 lesions, 135 (38%) were biopsied; lesions downgraded to BI-RADS categories 1, 2, or 3 on MRI were followed with imaging for 18 months, except for pure clustered microcalcifications (without accompanying mass), which were biopsied or followed with imaging for 24 months at patient discretion; none of the lesions monitored progressed during follow-up. The overall incidence of malignancy including DCIS was 20% (n=69). The MRI downgraded 256 (28%) of 353 lesions, confirmed 37 (11%) lesions, and upgraded 50 (14%) lesions. The PPV of MRI was 73% compared with 19% for conventional imaging. The negative predictive value (NPV) of MRI was 99% (and could not be calculated for conventional imaging). For pure clustered microcalcifications, sensitivity was 89% (25/28 lesions) and the false-negative rate was 12% (3/28 lesions). False-positive MRI findings resulted in a biopsy for 5 (1.5%) of 340 women.

In a similar study, Li et al (2014) in China included 84 women with BI-RADS categories 3, 4, or 5 microcalcifications on mammography.<sup>22</sup> Most patients were premenopausal (81%), had no family history of breast cancer (83%), and underwent initial evaluation for routine screening (56%). All lesions were biopsied surgically (n=91). The incidence of malignancy including DCIS was 46%. The PPV of MRI was 87% compared with 60% for mammography. The NPV of the MRI was 91%.

de Oliveira Pereira et al (2020) performed a cross-sectional study in Brazil of 32 women with suspected breast tumor based on findings from mammography, ultrasonography, or MRI.<sup>23</sup> The mean age of patients was 54.6 years, and the mean breast lump size was 1.6 cm. The sensitivity, specificity, PPV, and NPV were 100%, 50%, 66.7%, and 100%, respectively, for MRI; 56.2%, 87.5%, 81.8%, and 66.7% for mammography; and 75%, 18.8%, 48%, and 42.8% for ultrasonography.

### Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

## REVIEW OF EVIDENCE

### Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

No RCTs assessing diagnostic breast MRI in individuals to further characterize suspicious breast lesions were identified.

**Chain of Evidence**

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Available evidence has not shown this use of breast MRI would improve health outcomes. Considering the relative ease of breast biopsy, the sensitivity of breast MRI would have to be virtually 100% to confidently avoid biopsy. Although MRI performs well, it is clear that the sensitivity is not 100%. False-negative results tend to occur, particularly in certain subcategories, such as DCIS, but invasive carcinomas may not be detected on MRI, also leading to false-negative results. The potential harm to health outcomes of failing to diagnose breast cancer or at least of delaying the diagnosis of breast cancer is of significant concern.

**Section Summary: Detecting Breast Cancer by Further Characterizing Suspicious Breast Lesions**

Use of MRI for evaluation of suspicious breast lesions has relatively high sensitivity and a moderately high specificity. However, it has not yet been established whether the NPV is sufficient to preclude the need for biopsy. Although 3 more recent studies have reported NPVs greater than 90% in certain types of breast lesions, these studies were conducted in single, non-U.S. institutions that require replication in larger, multicenter trials. Therefore, the use of MRI to further characterize suspicious lesions is currently unlikely to alter clinical management. In addition, the fairly high rate of false-positives will lead to substantial numbers of unnecessary biopsies.

**Treatment-Related Uses**

Treatment-related uses addressed here are surgical planning, evaluating tumor response to neoadjuvant therapy, and evaluating residual tumor after BCT. Preoperative planning includes identification of multicentric disease in clinically localized breast cancer; surgical decisions after neoadjuvant chemotherapy; evaluation of suspected chest wall involvement; and localizing lesions prior to biopsy.

For each of these indications, study selection prioritized systematic reviews focusing on the relevant patient population and purpose. Systematic reviews were supplemented by studies of clinical validity. For the evaluation of clinical validity of MRI examination for the proposed purpose, studies that met the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard
- Patient/sample clinical characteristics were described
- Patient/sample selection criteria were described.

In addition, we sought studies of clinical usefulness. These are studies that report the outcomes of using MRI for the proposed purpose, with preference for RCTs.

**Objective: Surgical Planning**

The question addressed in this portion of the evidence review is whether the use of MRI evaluation as an adjunct to guide treatment planning (eg, surgical approach) for patients with

known or suspected breast cancer improves the net health outcome compared with standard techniques.

The sections on surgical planning address 4 specific indications (1) identification of multicentric disease in clinically localized breast cancer; (2) surgical decisions after neoadjuvant chemotherapy; (3) evaluation of suspected chest wall involvement; and (4) localizing lesions prior to biopsy.

## **PREOPERATIVE MAPPING TO IDENTIFY MULTICENTRIC DISEASE WITH CLINICALLY LOCALIZED BREAST CANCER**

### **Clinical Context and Test Purpose**

Patients with clinically localized breast cancer are considered candidates for BCS followed by radiotherapy. However, mastectomy may be considered in patients with multicentric disease (in a separate quadrant of the breast). Breast MRI has been investigated as a technique to assess the extent of the tumor in the breast, specifically to detect multicentric disease as an aid to surgical planning.

The question addressed in this evidence review is: Does the use of MRI for preoperative mapping to identify multicentric disease improve net health outcomes?

The following PICO was used to select literature to inform this review.

### ***Populations***

The populations of interest is individuals with clinically localized breast cancer.

### ***Interventions***

The intervention of interest is MRI as an adjunct to standard evaluation methods.

### ***Comparators***

The following tests and practices are currently being used to make decisions about managing breast cancer: standard workup without MRI.

### ***Outcomes***

Relevant outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility include avoidance of invasive procedures (eg, biopsy, mastectomy), the ability to detect cancer requiring additional or earlier treatment, and overall mortality and breast cancer-specific mortality rates.

Breast MRI is performed after identification of suspicious breast lesions, or before or after treatment for breast cancer.

### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.



- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer duration were preferred.
- Studies with duplicative or overlapping populations were excluded.

## REVIEW OF EVIDENCE

### Systematic Reviews

Several meta-analyses have evaluated evidence on additional disease detected by MRI and changes in clinical management, most of which were by the same research group.<sup>24,25,26,27,28,29,</sup>

Li et al (2022) conducted a systematic review of 19 studies (4 RCTs, 15 observational) that evaluated the efficacy of preoperative MRI in patients with invasive breast cancer.<sup>29</sup> All breast cancer types were included but patients had to be undergoing curative surgery (eg, excision or BCS). All studies included a control group. The primary outcome, mastectomy rate, was significantly increased with preoperative MRI (odds ratio [OR], 1.36; 95% CI, 1.13 to 1.64;  $p=.001$ ;  $I^2=91\%$ ) based on data from 16 studies ( $n=86,075$ ). Preoperative MRI significantly reduced the rate of reoperation (OR, 0.77; 95% CI, 0.62 to 0.97;  $p=.02$ ;  $I^2=71\%$ ). Other outcomes, including primary BCS, secondary mastectomy, and the rate of positive margins, were not significantly different between groups. An analysis of 3 studies in patients with invasive lobular carcinoma found similar results for all outcomes among patients who did and did not receive preoperative MRI.

The most recent meta-analysis published by Houssami et al was in 2017.<sup>26</sup> Studies included in the review were comparative (randomized or nonrandomized), evaluated preoperative MRI versus an alternative approach that did not include MRI, and reported quantitative data on surgical outcomes. The primary endpoint for the meta-analysis was whether patients underwent mastectomy as surgical treatment. Secondary endpoints were re-excision rates after BCS, positive margins after BCS, and receipt of contralateral prophylactic mastectomy. Nineteen studies met the inclusion criteria-3 RCTs and 16 nonrandomized comparative studies. For the primary study endpoint, a pooled analysis of 15 studies ( $N=85,975$ ) found significantly greater odds of receiving a mastectomy after preoperative MRI than after no MRI OR, 1.39; 95% CI, 1.23 to 1.57;  $p<.001$ ). Findings were the same in analyses stratified by publication dates, suggesting that the higher mastectomy rates were not limited to older studies conducted when the MRI-guided biopsy was less common. In an analysis limited to patients with invasive lobular cancer, there was no significant difference in the odds of mastectomy (6 studies: pooled OR, 1.00; 95% CI, 0.75 to 1.33;  $p=.988$ ) or the odds of re-excision (5 studies: OR, 0.65; 95% CI, 0.35 to 1.24;  $p=.192$ ). Among the secondary outcomes, a pooled analysis of 3 studies found a significantly higher odds of contralateral prophylactic mastectomy after MRI (OR, 1.91; 95% CI, 1.25 to 2.91). There were no significant differences between groups on other secondary outcomes (i.e., re-excision rates, positive margins, reoperation rates).

One meta-analysis has addressed breast cancer recurrence rates. This meta-analysis, by Houssami et al (2014), analyzed individual patient data from 4 studies-1 RCT and 3 nonrandomized comparative studies ( $N=3180$ ).<sup>28</sup> Most patients (62% to 93%) had localized, invasive disease and received BCT and systemic chemotherapy. After a median follow-up of 2.9 years (interquartile range [IQR], 1.6 to 4.5 years), there was no difference in estimated 8-year

ipsilateral local (adjusted hazard ratio [HR], 0.88; 95% CI, 0.52 to 1.51;  $p=.65$ ) or distant (adjusted HR, 1.18; 95% CI, 0.76 to 2.27;  $p=.48$ ) recurrence-free survival overall or in patients who received BCT only.

### Randomized Controlled Trials

Since the publication of the Houssami et al (2017) meta-analysis, Bruck et al (2018) reported on the results of an RCT to evaluate the diagnostic value of preoperative MRI in 100 patients with newly diagnosed unifocal stage I invasive ductal carcinoma.<sup>30</sup> Patients were randomized in a 1:1 ratio to preoperative breast MRI or surgery without MRI. Breast MRI detected an additional finding in 14 patients (28%) and MRI detected lesions in 7 (14%) patients, that were confirmed to be malignant. Seven (14%) patients underwent breast reoperation in the MRI group compared with 12 (24%) patients in the control group ( $p=.20$ ). Definitive mastectomy was performed in 6 (12%) patients in the MRI group compared with 2 (4%) in the control group ( $p=.14$ ).

Mota et al (2023) conducted a single-center, open-label RCT (BREAST-MRI) in patients with breast cancer undergoing breast conserving surgery.<sup>31</sup> Two hundred fifty seven patients received preoperative MRI and 267 patients served as controls. Local relapse-free survival ( $p=.7$ ), overall survival ( $p=.8$ ), and reoperation rates ( $p=.85$ ) were similar between groups; however, 21 patients underwent mastectomy in the MRI group compared to 1 patient in the control group.

A discussion of the 3 RCTs included in the Houssami et al (2017) meta-analysis (described above) is as follows.

The RCT by Gonzalez et al (2014) in Sweden assessed 440 women who underwent surgical treatment of invasive breast cancer with or without presurgical breast MRI.<sup>32</sup> Breast MRI provided incremental information that altered the treatment plan in 40 (18%) of 220 patients in the MRI group. Conversion from planned BCS to mastectomy occurred more often in the MRI group (20%) than in the control group (10%;  $p=.024$ ). However, more patients in the MRI group had planned BCS at baseline (70%) than in the control group (60%;  $p=.036$ ). The ipsilateral reoperation rate was 5% in the MRI group versus 15% in the control group ( $p<.001$ ). Reoperation rates among those initially planned for BCS were 5% and 22%, respectively ( $p<.001$ ).

A second RCT, the preoperative MRI and surgical management in patients with nonpalpable breast cancer trial, was reported by Peters et al (2011).<sup>33</sup> It randomized 463 patients with suspicious, nonpalpable breast lesions identified by mammography or ultrasound to prebiopsy MRI or usual care. Of 207 evaluable patients in the MRI group, 11 additional suspicious lesions were identified on MRI and were occult on other imaging studies. All 11 additional lesions underwent biopsy, with 2 (18%) positive for malignancy. The incidence of mastectomy was similar between groups (32% vs. 34% ;  $p=.776$ ), as was the incidence of BCS (68% vs. 66%). The incidence of re-excisions due to positive tumor margins was significantly greater in the MRI group (34%) than in the control group (12%;  $p=.008$ ).

A multicenter RCT from the U.K., Comparative effectiveness of MRI in breast cancer trial, reported by Turnbull et al (2010), examined the impact of presurgical MRI on the need for additional treatment within 6 months.<sup>34</sup> This study was an open, parallel-group trial conducted at 45 centers in the U.K. and enrolled 1623 women with biopsy-proven breast cancer who were scheduled for wide local excision BCT. Of 816 patients in the MRI group, 58 (7%) underwent

mastectomy as a result of MRI findings and/or patient choice, compared with 10 (1%) patients in the no-MRI group who underwent mastectomy by patient choice. There was no statistically significant reduction in reoperation rates in those who received MRI scans (19% in both groups; OR, 0.96; 95% CI, 0.75 to 1.24;  $p=.77$ ). In the MRI group, 19 (2%) patients had a "pathologically avoidable" mastectomy, defined as a mastectomy based on MRI results showing more extensive disease but histopathology showing only localized disease. Twelve months after surgery, there was no statistically significant difference in the quality of life between groups.

### Observational Studies

In addition to the RCTs, Onega et al (2018) reported on the association between preoperative MRI and all-cause mortality in 5 registries (N=4454) of the National Cancer Institute-sponsored Breast Cancer Surveillance Consortium.<sup>35</sup> Data from the Breast Cancer Surveillance Consortium registries were linked to Medicare claims data or electronic health records; women ages 66 years and older with initial nonmetastatic breast cancer (stage I to III) diagnosed from 2005 to 2010 were included with follow-up continuing through 2014. Nine hundred seventeen (21%) women underwent preoperative MRI. The unadjusted 5-year cumulative probability of death was 0.12 for women with MRI and 0.17 for those without (HR, 0.67; 95% CI, 0.54 to 0.82). However, after adjustment for age, sociodemographic, and clinical factors, the association was attenuated (HR, 0.90; 95% CI, 0.72 to 1.12).

Fortune-Greeley et al (2014) retrospectively examined case records of 20,332 women with invasive breast cancer in the Surveillance Epidemiology and End Results-Medicare-linked dataset.<sup>36</sup> Twelve percent of patients had a preoperative MRI. Among patients with invasive lobular carcinoma, but no other histologic types, preoperative breast MRI was associated with lower odds of reoperation after initial partial mastectomy (adjusted OR, 0.59; 95% CI, 0.40 to 0.86).

Zeng et al (2020) performed a retrospective analysis of 512 women age  $\leq 50$  years undergoing BCT.<sup>37</sup> Preoperative MRI was performed in 64.5% of women. In patients who did versus did not receive preoperative MRI, mean age was 43.4 and 43.6 years, and tumor size was 1.64 and 1.80 cm, respectively. In those who received MRI versus no MRI, local recurrence occurred in 7.9% versus 8.2% of patients, respectively (adjusted HR with MRI vs. no MRI, 1.03; 95% CI, 0.53 to 1.99), and was associated with distant recurrence in 6.4% versus 6.6% of patients (adjusted HR with MRI vs. no MRI, 0.89; 95% CI, 0.43 to 1.84).

### Section Summary: Preoperative Mapping to Identify Multicentric Disease With Clinically Localized Breast Cancer

Preoperative MRI as an adjunct to mammography and clinical assessment identifies additional foci of ipsilateral breast cancer and results in a higher rate of mastectomy. For example, a 2017 meta-analysis of 17 studies found significantly higher odds of receiving a mastectomy after preoperative MRI versus no MRI in women with breast cancer. Follow-up studies have reported mixed results, including no significant reduction in reoperation rates after MRI while other studies have reported lower odds of reoperation in patients with invasive lobular carcinoma. No significant differences in ipsilateral local or distant recurrence-free survival after MRI-guided treatment were found in meta-analyses. While there is limited evidence that use of MRI to identify multicentric disease improves recurrence free survival or reduces operations in the overall population, benefit might accrue to sub populations, particularly high risk individuals.

## **GUIDING SURGICAL DECISIONS AFTER NEOADJUVANT CHEMOTHERAPY**

### **Clinical Context and Test Purpose**

Patients with locally advanced breast cancer are usually offered neoadjuvant chemotherapy to reduce tumor size and permit BCT. Evaluation of tumor size and extent using conventional techniques (i.e., mammography, clinical examination, ultrasonography) is suboptimal, and breast MRI has been proposed as a means to more accurately determine tumor size for surgical planning. Breast MRI before chemotherapy is used to document tumor location so that the tumor can be optimally evaluated after chemotherapy, especially if the size and degree of contrast enhancement are greatly reduced. Tumors that respond to chemotherapy get smaller and may even disappear; however, the actual reduction in size is a delayed finding, and earlier changes in tumor vascularity have been observed in chemotherapy-responsive tumors. A decline in contrast enhancement on MRI has been noted in tumors relatively early in the course of chemotherapy. This MRI finding as an early predictor of tumor response has been explored as a means to optimize the choice of the chemotherapeutic agent (eg, to alter chemotherapy regimen if the tumor appears unresponsive).

The question addressed in this evidence review is: Does the use of MRI for guiding surgical decisions after neoadjuvant chemotherapy in patients with locally advanced breast cancer improve net health outcomes?

The following PICO was used to select literature to inform this review.

### ***Populations***

The population of interest is individuals with locally advanced breast cancer undergoing neoadjuvant chemotherapy.

### ***Interventions***

The intervention of interest is MRI to guide surgical decisions after neoadjuvant chemotherapy.

### ***Comparators***

The following tests and practices are currently being used to make decisions about managing breast cancer: mammography and clinical assessment.

### ***Outcomes***

Relevant outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility include avoidance of invasive procedures (eg, biopsy, mastectomy), the ability to detect cancer requiring additional or earlier treatment, and overall mortality and breast cancer-specific mortality rates.

Breast MRI is performed after identification of suspicious breast lesions, or before or after treatment for breast cancer.

### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;

- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer duration were preferred.
- Studies with duplicative or overlapping populations were excluded.

## REVIEW OF EVIDENCE

### Systematic Reviews

Compared with conventional methods of evaluating tumor size and extent (i.e., mammography, clinical exam, ultrasound), MRI of the breast provides an estimation of tumor size and extent that is at least as good as or better than that based on alternatives. Drew et al (2001) found MRI to be 100% sensitive and specific for defining residual tumor after chemotherapy.<sup>38</sup> Conversely, mammography achieved 90% sensitivity and 57% specificity (mammography results considered equivocal), and the clinical exam was only 50% sensitive and 86% specific. Similarly, Partridge et al (2002) reported on correlations of residual tumor size by histopathology of 0.89 with MRI and 0.60 with a clinical exam.<sup>39</sup> The MRI results were well-correlated with results of the histopathologic assessment (criterion standard) with correlation coefficients ranging from 0.72 to 0.98; however, MRI is not intended as a replacement for histopathologic assessment.

Marinovich et al (2015) published an individual patient data meta-analysis of agreement between MRI and pathologic tumor size and other evaluation methods after neoadjuvant chemotherapy.<sup>40</sup> To be eligible for inclusion, studies had to evaluate at least 15 patients undergoing neoadjuvant chemotherapy who were evaluated with MRI and at least 1 other test (i.e., mammography, ultrasound, clinical examination) after surgery. Studies also had to report residual tumor size (i.e., longest diameter). Twenty-four studies met inclusion criteria, and individual patient data were available for 8 of these studies (N=300). The pooled mean difference (MD) in size estimates between MRI and pathology (8 studies, n=243) was 0.0 cm (95% CI, -0.1 to 0.2 cm). In 4 studies comparing size estimates of mammography and pathology, the MD was 0.0 cm, but the 95% CI was wider (-0.3 to 0.4 cm). In 5 studies (n=123) reporting on the MD between ultrasound and pathology, the pooled estimate was -0.3 cm (95% CI, -0.6 to 0.1 cm). The largest size variance was for studies (3 studies, n=107) comparing clinical examination with pathology (pooled MD, -0.8 cm; 95% CI, -1.5 to -0.1 cm).

Previously, Lobbes et al (2013) reported on a systematic review of 35 studies (N=2359) reporting on the ability of MRI to predict tumor size after neoadjuvant chemotherapy.<sup>41</sup> Literature was searched to July 2012. Median correlation coefficient was 0.70 (range, 0.21 to 0.98). Variation in size between MRI and pathology ranged from -1.4 to +2.0 cm.

### Section Summary: Guiding Surgical Decisions After Neoadjuvant Chemotherapy

Studies, including a 2015 meta-analysis, have found that MRI results are well-correlated with pathologic assessment for measuring residual tumor size after neoadjuvant chemotherapy and that MRI performed better than conventional methods. Using breast MRI instead of conventional methods to guide surgical decisions regarding BCT versus mastectomy after neoadjuvant chemotherapy would be at least as beneficial and might lead more frequently to appropriate surgical treatment.

## EVALUATING SUSPECTED CHEST WALL INVOLVEMENT

### Clinical Context and Test Purpose

Tumors located near the chest wall may invade the pectoralis major muscle or extend deeper into chest wall tissues. Typically, modified radical mastectomy removes only the fascia of the pectoralis muscle; however, tumor involvement of the muscle would also necessitate the removal of the muscle (or a portion of it). In smaller tumors, it is necessary to determine how closely the tumor abuts the pectoralis muscle and whether it invades the muscle to determine whether there is an adequate margin of normal breast tissue to permit BCT. Breast MRI has been suggested as a means of determining pectoralis muscle/chest wall involvement for surgical planning and to assist in the decision whether to use neoadjuvant chemotherapy.

The question addressed in this evidence review is: Does the use of MRI to diagnose chest wall involvement of posteriorly located breast tumors improve net health outcomes?

The following PICO was used to select literature to inform this review.

### ***Populations***

The population of interest is individuals with posteriorly located breast tumors.

### ***Interventions***

The intervention of interest is MRI to diagnose chest wall involvement.

### ***Comparators***

The following tests and practices are currently being used to make decisions about managing breast cancer: mammography.

### ***Outcomes***

Relevant outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility include avoidance of invasive procedures (eg, biopsy, mastectomy), the ability to detect cancer requiring additional or earlier treatment, and overall mortality and breast cancer-specific mortality rates.

Breast MRI is performed after identification of suspicious breast lesions, or before or after treatment for breast cancer.

### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

## REVIEW OF EVIDENCE

### Observational Studies

Morris et al (2000) prospectively studied 19 patients with posteriorly located breast tumors suspected to involve the pectoralis major muscle based on either mammography or clinical exam.<sup>42</sup> Thirteen tumors were thought to be fixed to the chest wall on clinical exam, and 12 appeared to have pectoral muscle involvement on mammography. The MRI results were compared with surgical and pathologic findings. The presence of abnormal enhancement within the pectoralis major muscle on MRI was 100% sensitive and 100% specific for identifying 5 tumors that actually involved the pectoralis major muscle.

Two other retrospective studies have reported on 4 cases in which MRI was able to determine the involvement of the chest wall with 100% accuracy.<sup>43,44</sup>

### Section Summary: Evaluating Suspected Chest Wall Involvement

Evidence on MRI for evaluating suspected chest wall involvement with posteriorly located tumors is based on prospective and retrospective observational studies. All studies found that MRI was able to detect chest wall involvement with 100% accuracy. Given the high level of diagnostic accuracy for MRI compared with criterion standard and conventional alternative techniques, the evidence is considered sufficient to conclude that breast MRI improves net health outcome.

## EVALUATING AND LOCALIZING LESIONS PRIOR TO BIOPSY

### Clinical Context and Test Purpose

The question addressed in this evidence review is: Does the use of MRI to evaluate and localize breast lesions prior to biopsy improve net health outcomes?

The following PICO's were used to select literature to inform this review.

### Populations

The populations of interest is individuals with a suspicious breast lesion recommended for biopsy but not localizable by mammography or ultrasonography.

### Interventions

The intervention of interest is MRI to evaluate and localize breast lesion prior to biopsy.

### Comparators

The following tests and practices are currently being used to make decisions about managing breast cancer: waiting until lesion becomes palpable or visible on mammography or ultrasonography.

### Outcomes

Relevant outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility include avoidance of invasive procedures (eg, biopsy, mastectomy), the ability to detect cancer requiring additional or earlier treatment, and overall mortality and breast cancer-specific mortality rates.

Breast MRI is performed after identification of suspicious breast lesions recommended for biopsy.

### Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer duration were preferred.
- Studies with duplicative or overlapping populations were excluded.

## REVIEW OF EVIDENCE

### Observational Studies

Use of MRI to evaluate lesions prior to biopsy is infrequent. An MRI is used in this situation to permit biopsy and breast cancer diagnosis sooner than waiting until the lesion is visible on 2 mammographic views or on ultrasound or becomes palpable. The evidence base addressing this use is mainly anecdotal.

Xie et al (2023) retrospectively evaluated the value of breast MRI to downgrade suspicious lesions (BI-RADS 4A or 4B) found on ultrasound in 167 patients with 186 lesions.<sup>45</sup> Compared to pathology and imaging findings over the subsequent 12 months, MRI had 100% sensitivity, 92.6% specificity, 87.8% PPV, and 100% NPV. Four additional suspicious lesions were detected by MRI, of which 3 (75%) were malignant. Survival was not mentioned. The authors concluded that MRI could allow suspicious lesions to be downgraded and prevent unneeded biopsies.

De Lima Docema et al (2014) used contrast-enhanced MRI to locate occult tumors in 25 patients selected from a group who had undergone breast MRI for suspicious incidental MRI findings at a single-institution in Brazil.<sup>46</sup> Sentinel lymph node mapping and tumor resection were done simultaneously. Malignant tumors were confirmed in 15 (60%) patients, including 4 patients with DCIS. Survival outcomes were not reported.

### Section Summary: Evaluating and Localizing Lesions Prior to Biopsy

A small cohort study in Brazil identified malignant tumors in 60% of patients with MRI-detected occult lesions using contrast-enhanced MRI. A retrospective study of patients with suspicious lesions on ultrasound reported high sensitivity, specificity, PPV, and NPV of MRI to downgrade lesion status and prevent biopsies.

## EVALUATING RESPONSE TO NEOADJUVANT CHEMOTHERAPY WITH LOCALLY ADVANCED BREAST CANCER

### Clinical Context and Test Purpose

The question addressed in this evidence review is: Does the use of MRI to evaluate response to chemotherapy for locally advanced breast cancer improve net health outcomes?

The following PICO was used to select literature to inform this review.



**Populations**

The population of interest is individuals with locally advanced breast cancer undergoing neoadjuvant chemotherapy.

**Interventions**

The intervention of interest is MRI to evaluate the response to chemotherapy.

**Comparators**

The comparator of interest is clinical assessment alone.

**Outcomes**

Relevant outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility include avoidance of invasive procedures (eg, biopsy, mastectomy), the ability to detect cancer requiring additional or earlier treatment, and overall mortality and breast cancer-specific mortality rates.

Breast MRI is performed after a period of undergoing adjuvant chemotherapy.

**Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer duration were preferred.
- Studies with duplicative or overlapping populations were excluded.

**REVIEW OF EVIDENCE****Systematic Reviews**

Four systematic reviews of MRI to evaluate response to neoadjuvant chemotherapy have been published.<sup>41,47,48,49</sup> Characteristics of the reviews are shown in Table 11 and described briefly in the following paragraphs. Li et al (2018) compared the performance of MRI with positron emission tomography (PET) plus computed tomography (CT).<sup>48</sup>

**Table 11. Characteristics of Systematic Reviews Assessing Magnetic Resonance Imaging to Evaluate Response to Neoadjuvant Chemotherapy**

Study	Dates	Studies	Participants	N (Range)	Design	Reference Standard
Janssen et al (2022) <sup>49</sup> ,	2000 to 2019	26	Patients with early-stage breast cancer who received MRI after NAC	4497 (NR)	Observational (prospective, retrospective)	Pathologic response
Li et al (2018) <sup>48</sup> ,	Up to 2017	13	Had both PET/CT and MRI after preoperative NAC with at least 10 patients	MRI: 575 (16 to 142); PET/CT: 618 (16 to 142)	Observational (prospective, retrospective)	Postoperative pathologic result (pCR vs. non-pCR)
Marinovich et al (2013) <sup>47</sup> ,	Up to 2011	44	Newly diagnosed breast cancer undergoing NAC, with MRI undertaken after NAC	2949 (14 to 869)	Observational (prospective, retrospective)	Pathologic response based on surgical excision preferred; other references standards allowed
Lobbes et al (2013) <sup>41</sup> ,	Up to 2012	8	Newly diagnosed breast cancer for whom breast MRI was not performed at baseline or prior to surgery but after completion of NAC with at least 25 patients	560 (31 to 195)	Observational (prospective, retrospective)	NR

CT: computed tomography; MRI: magnetic resonance imaging; NAC: neoadjuvant chemotherapy; NR: not reported; pCR: pathologic complete response; PET: positron emission tomography.

Results of the systematic reviews are shown in Table 12. Janssen et al (2022) reported the results of a systematic review that evaluated the accuracy of MRI for detecting pCR after neoadjuvant chemotherapy.<sup>49</sup> Risk of bias was assessed using the QUADAS-2 tool. Sensitivity was highest for hormone receptor (HR)-negative/*HER2*-negative cancer (0.67), followed by HR-negative/*HER2*-positive (0.65), HR-positive/*HER2*-positive (0.60), and HR-positive/*HER2*-negative (0.55). None of the differences in sensitivity were significant between groups. Specificity results were 0.85, 0.81, 0.74, and 0.88, respectively. Specificity was significantly different between the HR-negative/*HER2*-positive and R-positive/*HER2*-negative groups ( $p=.046$ ).

Li et al (2018) reported on a systematic review comparing MRI with PET/CT to evaluate pathologic response to neoadjuvant chemotherapy and included studies in which patients underwent both PET/CT and MRI after preoperative neoadjuvant chemotherapy; postoperative pathologic complete response (pCR vs. non-pCR) was used as the reference standard; and the study included at least 10 patients.<sup>48</sup> Methodologic quality was assessed using QUADAS-2. Most domains were rated as low-risk of bias in all studies; however, only 2 studies enrolled consecutive or random samples and in only 3 studies were the reference standard results

interpreted without knowledge of the results of the index tests. There was a high level of heterogeneity in the pooled estimate of both sensitivity (88%; 95% CI, 78 to 94;  $I^2=83\%$ ) and specificity (69%; 95% CI, 51 to 83;  $I^2=72\%$ ) for MRI.

Marinovich et al (2013) conducted a systematic review with meta-analysis.<sup>47</sup> Forty-four studies (N=2949) assessing the ability of MRI to discriminate residual breast tumor after neoadjuvant chemotherapy from pCR were identified. Studies were heterogeneous in MRI parameters used, thresholds for identifying a response, and definitions of pathologic response. Median MRI sensitivity, defined as the proportion of patients with residual tumor correctly classified by MRI, and specificity, defined as the proportion of patients with pCR classified by MRI as the absence of residual tumor was 0.92 (IQR, 0.85 to 0.97) and 0.60 (IQR, 0.39 to 0.96), respectively. Specificity increased when a *relative* threshold for defining negative MRI (i.e., contrast enhancement was less than or equal to normal breast tissue) was used rather than an *absolute* threshold (complete absence of MRI enhancement) with little decrement to sensitivity. The pooled area under the receiver operating characteristic curve was 0.88, and the diagnostic OR was 17.9 (95% CI, 11.5 to 28.0). A diagnostic OR of 1 indicates no discriminatory ability; higher values indicate better test performance. Accuracy decreased when residual DCIS was included in the definition of pCR. Statistical measures of between-study heterogeneity were not reported. A subset of studies compared MRI with other imaging modalities (mammography, ultrasound) and clinical exam; however, 95% CIs for pooled analyses were very large, rendering conclusions uncertain.

In the systematic review by Lobbes et al (2013), 8 studies reported on measures of diagnostic accuracy.<sup>41</sup> Median sensitivity, defined as the proportion of patients with pCR correctly classified by MRI, was 42% (range, 25% to 92%). Median specificity, defined as the proportion of patients without pCR correctly classified by MRI, was 89% (range, 50% to 97%). Median (range) PPV and NPV were 64% (50% to 73%) and 87% (71% to 96%), respectively.

**Table 12. Results of Systematic Reviews Assessing Magnetic Resonance Imaging to Evaluate Response to Neoadjuvant Chemotherapy**

Study	MRI		Mammography		PET/CT	
	Sensitivity, %	Specificity, %	Sensitivity, %	Specificity, %	Sensitivity, %	Specificity, %
Janssen et al (2022) <sup>49</sup>						
HR-/HER2- (n=1646), PE (95% CI)	0.67 (0.58 to 0.74)	0.85 (0.81 to 0.88)	NR	NR	NR	NR
HR-/HER2+ (n=1013), PE (95% CI)	0.65 (0.56 to 0.73)	0.81 (0.74 to 0.86)	NR	NR	NR	NR
HR+/HER2- (n=2273), PE (95% CI)	0.55 (0.45 to 0.64)	0.88 (0.84 to 0.91)	NR	NR	NR	NR

Study	MRI		Mammography		PET/CT	
HR+/HER2+ (n=1144), PE (95% CI)	0.60 (0.50 to 0.70)	0.74 (0.63 to 0.83)	NR	NR	NR	NR
Li et al (2018) <sup>48</sup> ,						
Total N	575	575			618	618
PE (95% CI)	88 (78 to 94)	69 (51 to 83)	NR	NR	77 (58 to 90)	78 (63 to 88)
Marinovich et al (2013) <sup>47</sup> ,						
Total N	2949	2949				
Median (IQR)	92 (85 to 97)	60 (39 to 96)	NR	NR	NR	NR
Lobb et al (2013) <sup>41</sup> ,						
Total N	560	560				
Median (range)	42 (25 to 92)	89 (50 to 97)	NR	NR	NR	NR

CI: confidence interval; CT: computed tomography; HR: hormone receptor; IQR: interquartile range; MRI: magnetic resonance imaging; NR: not reported; PE: pooled estimate; PET: positron emission tomography.

### Trials

The ACRIN 6657/I-SPY trial (2012) enrolled 206 women aged 26 to 68 years with invasive breast cancer 3 cm or larger who were receiving anthracycline-based neoadjuvant chemotherapy, with or without a taxane.<sup>50</sup> Of the patients included in the study, 74.4% were White, 19.2% were Black, 4% were Asian, and 2.4% were more than one race or unknown race; 4.2% of patients were Hispanic or Latino. The MRI was performed at 4 time points: before chemotherapy, after 1 cycle of chemotherapy, between the anthracycline-based regimen and the taxane, and after all chemotherapy but before surgery. Various MRI parameters were evaluated for their ability to predict the pathologic outcome. Results were reported as the difference in the predictive ability for residual cancer burden, a composite pathologic index, between MRI parameters and clinical size predictors at the same time points. The MRI findings were a stronger predictor of pathologic outcomes than clinical assessment, with the largest difference being tumor volume after the first chemotherapy cycle and a difference in the area under the receiver operating characteristic curve of 0.09; the corresponding area under the receiver operating characteristic curve values after the third and fourth MRIs were 0.07 and 0.05. Similar findings were reported for predicting pCR.

### Section Summary: Evaluating Response to Neoadjuvant Chemotherapy With Locally Advanced Breast Cancer

Studies, including systematic reviews, have not found sufficient evidence to determine whether breast MRI can reliably predict lack of response to neoadjuvant chemotherapy. There is a large amount of variability in reported performance characteristics of MRI in published studies, leaving uncertain the true accuracy of MRI for this purpose. Furthermore, evidence would need to show that any resulting change in patient management (eg, discontinuation of chemotherapy or change to a different regimen) would improve outcomes.

## **EVALUATING RESIDUAL TUMOR AFTER LUMPECTOMY OR BREAST CONSERVATION SURGERY**

### **Clinical Context and Test Purpose**

In BCT there is complete removal of the primary tumor along with a rim of normal surrounding tissue. Pathologic assessment of surgical margins is performed on excisional specimens to determine whether the tumor extends to the margins of resection. Surgical specimens are oriented and marked to direct re-excision if margins are shown to contain tumor; however, when the tumor is not grossly visible, the extent of a residual tumor within the breast can only be determined through repeat excision and pathologic assessment. Use of MRI has been proposed to evaluate the presence and extent of the residual tumor as a guide to re-excision when surgical margins are positive for tumor.

The question addressed in this evidence review is: Does the use of MRI to evaluate residual tumor after lumpectomy or BCT improve net health outcomes?

The following PICO was used to select literature to inform this review.

### ***Populations***

The population of interest are individuals with positive surgical margins after lumpectomy or BCT.

### ***Interventions***

The intervention of interest is MRI to evaluate the residual tumor.

### ***Comparators***

The comparator of interest is pathologic inspection.

### ***Outcomes***

Relevant outcomes of interest for diagnostic accuracy include test accuracy and test validity (i.e., sensitivity, specificity). Primary outcomes of interest for clinical utility include avoidance of invasive procedures (eg, biopsy, mastectomy), the ability to detect cancer requiring additional or earlier treatment, and overall mortality and breast cancer-specific mortality rates.

Breast MRI is performed after lumpectomy or BCT.

### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer duration were preferred.
- Studies with duplicative or overlapping populations were excluded.

## **REVIEW OF EVIDENCE**

### Observational Studies

Evidence on evaluating residual tumor includes several observational studies, most of which are retrospective.<sup>51,52,53,54,55,56,57,58,59</sup> Histopathologic examination on re-excision was used as the criterion standard. Three studies were conducted at the same institution and accrued patients during similar time periods, so overlap reporting may exist.<sup>52,54,55</sup> Most of the studies were published before 2005 and are not discussed further. Characteristics of studies published since 2015 are shown in Table 13 and described briefly in the following paragraphs.<sup>56,57</sup>

**Table 13. Characteristics of Clinical Validity Studies Assessing Magnetic Resonance Imaging to Evaluate Residual Tumor After Surgery**

Study	Study Population	Design	Reference Standard	Threshold for Positive Index Test	Timing of Reference and Index Tests	Blinding of Assessors	Comment
Lee et al (2018) <sup>57</sup>	Patients in Taiwan with LCIS who had initial excision from 2011 to 2015; race or ethnicity were not described	Unclear	Histopathology	NR	NR	NR	Few details on study design or conduct provided
Krammer et al (2017) <sup>56</sup>	Women with positive margins after initial surgery for breast cancer from 2004 to 2013; race or ethnicity were not described	Retrospective	Histopathology	<ul style="list-style-type: none"> <li>Read independently by 2 radiologists</li> <li>Criteria for suspected residual disease: asymmetric thickening or nodular enhancement with irregular or spiculated margins or extensive focal non-mass enhancement</li> </ul>	NR	Radiologists had access to other imaging results, when available	

LCIS: lobular carcinoma in situ; NR: not reported.

Results of the clinical validity studies published after 2015 are shown in Table 14. Lee et al (2018) reported on the results of a study comparing breast MRI with ultrasonography for detecting remnant lobular carcinoma in situ lesions after initial excision.<sup>57</sup> Twenty-nine patients with lobular carcinoma in situ were enrolled between 2011 and 2015. Methods are poorly described. Residual lesions were identified by pathology in 12 (41%) cases. The sensitivity of ultrasonography was 58% compared with 83% for breast MRI; precision estimates were not reported. Specificity was 100% for both modalities.

Krammer et al (2017) published a retrospective study evaluating breast MRI to assess residual disease in 175 patients who had been candidates for BCS and had positive surgical margins.<sup>56</sup> The MRIs were read independently by 2 radiologists, both of whom had access to the pathology report from the initial surgery and any prior breast imaging. Pathology findings served as the criterion standard. For reader 1, the sensitivity and specificity of detecting residual disease was 63% and 75%, respectively. For reader 2, sensitivity and specificity were 83% and 64%, respectively. The inter-observer agreement was moderate ( $k=0.56$ ).

**Table 14. Results of Clinical Validity Studies Assessing Magnetic Resonance Imaging to Evaluate Residual Tumor After Surgery**

Study	Initial N	Final N	Excluded Samples	Prevalence of Condition, %	Clinical Validity (95% Confidence Interval), %			
					Sensitivity	Specificity	PPV	NPV
Lee et al (2018) <sup>57</sup>	NR	29	Any invasive focus or other malignancy	41				
MRI					83% (NR)	100% (NR)	NR	NR
Ultrasonography					58% (NR)	100% (NR)	NR	NR
Krammer et al (2017) <sup>56</sup>	180	175	Received chemotherapy prior to postoperative MRI (n=4), poor MRI image quality (n=1)	79				
MRI					73% (NR)	72% (NR)	91% (NR)	45% (NR)

MRI: magnetic resonance imaging; NPV: negative predictive value; NR: not reported; PPV: positive predictive value.

Tables 15 and 16 display notable limitations identified in each study.

**Table 15. Study Relevance Limitations of Clinical Validity Studies of Magnetic Resonance Imaging to Evaluate Residual Tumor After Surgery**

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Duration of Follow-Up <sup>e</sup>
Lee et al (2018) <sup>57</sup> ,	2. Study population is unclear	1,2. No description provided	1. No description provided	1. Health outcomes not reported	
Krammer et al (2017) <sup>56</sup> ,	2. Study population is unclear		3. No comparator	1. Health outcomes not reported	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup>Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

<sup>b</sup>Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest.

<sup>c</sup>Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

<sup>d</sup>Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

<sup>e</sup>Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true-positives, true-negatives, false-positives, false-negatives cannot be determined).

**Table 16. Study Design and Conduct Limitations of Clinical Validity Studies Assessing Magnetic Resonance Imaging to Evaluate Residual Tumor After Surgery**

Study	Selection <sup>a</sup>	Blinding <sup>b</sup>	Delivery of Test <sup>c</sup>	Selective Reporting <sup>d</sup>	Data Completeness <sup>e</sup>	Statistical <sup>f</sup>
Lee et al (2018) <sup>57</sup> ,		1. Not described	1,3,4. Not described			1. No precision estimates provided 2. No statistical comparison to other methods
Krammer et al (2017) <sup>56</sup> ,		1. Not blinded to other imaging results				

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup>Selection key: 1. Selection not described; 2. Selection not random or consecutive (i.e., convenience).

<sup>b</sup>Blinding key: 1. Not blinded to results of reference or other comparator tests.

<sup>c</sup>Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

<sup>d</sup>Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>e</sup>Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

<sup>f</sup>Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison with other tests not reported.



**Section Summary: Evaluating Residual Tumor After Lumpectomy or Breast Conservation Surgery**

The available evidence is not sufficient to permit conclusions whether the use of MRI identifies the presence and/or extent of residual disease after lumpectomy or BCS and before re-excision. Most studies were retrospective, and most reported moderate sensitivity and specificity of MRI for detection of residual disease. One study published after 2015 reported the sensitivity and specificity of MRI to be over 70%. The other study published after 2015 reported a sensitivity of 83% and a specificity of 100% but offered very few details on methods, so study quality cannot be assessed.

**SUPPLEMENTAL INFORMATION**

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

**Practice Guidelines and Position Statements**

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

**National Comprehensive Cancer Network**

Current National Comprehensive Cancer Network (NCCN) guidelines on breast cancer (v. 4.2023),<sup>60</sup> breast cancer screening and diagnosis (v. 1.2023),<sup>61</sup> and genetic assessment of those at high-risk of breast, ovarian, and pancreatic cancer (v. 3.2023)<sup>62</sup>, list the following indications for breast magnetic resonance imaging (MRI)

Screening (as an adjunct to mammography):<sup>61</sup>,

"Recommend Annual MRI Screening

- For individuals with a genetic mutation, or an untested first-degree relative of gene mutation carrier
- For individuals who received thoracic RT [radiation therapy] between the ages of 10 and 30 years
- For individuals with a residual lifetime risk >20% as defined by models that are largely dependent on family history; based on the extent of family history, consider referral for genetic testing.
- Consider annual MRI screening for individuals with ADH [atypical ductal hyperplasia] or lobular neoplasia (LCIS [lobular carcinoma in situ]/ALH [atypical lobular hyperplasia]) and ≥20% lifetime risk

Insufficient Evidence to Recommend for or Against Routine Population-Based MRI Screening:

- Residual lifetime risk 15%-20%, as defined by models that are largely dependent on family history
- Heterogeneously or extremely dense breast on mammography

Recommend Against MRI Screening (Based on Expert Consensus Opinion):

- Individuals at <15% residual lifetime risk"

The NCCN guidelines for breast cancer screening and diagnosis also state that individuals assigned female at birth at "increased risk" of breast cancer include the following groups:<sup>61,</sup>

- " those with a prior history of breast cancer;
- those  $\geq 35$  years of age with a 5-year risk of invasive breast carcinoma  $\geq 1.7\%$  (per the Modified Gail Model);
- those who have a lifetime risk  $>20\%$  based on history of LCIS or ADH/ALH;
- those who have a lifetime risk  $>20\%$  as defined by models that are largely dependent on family history;
- those who received prior thoracic irradiation between the ages of 10 and 30 years
- those with a pedigree suggestive of or with a known genetic predisposition"

The NCCN guidelines for genetic or familial high-risk assessment for breast cancer recommend MRI screening with contrast for patients with *BRCA* pathogenic or likely pathogenic variants starting at age 25 to 29 years or individualized if the family had breast cancer diagnosis before age 30. The guidelines further state that MRI with contrast can be considered for patients with the following genetic variants:<sup>62,</sup>

- *ATM*, *BARD1*, and *CHEK2* starting at age 30 to 35 years
- *CDH1*, *STK11*, and *PALB2*, starting at age 30 years
- *NF1*, from ages 30 to 50 years
- *TP53* pathogenic/likely pathogenic variant who are treated for breast cancer and have not had a bilateral mastectomy, starting at age 20 to 29 years
- *RAD51C* and *RAD51D*, starting at age 40 years
- *PTEN* pathogenic/likely pathogenic variant who are treated for breast cancer and have not had a bilateral mastectomy, starting at age 30 to 35 years or 5 to 10 years before the earliest breast cancer in the family

The NCCN guidelines for genetic or familial high-risk assessment for breast cancer also state there is insufficient evidence for any recommendations for use of breast MRI for patients with the following genetic variants: *BRIP1*, *MLH1*, *MSH2*, *MSH6*, *PMS2*, *EPCAM*, *FANCC*, *MRE11A*, *MUTYH* heterozygotes, *RECQL*, *RAD50*, *RINT1*, *SLX4*, *SMARCA4*, or *XRCC2*.

Guidelines on breast cancer screening and diagnosis make the following recommendations on diagnosis:<sup>61,</sup>

- Optional MRI for women with nipple discharge, no palpable mass, and a Breast Imaging Reporting and Data System (BI-RADS) rating of 1 to 3.
- For patients with skin changes consistent with serious breast disease, consideration of breast MRI is included in the guidelines for those with benign biopsy of skin or nipple following BI-RADS category 1 to 3 assessment. Since a benign skin punch biopsy in a patient with clinical suspicion of inflammatory breast cancer (IBC) does not rule out malignancy, further evaluation is recommended...[and] MRI may be used for suspicious nipple discharge when mammography and ultrasound are not diagnostic.

Guidelines on breast cancer make the following recommendations on pretreatment evaluation with breast MRI:<sup>60,</sup>

- "May be useful in identifying otherwise clinically occult disease in patients presenting with axillary nodal metastases (cT0, cN+), with Paget disease, or with invasive lobular

carcinoma poorly (or inadequately) defined on mammography, ultrasound, or physical examination."

- "May be used for staging evaluation to define extent of cancer or presence of multifocal or multicentric cancer in the ipsilateral breast, or as screening of the contralateral breast cancer at time of initial diagnosis."

Guidelines on breast cancer make the following recommendations related to MRI surrounding treatment:<sup>60</sup>

- "May be helpful for breast cancer evaluation before and after preoperative systemic therapy to define extent of disease, response to treatment, and potential for breast-conservation therapy."
- "False-positive findings on breast MRI are common. Surgical decisions should not be based solely on the MRI findings. Additional tissue sampling of areas of concern identified by breast MRI is recommended."

Guidelines on breast cancer make the following recommendations on MRI related to surveillance:<sup>60</sup>

- "The utility of MRI in follow-up screening of patients with prior breast cancer is undefined. It should generally be considered for: patients with dense breasts treated with breast-conserving surgery and radiation therapy, those diagnosed before the age of 50, and those whose lifetime risk of a second primary breast cancer is >20% based on models largely dependent on family history, such as in those with the risk associated with inherited susceptibility to breast cancer."

### **American Cancer Society**

The American Cancer Society recommendations for the early detection of breast cancer, most recently updated in 2022, has recommended the following on MRI:<sup>63</sup>

"Women who are high risk for breast cancer based on certain factors should get a breast MRI and a mammogram every year, typically starting at age 30. This includes women who:

- Have a lifetime risk of breast cancer of about 20% to 25% or greater, according to risk assessment tools that are based mainly on family history
- Have a known *BRCA1* or *BRCA2* gene mutation (based on having had genetic testing)
- Have a first-degree relative (parent, brother, sister, or child) with a *BRCA1* or *BRCA2* gene mutation, and have not had genetic testing themselves
- Had radiation therapy to the chest when they were between the ages of 10 and 30 years
- Have Li-Fraumeni syndrome, Cowden syndrome, or Bannayan-Riley-Ruvalcaba syndrome, or have first-degree relatives with one of these syndromes

The American Cancer Society recommends against MRI screening for women whose lifetime risk of breast cancer is less than 15%.

There's not enough evidence to make a recommendation for or against yearly MRI screening for women who have a higher lifetime risk based on certain factors, such as:

- Having a personal history of breast cancer, ductal carcinoma in situ (DCIS), lobular carcinoma in situ (LCIS), atypical ductal hyperplasia (ADH), or atypical lobular hyperplasia (ALH)
- Having 'extremely' or 'heterogeneously' dense breasts as seen on a mammogram

If MRI is used, it should be in addition to, not instead of, a screening mammogram. This is because although an MRI is more likely to find cancer than a mammogram, it may still miss some cancers that a mammogram would find.

Most women at high risk should begin screening with MRI and mammograms when they are 30 and continue for as long as they are in good health. But this is a decision that should be made with a woman's health care providers, taking into account her personal circumstances and preferences."

### American College of Radiology

The American College of Radiology has appropriateness criteria for breast cancer screening, which were developed in 2012 and revised in 2017;<sup>64</sup> palpable breast masses<sup>65</sup>, revised in 2022 ; initial workup and surveillance for stage I breast cancer, reviewed in 2019<sup>66</sup>; monitoring response to neoadjuvant therapy, revised 2022 ;<sup>67</sup> transgender breast cancer screening, 2021<sup>68</sup>; and supplemental breast cancer screening based on breast density, 2021<sup>69</sup>, (see Table 17).

**Table 17. Magnetic Resonance Imaging-Related Criteria for Breast Cancer Screening, Diagnosis, and Monitoring Response**

Specific Indications	MRI Rating
High-risk women: women with a <i>BRCA</i> gene variant and their untested first-degree relatives, women with a history of chest irradiation between the ages of 10 and 30 years, women with 20% or greater lifetime risk of breast cancer	Usually appropriate with and without contrast (with mammography)
Intermediate-risk women: women with personal history of breast cancer, lobular neoplasia, atypical ductal hyperplasia, or 15% to 20% lifetime risk of breast cancer	May be appropriate with and without contrast (with mammography)
Average-risk women: women with <15% lifetime risk of breast cancer, breasts not dense	Usually not appropriate with and without contrast
Evaluating palpable breast mass. All indications reviewed	Usually not appropriate with and without contrast
Known breast cancer. Initial determination of tumor size and extent within the breast prior to neoadjuvant chemotherapy.	Usually appropriate without and with contrast
Known breast cancer. Imaging of the breast after initiation or completion of neoadjuvant chemotherapy.	Usually appropriate without and with contrast
Known breast cancer, clinically node-negative. Axillary evaluation prior to neoadjuvant chemotherapy.	Usually not appropriate
Known breast cancer, clinically node-positive. Axillary evaluation prior to neoadjuvant chemotherapy.	May be appropriate without and with contrast
Known breast cancer, clinically node-negative. Axillary evaluation after completion of neoadjuvant chemotherapy, axilla not previously evaluated.	Usually not appropriate
Known breast cancer, clinical suspicion of metastatic disease. Staging or assessment of response to neoadjuvant chemotherapy.	Usually not appropriate

<b>Specific Indications</b>	<b>MRI Rating</b>
Known axillary lymph node-positive breast cancer on prior mammography, ultrasound, or MRI. Axillary evaluation after completion of neoadjuvant chemotherapy, axilla previously evaluated.	Usually not appropriate
Known breast cancer. Axillary imaging suspicious for metastatic disease on mammography, ultrasound, or MRI during initial evaluation.	Usually not appropriate
Surveillance. Rule out local recurrence.	May be appropriate without and with contrast
Transfeminine (male-to-female) patient, 40 years of age or older with past or current hormone use $\geq 5$ years; average risk patient.	Usually not appropriate without and with contrast
Transfeminine (male-to-female) patient, 25 to 30 years of age or older with past or current hormone use $\geq 5$ years; higher-than-average risk.	Usually not appropriate without and with contrast
Transfeminine (male-to-female) patient with no hormone use (or hormone use $< 5$ years) at any age; average-risk patient	Usually not appropriate without and with contrast
Transfeminine (male-to-female) patient, 25 to 30 years of age or older with no hormone use (or hormone use $< 5$ years); higher-than-average risk.	Usually not appropriate without and with contrast
Transmasculine (female-to-male) patient with bilateral mastectomies ("top surgery") at any age and any risk.	Usually not appropriate without and with contrast
Transmasculine (female-to-male) patient with reduction mammoplasty or no chest surgery, 40 years of age or older; average-risk patient (less than 15% lifetime risk of breast cancer).	Usually not appropriate without and with contrast
Transmasculine (female-to-male) patient with reduction mammoplasty or no chest surgery, $\geq 30$ years of age. Intermediate risk (patient with personal history of breast cancer, lobular neoplasia, atypical ductal hyperplasia, or 15% to 20% lifetime risk of breast cancer).	May be appropriate without and with contrast; usually not appropriate without contrast
Transmasculine (female-to-male) patient with reduction mammoplasty or no chest surgery, 25 to 30 years of age or older. High risk (with genetic predisposition to breast cancer or untested patient with a first-degree relative with genetic predisposition to breast cancer, patient with a history of chest irradiation between 10 to 30 years of age, patient with 20% or greater lifetime risk of breast cancer).	Usually appropriate without and with contrast; usually not appropriate without contrast
Average-risk females with nondense breasts	Usually not appropriate without and with contrast
Intermediate-risk females with nondense breasts	Usually not appropriate without and with contrast
High-risk females with nondense breasts	Usually not appropriate without and with contrast
Average-risk females with dense breasts	May be appropriate without and with contrast; usually not appropriate without contrast
Intermediate-risk females with dense breasts	May be appropriate without and with contrast;

Specific Indications	MRI Rating
	usually not appropriate without contrast
High-risk females with dense breasts	Usually appropriate without and with contrast; usually not appropriate without contrast

MRI: magnetic resonance imaging.

The College (2018) issued recommendations for breast cancer screening in women at higher-than-average risk.<sup>70</sup> The recommendations for MRI are as follows:

- "For women with genetics-based increased risk (and their untested first-degree relatives), history of chest radiation, calculated lifetime risk of 20% or more, breast MRI should be performed annually beginning at age 25 to 30."
- "For women with personal histories of breast cancer and dense breast tissue, or those diagnosed before age 50, annual surveillance with breast MRI is recommended."
- "For women with personal histories of breast cancer not included in the above, or with LCIS or atypia on prior biopsy, MRI should be considered, especially if other risk factors are present."

### American Society of Clinical Oncology

The American Society of Clinical Oncology (2006) has published guidelines for follow-up and management after primary treatment of breast cancer.<sup>71</sup> In 2013, the guidelines were updated with a systematic review of the literature through March 2012, and no revisions were made.<sup>72</sup> The guidelines recommended against the use of breast MRI "for routine follow-up in an otherwise asymptomatic patient with no specific findings on clinical examination."<sup>72</sup> Furthermore, "The decision to use breast MRI in high-risk patients should be made on an individual basis depending on the complexity of the clinical scenario."<sup>71</sup>

### International Late Effects of Childhood Cancer Guideline Harmonization Group

The International Late Effects of Childhood Cancer Guideline Harmonization Group from 9 countries (2020) published evidence-based recommendations for breast cancer surveillance in female survivors of childhood, adolescent, and young adult cancer who received chest irradiation before age 30 years and have no genetic predisposition to breast cancer.<sup>73</sup> The guideline recommends to initiate annual breast MRI exams beginning at age 25 or 8 years after radiation. Based on a systematic review of the literature to June 2019, the authors recommended mammography and breast MRI for surveillance (strong recommendation based on high-quality evidence with a low degree of uncertainty). The authors acknowledged that "there are no studies of survivors of [childhood, adolescent, and young adult] cancer that investigated whether early detection by MRI or mammography results in better prognosis." However, the panel concluded that the benefits of initiating early annual mammography and MRI are expected to outweigh the harms.

### U.S. Preventive Services Task Force Recommendations

The U.S. Preventive Services Task Force (2016) updated its recommendations on breast cancer screening. The Task Force concluded the following on breast MRI:<sup>74</sup>

"... the current evidence is insufficient to assess the balance of benefits and harms of adjunctive screening for breast cancer using breast ultrasonography, magnetic resonance imaging, DBT [digital breast tomosynthesis], or other methods in women identified to have dense breasts on an otherwise negative screening mammogram."

These guidelines are currently undergoing an update and updated recommendations may be forthcoming.

### Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review are listed in Table 18.

**Table 18. Summary of Key Trials**

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT05797545	Comparison of Ultrasound and Breast MRI for Breast Cancer Detection Among Women With Dense Breasts and a Personal History of Breast Cancer	1464	May 2028
NCT05704062	Multi-Functional Magnetic Resonance Imaging Modalities for Assessment of Breast Cancer Response to Neoadjuvant Chemotherapy	135	Nov 2026
NCT05825768	Preoperative Magnetic Resonance Imaging to Obtain Adequate Resection Margins (PRIMAR) Trial	440	Aug 2026
NCT01805076	Effect of Preoperative Breast MRI on Surgical Outcomes, Costs and Quality of Life of Women With Breast Cancer	317	Feb 2025
NCT01035112	Magnetic Resonance Imaging of Breast Cancer	445	May 2027
NCT00474604	MRI Evaluation of Breast Tumor Growth and Treatment Response	209	Dec 2025
<i>Unpublished</i>			
NCT01716247	Comparison of Contrast Enhanced Mammography to Breast MRI in Screening Patients at Increased Risk for Breast Cancer	1000	Jun 2018
NCT01929395	A Study to Evaluate the Use of Supine MRI Images in Breast Conserving Surgery	159	Jul 2018

NCT: national clinical trial.

**CODING**

**The following codes for treatment and procedures applicable to this policy are included below for informational purposes. This may not be a comprehensive list of procedure codes applicable to this policy.**

**Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.**

**The code(s) listed below are medically necessary ONLY if the procedure is performed according to the "Policy" section of this document.**

<b>CPT/HCPCS</b>	
77046	Magnetic resonance imaging, breast, without contrast material; unilateral
77047	Magnetic resonance imaging, breast, without contrast material; bilateral
77048	Magnetic resonance imaging, breast, without and with contrast material(s), including computer-aided detection (CAD real-time lesion detection, characterization and pharmacokinetic analysis), when performed; unilateral
77049	Magnetic resonance imaging, breast, without and with contrast material(s), including computer-aided detection (CAD real-time lesion detection, characterization and pharmacokinetic analysis), when performed; bilateral

<b>REVISIONS</b>	
06-10-2004	In "Policy" section added 2, 3, 4, 5, and 6.
04-21-2005	In "Policy" section added, "All of the following policy statements refer to performing MRI of the breast with a breast coil. MRI of the breast without the use of a breast coil, regardless of the clinical indication, is considered investigational."
	In "Policy" section added #5 a, b, c, and d – "MRI breast biopsy".
11-03-2005	In "Policy" section changed the wording (not concept) in #1, 2, 3, and 4.
	In "Policy" section #5 is now the new #11. Deleted the fourth bullet and added a statement at the beginning of the policy to address the breast coil.
	In "Policy" section deleted #6 and 7.
	In "Policy" section added new #5, 6, 7, 8, 9, and 10.
	In "Policy" section added, "Breast MRI is considered experimental/investigational as a screening technique in average risk patients."
12-28-2005 with an effective date of 02-01-2006	In "Documentation" section deleted 'The ordering physician should retain in the patient's medical record, history and physical, examination notes documenting evaluation and management of one of the covered conditions/diagnoses, with relevant clinical signs/symptoms or abnormal laboratory test results, appropriate to one of the covered indications. The patient's clinical record should further indicate changes/alterations in medications prescribed for the treatment of the patient's condition. There must be an attending/treating physician's order for each test documented in the patient's medical/clinical record' at the request of the Associate Medical Director.
01-12-2007 with an effective date of 01-01-2007	In "Coding" section, CPT Codes, deleted 76093 and 76094 and added CPT Codes 77058 and 77059 due to the 2007 CPT changes.



<b>REVISIONS</b>	
12-07-2012	Revision posted to BCBSKS website, December 7, 2012.
	Description section updated.
	In the Policy section:
	<ul style="list-style-type: none"> <li>▪ Revised the following medical policy language: MRI of the breast using scanners equipped with breast coils is medically necessary for the following:               <ol style="list-style-type: none"> <li>1. For evaluation for rupture breast implants when there is breast pain and/or abnormal ultrasound of the breast.</li> <li>2. As a screening technique for breast cancer in women with known BRCA1 or BRCA2 mutation; at high risk of BRCA1 or BRCA2 mutation due to a known presence of the mutation in relatives; or with a pattern of breast cancer history in multiple first-degree relatives, often occurring at a young age and bilaterally, consistent with a high probability of harboring BRCA mutations or other hereditary breast cancer.</li> <li>3. For metastatic adenocarcinoma to an axillary node with unknown primary, negative physical exam, and negative standard mammogram.</li> <li>4. For patients who have dense breast tissue, negative mammograms and a strong family history of breast cancer.</li> <li>5. As a screening technique of the contralateral breast in patients who have breast cancer.</li> <li>6. For presurgical planning in patients with locally advanced breast cancer before and after completion of neoadjuvant chemotherapy to permit tumor localization and characterization.</li> <li>7. To determine the presence of pectoralis muscle or chest wall invasion in patients with posteriorly located tumors.</li> <li>8. To detect local tumor recurrence in individuals with breast cancer who have radiographically dense breasts or old scar from previous breast surgery that compromises the ability of combined mammography and ultrasonography.</li> <li>9. Further evaluation of suspicious clinical findings or imaging results, which remain indeterminate after complete mammographic and sonographic evaluations, combined with a thorough physical examination.</li> <li>10. To detect the extent of residual cancer in the recently post operative breast with positive pathological margins after incomplete lumpectomy when the member still desires breast conservation and local re-excision is planned.</li> <li>11. MRI breast biopsy:                   <ol style="list-style-type: none"> <li>a. May be performed if a suspicious lesion is identified only on MRI of the breast.</li> <li>b. Performed by a provider capable of interpreting breast MRI, performing needle biopsy of the breast, and interpreting mammographies.</li> <li>c. Requires only one person to perform a MRI breast biopsy.</li> </ol> </li> </ol> </li> </ul>
	Breast MRI is considered experimental/investigational as a screening technique in average risk patients.
	<ul style="list-style-type: none"> <li>▪ Added Item B, #2, "To Confirm the clinical diagnosis of rupture of silicone breast implants."</li> <li>▪ Added Item C, #7, "To monitor the integrity of silicone gel-filled breast implants when there are no signs or symptoms or rupture."</li> </ul>
	Policy Guidelines section added.
	Rationale section updated.
	Reference section updated.
09-12-2013	Updated Description section.
	In Policy section:

<b>REVISIONS</b>	
	<ul style="list-style-type: none"> <li>For clarification the following statement was revised from "All of the following policy statements refer to performing MRI of the breast with a breast coil. MRI of the breast without the use of breast coil, regardless of the clinical indication is considered experimental / investigational." to read "All of the policy statement above refer to performing MRI of the breast with a breast coil and the use of contrast. MRI of the breast without the use of a breast coil, regardless of the clinical indications, is considered experimental / investigational."</li> </ul>
	Updated Rationale section.
	In Coding section:
	<ul style="list-style-type: none"> <li>Added ICD-10 Diagnosis codes (<i>Effective October 1, 2014</i>)</li> </ul>
	Updated Reference section.
07-08-2015	In Policy title:
	<ul style="list-style-type: none"> <li>Revised from "Magnetic Resonance Imaging (MRI) Breast"</li> </ul>
	Updated Description section.
	In Policy section:
	<ul style="list-style-type: none"> <li>In Policy Guidelines, Item 2, removed "models" and "using family history, including" and added "risk assessment tools based mainly on family history", "and include", and "Cuzick", to read "A number of risk assessment tools based mainly on family history can assist practitioners in estimating breast cancer risk and include the Claus,(1) modified Gail,(2) Tyrer-Cuzick,(3) and BRCAPRO(4) models."</li> <li>In Policy Guidelines, Item 4, added "the use of contrast by" to read, "As noted, breast MRI exams require a dedicated breast coil and the use of contrast by radiologists familiar with the optimal timing sequences and other technical aspects of image interpretation."</li> <li>In Policy Guidelines, Item 5, removed "The use of" and "treatment" and added "apparently", "therapy", to read, "Preoperative MRI in patients with localized disease apparently results in higher rates of mastectomy and lower rates of breast-conserving therapy (BCT)." Also added, "If biopsies are performed on all MRI-identified lesions, and if shared patient decision making is used for altering the surgical approach, then the probability of improved outcomes is increased."</li> </ul>
	Updated Rationale section.
	Updated References section.
10-01-2016	In Coding section:
	<ul style="list-style-type: none"> <li>Added ICD-10 codes effective 10-01-2016: T85.848A, T85.848D, T85.848S, T85.898A, T85.898D, T85.898S</li> <li>Termed ICD-10 codes effective 09-30-2016: T85.84XA, T85.84XD, T85.84XS, T85.89XA, T85.89XD, T85.89XS</li> </ul>
11-09-2016	Updated Description section.
	Updated Rationale section.
	Updated References section.
10-01-2017	In Coding section:
	<ul style="list-style-type: none"> <li>Added ICD-10 codes: N63.11, N63.12, N63.13, N63.14, N63.21, N63.22, N63.23, N63.24, N63.31, N63.32, N63.41, N63.42.</li> <li>Removed ICD-10 code: N63.</li> </ul>
11-08-2017	Updated Description section.
	In Policy section:
	<ul style="list-style-type: none"> <li>In Item A 1, removed "mutation" and added "variant" to read, "With a known <i>BRCA1</i> or <i>BRCA2</i> variant;"</li> <li>In Item A 2, removed "mutation" and added "variant" to read, "At high risk of <i>BRCA1</i> or <i>BRCA2</i> variant due to a known presence of the variant in relatives;"</li> </ul>

<b>REVISIONS</b>	
	Updated Rationale section.
	In Coding section: ▪ Removed ICD-9 codes.
	Updated References section.
01-01-2019	Updated Description section.
	Updated Rationale section.
	In Coding section: ▪ Added new CPT codes: 77046, 77047, 77048, 77049. ▪ Removed deleted CPT codes: 77058, 77059.
	Updated References section.
05-22-2020	Updated Description section.
	Objective section: Changed the word "policy" to "evidence"
	Updated Rationale section.
	In Coding section: ▪ Added: Z15.01 Genetic susceptibility to malignant neoplasm of breast
	Updated References Section
11-05-2021	Updated Description Section
	Updated Rationale Section
	Updated Reference Section
11-22-2022	Updated Description Section
	Updated Policy Section ▪ Section A1: Added: "with high risk of breast cancer" and "including to but not limited to" ▪ Section B10: Removed repeat statement "For breast cancer screening individuals with high risk of breast cancer. (For definitions on each of the risk levels, see the Policy Guidelines section.)" ▪ Section C2: Added " (i.e., mammography using low-dose x-rays for imaging)" to the statement ▪ Removed the statement "MRI of the breast is considered experimental / investigational for evaluation of residual tumor in individuals with positive margins after initial lumpectomy or breast conservation surgery"
	Updated Rationale Section
	Updated Reference Section
10-24-2023	Updated Description Section
	Updated Coding Section ▪ Removed ICD-10 Codes
	Updated Rationale Section
	Updated Reference Section

## REFERENCES

1. Jatoi I, Sung H, Jemal A. The Emergence of the Racial Disparity in U.S. Breast-Cancer Mortality. *N Engl J Med*. Jun 23 2022; 386(25): 2349-2352. PMID 35713541
2. Yedjou CG, Sims JN, Miele L, et al. Health and Racial Disparity in Breast Cancer. *Adv Exp Med Biol*. 2019; 1152: 31-49. PMID 31456178
3. Food and Drug Administration (FDA). 510(k) Summary of Safety and Effectiveness: Access Breast Coil 4/SMS. 2006; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf6/K060547.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf6/K060547.pdf). Accessed July 19, 2023.

4. Warner E, Messersmith H, Causer P, et al. Systematic review: using magnetic resonance imaging to screen women at high risk for breast cancer. *Ann Intern Med.* May 06 2008; 148(9): 671-9. PMID 18458280
5. Phi XA, Houssami N, Obdeijn IM, et al. Magnetic resonance imaging improves breast screening sensitivity in BRCA mutation carriers age  $\geq 50$  years: evidence from an individual patient data meta-analysis. *J Clin Oncol.* Feb 01 2015; 33(4): 349-56. PMID 25534390
6. Phi XA, Houssami N, Hooning MJ, et al. Accuracy of screening women at familial risk of breast cancer without a known gene mutation: Individual patient data meta-analysis. *Eur J Cancer.* Nov 2017; 85: 31-38. PMID 28886475
7. Ding W, Fan Z, Xu Y, et al. Magnetic resonance imaging in screening women at high risk of breast cancer: A meta-analysis. *Medicine (Baltimore).* Mar 10 2023; 102(10): e33146. PMID 36897691
8. Nelson HD, Fu R, Cantor A, et al. Effectiveness of Breast Cancer Screening: Systematic Review and Meta-analysis to Update the 2009 U.S. Preventive Services Task Force Recommendation. *Ann Intern Med.* Feb 16 2016; 164(4): 244-55. PMID 26756588
9. Nikitovic-Jokic M, Holubowich C. Magnetic Resonance Imaging as an Adjunct to Mammography for Breast Cancer Screening in Women at Less Than High Risk for Breast Cancer: A Health Technology Assessment. *Ont Health Technol Assess Ser.* 2016; 16(20): 1-30. PMID 27990198
10. Bakker MF, de Lange SV, Pijnappel RM, et al. Supplemental MRI Screening for Women with Extremely Dense Breast Tissue. *N Engl J Med.* Nov 28 2019; 381(22): 2091-2102. PMID 31774954
11. Berg WA, Zhang Z, Lehrer D, et al. Detection of breast cancer with addition of annual screening ultrasound or a single screening MRI to mammography in women with elevated breast cancer risk. *JAMA.* Apr 04 2012; 307(13): 1394-404. PMID 22474203
12. Cho N, Han W, Han BK, et al. Breast Cancer Screening With Mammography Plus Ultrasonography or Magnetic Resonance Imaging in Women 50 Years or Younger at Diagnosis and Treated With Breast Conservation Therapy. *JAMA Oncol.* Nov 01 2017; 3(11): 1495-1502. PMID 28655029
13. Kim EJ, Kang BJ, Kim SH, et al. Diagnostic Performance of and Breast Tissue Changes at Early Breast MR Imaging Surveillance in Women after Breast Conservation Therapy. *Radiology.* Sep 2017; 284(3): 656-666. PMID 28419815
14. de Bresser J, de Vos B, van der Ent F, et al. Breast MRI in clinically and mammographically occult breast cancer presenting with an axillary metastasis: a systematic review. *Eur J Surg Oncol.* Feb 2010; 36(2): 114-9. PMID 19822403
15. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Breast magnetic resonance imaging (MRI) for detection or diagnosis of primary or recurrent breast cancer TEC Assessments. 2004;Volume 19:Tab 1.
16. Lehman CD, Gatsonis C, Kuhl CK, et al. MRI evaluation of the contralateral breast in women with recently diagnosed breast cancer. *N Engl J Med.* Mar 29 2007; 356(13): 1295-303. PMID 17392300
17. Liberman L, Morris EA, Kim CM, et al. MR imaging findings in the contralateral breast of women with recently diagnosed breast cancer. *AJR Am J Roentgenol.* Feb 2003; 180(2): 333-41. PMID 12540428
18. Lehman CD, Blume JD, Thickman D, et al. Added cancer yield of MRI in screening the contralateral breast of women recently diagnosed with breast cancer: results from the

- International Breast Magnetic Resonance Consortium (IBMC) trial. J Surg Oncol. Oct 01 2005; 92(1): 9-15; discussion 15-6. PMID 16180217
19. Medeiros LR, Duarte CS, Rosa DD, et al. Accuracy of magnetic resonance in suspicious breast lesions: a systematic quantitative review and meta-analysis. Breast Cancer Res Treat. Apr 2011; 126(2): 273-85. PMID 21221772
  20. Zhang J, Wang L, Liu HF. Imaging Features Derived From Dynamic Contrast-Enhanced Magnetic Resonance Imaging to Differentiate Malignant From Benign Breast Lesions: A Systematic Review and Meta-Analysis. J Comput Assist Tomogr. May-Jun 2022; 46(3): 383-391. PMID 35575651
  21. Strobel K, Schrading S, Hansen NL, et al. Assessment of BI-RADS category 4 lesions detected with screening mammography and screening US: utility of MR imaging. Radiology. Feb 2015; 274(2): 343-51. PMID 25271857
  22. Li E, Li J, Song Y, et al. A comparative study of the diagnostic value of contrast-enhanced breast MR imaging and mammography on patients with BI-RADS 3-5 microcalcifications. PLoS One. 2014; 9(11): e111217. PMID 25365327
  23. Pereira RO, Luz LAD, Chagas DC, et al. Evaluation of the accuracy of mammography, ultrasound and magnetic resonance imaging in suspect breast lesions. Clinics (Sao Paulo). 2020; 75: e1805. PMID 32725074
  24. Houssami N, Ciatto S, Macaskill P, et al. Accuracy and surgical impact of magnetic resonance imaging in breast cancer staging: systematic review and meta-analysis in detection of multifocal and multicentric cancer. J Clin Oncol. Jul 01 2008; 26(19): 3248-58. PMID 18474876
  25. Houssami N, Turner R, Morrow M. Preoperative magnetic resonance imaging in breast cancer: meta-analysis of surgical outcomes. Ann Surg. Feb 2013; 257(2): 249-55. PMID 23187751
  26. Houssami N, Turner RM, Morrow M. Meta-analysis of pre-operative magnetic resonance imaging (MRI) and surgical treatment for breast cancer. Breast Cancer Res Treat. Sep 2017; 165(2): 273-283. PMID 28589366
  27. Plana MN, Carreira C, Muriel A, et al. Magnetic resonance imaging in the preoperative assessment of patients with primary breast cancer: systematic review of diagnostic accuracy and meta-analysis. Eur Radiol. Jan 2012; 22(1): 26-38. PMID 21847541
  28. Houssami N, Turner R, Macaskill P, et al. An individual person data meta-analysis of preoperative magnetic resonance imaging and breast cancer recurrence. J Clin Oncol. Feb 10 2014; 32(5): 392-401. PMID 24395846
  29. Li L, Zhang Q, Qian C, et al. Impact of Preoperative Magnetic Resonance Imaging on Surgical Outcomes in Women with Invasive Breast Cancer: A Systematic Review and Meta-Analysis. Int J Clin Pract. 2022; 2022: 6440952. PMID 36081810
  30. Brück N, Koskivuo I, Boström P, et al. Preoperative Magnetic Resonance Imaging in Patients With Stage I Invasive Ductal Breast Cancer: A Prospective Randomized Study. Scand J Surg. Mar 2018; 107(1): 14-22. PMID 28401771
  31. Mota BS, Reis YN, de Barros N, et al. Effects of preoperative magnetic resonance image on survival rates and surgical planning in breast cancer conservative surgery: randomized controlled trial (BREAST-MRI trial). Breast Cancer Res Treat. Apr 2023; 198(3): 447-461. PMID 36786946
  32. Gonzalez V, Sandelin K, Karlsson A, et al. Preoperative MRI of the breast (POMB) influences primary treatment in breast cancer: a prospective, randomized, multicenter study. World J Surg. Jul 2014; 38(7): 1685-93. PMID 24817517

33. Peters NH, van Esser S, van den Bosch MA, et al. Preoperative MRI and surgical management in patients with nonpalpable breast cancer: the MONET - randomised controlled trial. *Eur J Cancer*. Apr 2011; 47(6): 879-86. PMID 21195605
34. Turnbull L, Brown S, Harvey I, et al. Comparative effectiveness of MRI in breast cancer (COMICE) trial: a randomised controlled trial. *Lancet*. Feb 13 2010; 375(9714): 563-71. PMID 20159292
35. Onega T, Zhu W, Weiss JE, et al. Preoperative breast MRI and mortality in older women with breast cancer. *Breast Cancer Res Treat*. Jul 2018; 170(1): 149-157. PMID 29516372
36. Fortune-Greeley AK, Wheeler SB, Meyer AM, et al. Preoperative breast MRI and surgical outcomes in elderly women with invasive ductal and lobular carcinoma: a population-based study. *Breast Cancer Res Treat*. Jan 2014; 143(1): 203-12. PMID 24305978
37. Zeng Z, Amin A, Roy A, et al. Preoperative magnetic resonance imaging use and oncologic outcomes in premenopausal breast cancer patients. *NPJ Breast Cancer*. 2020; 6: 49. PMID 33083528
38. Drew PJ, Kerin MJ, Mahapatra T, et al. Evaluation of response to neoadjuvant chemoradiotherapy for locally advanced breast cancer with dynamic contrast-enhanced MRI of the breast. *Eur J Surg Oncol*. Nov 2001; 27(7): 617-20. PMID 11669587
39. Partridge SC, Gibbs JE, Lu Y, et al. Accuracy of MR imaging for revealing residual breast cancer in patients who have undergone neoadjuvant chemotherapy. *AJR Am J Roentgenol*. Nov 2002; 179(5): 1193-9. PMID 12388497
40. Marinovich ML, Macaskill P, Irwig L, et al. Agreement between MRI and pathologic breast tumor size after neoadjuvant chemotherapy, and comparison with alternative tests: individual patient data meta-analysis. *BMC Cancer*. Oct 08 2015; 15: 662. PMID 26449630
41. Lobbes MB, Prevost R, Smidt M, et al. The role of magnetic resonance imaging in assessing residual disease and pathologic complete response in breast cancer patients receiving neoadjuvant chemotherapy: a systematic review. *Insights Imaging*. Apr 2013; 4(2): 163-75. PMID 23359240
42. Morris EA, Schwartz LH, Drotman MB, et al. Evaluation of pectoralis major muscle in patients with posterior breast tumors on breast MR images: early experience. *Radiology*. Jan 2000; 214(1): 67-72. PMID 10644103
43. Kerslake RW, Carleton PJ, Fox JN, et al. Dynamic gradient-echo and fat-suppressed spin-echo contrast-enhanced MRI of the breast. *Clin Radiol*. Jul 1995; 50(7): 440-54. PMID 7614789
44. Rodenko GN, Harms SE, Pruneda JM, et al. MR imaging in the management before surgery of lobular carcinoma of the breast: correlation with pathology. *AJR Am J Roentgenol*. Dec 1996; 167(6): 1415-9. PMID 8956569
45. Xie Z, Xu W, Zhang H, et al. The value of MRI for downgrading of breast suspicious lesions detected on ultrasound. *BMC Med Imaging*. Jun 05 2023; 23(1): 72. PMID 37271827
46. de Lima Docema MF, Costa PA, de Andrade FE, et al. Magnetic resonance imaging-guided occult breast lesion localization and simultaneous sentinel lymph node mapping. *World J Surg Oncol*. Oct 23 2014; 12: 320. PMID 25341393
47. Marinovich ML, Houssami N, Macaskill P, et al. Meta-analysis of magnetic resonance imaging in detecting residual breast cancer after neoadjuvant therapy. *J Natl Cancer Inst*. Mar 06 2013; 105(5): 321-33. PMID 23297042
48. Li H, Yao L, Jin P, et al. MRI and PET/CT for evaluation of the pathological response to neoadjuvant chemotherapy in breast cancer: A systematic review and meta-analysis. *Breast*. Aug 2018; 40: 106-115. PMID 29758503

49. Janssen LM, den Dekker BM, Gilhuijs KGA, et al. MRI to assess response after neoadjuvant chemotherapy in breast cancer subtypes: a systematic review and meta-analysis. *NPJ Breast Cancer*. Sep 19 2022; 8(1): 107. PMID 36123365
50. Hylton NM, Blume JD, Bernreuter WK, et al. Locally advanced breast cancer: MR imaging for prediction of response to neoadjuvant chemotherapy--results from ACRIN 6657/I-SPY TRIAL. *Radiology*. Jun 2012; 263(3): 663-72. PMID 22623692
51. Lee JM, Orel SG, Czerniecki BJ, et al. MRI before reexcision surgery in patients with breast cancer. *AJR Am J Roentgenol*. Feb 2004; 182(2): 473-80. PMID 14736685
52. Bedrosian I, Mick R, Orel SG, et al. Changes in the surgical management of patients with breast carcinoma based on preoperative magnetic resonance imaging. *Cancer*. Aug 01 2003; 98(3): 468-73. PMID 12879462
53. Kawashima H, Tawara M, Suzuki M, et al. Effectiveness of dynamic MRI for diagnosing pericatricial minimal residual breast cancer following excisional biopsy. *Eur J Radiol*. Oct 2001; 40(1): 2-9. PMID 11673001
54. Orel SG, Reynolds C, Schnall MD, et al. Breast carcinoma: MR imaging before re-excisional biopsy. *Radiology*. Nov 1997; 205(2): 429-36. PMID 9356624
55. Weinstein SP, Orel SG, Heller R, et al. MR imaging of the breast in patients with invasive lobular carcinoma. *AJR Am J Roentgenol*. Feb 2001; 176(2): 399-406. PMID 11159081
56. Krammer J, Price ER, Jochelson MS, et al. Breast MR imaging for the assessment of residual disease following initial surgery for breast cancer with positive margins. *Eur Radiol*. Nov 2017; 27(11): 4812-4818. PMID 28567547
57. Lee RK, Kim HJ, Lee J. Role of breast magnetic resonance imaging in predicting residual lobular carcinoma in situ after initial excision. *Asian J Surg*. May 2018; 41(3): 279-284. PMID 28286021
58. Frei KA, Kinkel K, Bonel HM, et al. MR imaging of the breast in patients with positive margins after lumpectomy: influence of the time interval between lumpectomy and MR imaging. *AJR Am J Roentgenol*. Dec 2000; 175(6): 1577-84. PMID 11090379
59. Soderstrom CE, Harms SE, Farrell RS, et al. Detection with MR imaging of residual tumor in the breast soon after surgery. *AJR Am J Roentgenol*. Feb 1997; 168(2): 485-8. PMID 9016232
60. National Comprehensive Cancer Network (NCCN). NCCN clinical practice guidelines in oncology (NCCN guidelines): breast cancer. Version 4.2023. Updated March 23, 2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/breast.pdf](https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf). Accessed July 20, 2023.
61. National Comprehensive Cancer Network (NCCN). NCCN clinical practice guidelines in oncology (NCCN guidelines): breast cancer screening and diagnosis. Version 1.2023. Updated June 19, 2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/breast-screening.pdf](https://www.nccn.org/professionals/physician_gls/pdf/breast-screening.pdf). Accessed July 19, 2023.
62. National Comprehensive Cancer Network (NCCN). NCCN clinical practice guidelines in oncology (NCCN guidelines): genetic/familial high-risk assessment: breast, ovarian, and pancreatic. Version 3.2023. Updated February 13, 2023. [https://www.nccn.org/professionals/physician\\_gls/pdf/genetics\\_bop.pdf](https://www.nccn.org/professionals/physician_gls/pdf/genetics_bop.pdf). Accessed July 21, 2023.
63. American Cancer Society. Breast MRI. Updated January 14, 2022; <https://www.cancer.org/cancer/breast-cancer/screening-tests-and-early-detection/american-cancer-society-recommendations-for-the-early-detection-of-breast-cancer.html>. Accessed July 19, 2023.
64. American College of Radiology. Appropriateness criteria - breast cancer screening, revised 2017. <https://acsearch.acr.org/docs/70910/Narrative/>. Accessed July 24, 2023.

65. American College of Radiology. Appropriateness criteria - Palpable Breast Masses, revised 2022. <https://acsearch.acr.org/docs/69495/Narrative/>. Accessed July 23, 2023.
66. American College of Radiology. ACR Appropriateness Criteria stage I breast cancer: initial workup and surveillance for local recurrence and distant metastases in asymptomatic women. Revised 2019. <https://acsearch.acr.org/docs/69496/Narrative/> Accessed July 19, 2023.
67. American College of Radiology. Appropriateness criteria - Monitoring Response to Neoadjuvant Systemic Therapy for Breast Cancer, 2022. <https://acsearch.acr.org/docs/3099208/Narrative/>. Accessed July 22, 2023.
68. American College of Radiology. ACR Appropriateness Criteria - Transgender breast cancer screening. American College of Radiology. Published 2021. <https://acsearch.acr.org/docs/3155692/Narrative/> Accessed July 21, 2023.
69. American College of Radiology. ACR Appropriateness Criteria - Supplemental breast cancer screening based on breast density. American College of Radiology. Published 2021. <https://acsearch.acr.org/docs/3158166/Narrative/> Accessed July 20, 2023.
70. Monticciolo DL, Newell MS, Moy L, et al. Breast Cancer Screening in Women at Higher-Than-Average Risk: Recommendations From the ACR. *J Am Coll Radiol*. Mar 2018; 15(3 Pt A): 408-414. PMID 29371086
71. Khatcheressian JL, Wolff AC, Smith TJ, et al. American Society of Clinical Oncology 2006 update of the breast cancer follow-up and management guidelines in the adjuvant setting. *J Clin Oncol*. Nov 01 2006; 24(31): 5091-7. PMID 17033037
72. Khatcheressian JL, Hurley P, Bantug E, et al. Breast cancer follow-up and management after primary treatment: American Society of Clinical Oncology clinical practice guideline update. *J Clin Oncol*. Mar 01 2013; 31(7): 961-5. PMID 23129741
73. Mulder RL, Hudson MM, Bhatia S, et al. Updated Breast Cancer Surveillance Recommendations for Female Survivors of Childhood, Adolescent, and Young Adult Cancer From the International Guideline Harmonization Group. *J Clin Oncol*. Dec 10 2020; 38(35): 4194-4207. PMID 33078972
74. U.S. Preventive Services Task Force. Final Recommendation Statement. Breast Cancer: Screening. 2016; <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/breast-cancer-screening>. Accessed July 19, 2023.

## OTHER REFERENCES

1. Blue Cross and Blue Shield of Kansas Radiology Liaison Committee, February 10, 2004 (see Blue Cross and Blue Shield of Kansas Newsletter, Blue Shield Report MAC-01-04); February 2013; January 2014; January 2018, July 2023.
2. Blue Cross and Blue Shield of Kansas Medical Advisory Committee meeting, April 22, 2004 (see Blue Cross and Blue Shield of Kansas Newsletter, Blue Shield Report MAC-01-04).
3. Blue Cross and Blue Shield of Kansas Surgery Liaison Committee meeting, August 17, 2005 (see Blue Cross and Blue Shield of Kansas Newsletter, Blue Shield Report. MAC-03-05).
4. Blue Cross and Blue Shield of Kansas Medical Advisory Committee meeting, November 3, 2005 (see Blue Cross and Blue Shield of Kansas Newsletter, Blue Shield Report. MAC-03-05).
5. Blue Cross and Blue Shield of Kansas Consent Ballot (CB); Radiology Liaison Committee, September 2012; Surgery Liaison Committee, September 2012; November 2012.