

Medical Policy



Title: **Recombinant and Autologous Platelet-Derived Growth Factors for Wound Healing and Other Non-Orthopedic Conditions**

Related Policies:	<ul style="list-style-type: none">▪ <i>Orthopedic Applications of Platelet-Rich Plasma</i>▪ <i>Negative Pressure Wound Therapy in the Outpatient Setting</i>▪ <i>Bio-Engineered Skin and Soft Tissue Substitutes</i>
-------------------	--

Professional / Institutional
Original Effective Date: February 13, 2007 / December 15, 2008
Latest Review Date: March 12, 2024
Current Effective Date: March 1, 2018

State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).

The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.

The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.

If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none">• With diabetic lower-extremity ulcers	Interventions of interest are: <ul style="list-style-type: none">• Recombinant platelet-derived growth factor	Comparators of interest are: <ul style="list-style-type: none">• Standard wound care	Relevant outcomes include: <ul style="list-style-type: none">• Symptoms• Change in disease status• Morbid events• Quality of life• Treatment-related morbidity
Individuals:	Interventions of interest are:	Comparators of interest are:	Relevant outcomes include:

Populations	Interventions	Comparators	Outcomes
<ul style="list-style-type: none"> • With pressure ulcers 	<ul style="list-style-type: none"> • Recombinant platelet-derived growth factor 	<ul style="list-style-type: none"> • Standard wound care 	<ul style="list-style-type: none"> • Symptoms • Change in disease status • Morbid events • Quality of life • Treatment-related morbidity
Individuals: <ul style="list-style-type: none"> • With venous stasis leg ulcers 	Interventions of interest are: <ul style="list-style-type: none"> • Recombinant platelet-derived growth factor 	Comparators of interest are: <ul style="list-style-type: none"> • Standard wound care 	Relevant outcomes include: <ul style="list-style-type: none"> • Symptoms • Change in disease status • Morbid events • Quality of life • Treatment-related morbidity
Individuals: <ul style="list-style-type: none"> • With acute surgical or traumatic wounds 	Interventions of interest are: <ul style="list-style-type: none"> • Recombinant platelet-derived growth factor 	Comparators of interest are: <ul style="list-style-type: none"> • Standard wound care 	Relevant outcomes include: <ul style="list-style-type: none"> • Symptoms • Change in disease status • Morbid events • Quality of life • Treatment-related morbidity
Individuals: <ul style="list-style-type: none"> • With chronic wounds 	Interventions of interest are: <ul style="list-style-type: none"> • Platelet-rich plasma 	Comparators of interest are: <ul style="list-style-type: none"> • Standard wound care 	Relevant outcomes include: <ul style="list-style-type: none"> • Symptoms • Change in disease status • Morbid events • Quality of life • Treatment-related morbidity
Individuals: <ul style="list-style-type: none"> • With acute surgical or traumatic wounds 	Interventions of interest are: <ul style="list-style-type: none"> • Platelet-rich plasma 	Comparators of interest are: <ul style="list-style-type: none"> • Standard wound care 	Relevant outcomes include: <ul style="list-style-type: none"> • Symptoms • Change in disease status • Morbid events • Quality of life • Treatment-related morbidity

DESCRIPTION

The use of blood-derived growth factors, including recombinant platelet-derived growth factors (PDGFs) and platelet-rich plasma (PRP), has been suggested as a treatment for wounds or other miscellaneous non-orthopedic conditions, including but not limited to, diabetic ulcers, pressure ulcers, venous stasis ulcers, and surgical and traumatic wounds.

OBJECTIVE

The objective of this evidence review is to evaluate whether the use of recombinant platelet-derived growth factor or platelet-rich plasma improves health outcomes compared with standard care for diabetic ulcers, pressure ulcers, venous stasis ulcers, and surgical and traumatic wounds.

BACKGROUND

Wound Healing Treatment

A variety of growth factors have been found to play a role in wound healing, including platelet-derived growth factor (PDGF), epidermal growth factor, fibroblast growth factors, transforming growth factors, and insulin-like growth factors. Autologous platelets are a rich source of PDGF, transforming growth factors (that function as a mitogen for fibroblasts, smooth muscle cells, and osteoblasts), and vascular endothelial growth factors. Recombinant PDGF also has been extensively investigated for clinical use in wound healing.

Autologous platelet concentrate suspended in plasma, also known as platelet-rich plasma (PRP), can be prepared from samples of centrifuged autologous blood. Exposure to a solution of thrombin and calcium chloride degranulates platelets (releasing various growth factors) and results in the polymerization of fibrin from fibrinogen, creating a platelet gel. The platelet gel can then be applied to wounds or may be used as an adjunct to surgery to promote hemostasis and accelerate healing. In the operating room setting, PRP has been investigated as an adjunct to a variety of periodontal, reconstructive, and orthopedic procedures. For example, bone morphogenetic proteins are a transforming growth factor, and thus PRP has been used in conjunction with bone-replacement grafting (using either autologous grafts or bovine-derived xenograft) in periodontal and maxillofacial surgeries.

PRP is distinguished from fibrin glues or sealants, which have been used for many years as a surgical adjunct to promote local hemostasis at incision sites. Fibrin glue is created from platelet-poor plasma and consists primarily of fibrinogen. Commercial fibrin glues are created from pooled homologous human donors; Tisseel® (Baxter International) and Hemaseel® (Haemacure Corp.) are examples of commercially available fibrin sealants. Autologous fibrin sealants can also be created from platelet-poor plasma. This evidence review does not address the use of fibrin sealants.

Wound Closure Outcomes

This review addresses the use of recombinant PDGF products and PRP for non-orthopedic indications, which include a number of wound closure-related indications.

For this review, the primary endpoints of interest for the study of wound closure are as follows, consistent with guidance from the U.S. Food and Drug Administration (FDA) for the industry in developing products for the treatment of chronic cutaneous ulcer and burn wounds¹:

- Incidence of complete wound closure;
- Time to complete wound closure (reflecting accelerated wound closure);
- Incidence of complete wound closure following surgical wound closure;
- Pain control.

REGULATORY STATUS

Becaplermin

In 1997, becaplermin gel (Regranex®; Smith & Nephew), a recombinant PDGF product, was approved by the FDA for the following labeled indication:

"Regranex Gel is indicated for the treatment of lower extremity diabetic neuropathic ulcers that extend into the subcutaneous tissue or beyond and have an adequate blood supply. When used as an adjunct to, and not a substitute for, good ulcer care practices including initial sharp debridement, pressure relief and infection control, Regranex Gel increases the complete healing of diabetic ulcers.

The efficacy of Regranex Gel for the treatment of diabetic neuropathic ulcers that do not extend through the dermis into subcutaneous tissue or ischemic diabetic ulcers ... has not been evaluated...Regranex is not intended to be used in wounds that close by primary intention."

In 2008, the manufacturer added the following black box warning to the labeling for Regranex®: "An increased rate of mortality secondary to malignancy was observed in patients treated with 3 or more tubes of Regranex Gel in a postmarketing retrospective cohort study. Regranex Gel should only be used when the benefits can be expected to outweigh the risks. Regranex Gel should be used with caution in patients with known malignancy."

In 2018, the "Boxed Warning" and "Warnings and Precautions" were changed to remove "increased rate of cancer mortality" and "cancer mortality," respectively.

Platelet-Rich Plasma

The FDA regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation, Title 21, parts 1270 and 1271. Blood products such as PRP are included in these regulations.

Under these regulations, certain products including blood products such as PRP are exempt and therefore, do not follow the traditional FDA regulatory pathway. To date, the FDA has not attempted to regulate activated PRP.²

Numerous PRP preparation systems have been cleared for marketing by the FDA through the 510(k) process. These devices are intended to concentrate patient plasma at the point of care during bone grafting procedures. The use of different devices and procedures can lead to variable concentrations of active platelets and associated proteins, increasing variability between studies of clinical efficacy.

POLICY

- A. Recombinant platelet-derived growth factor (i.e., becaplermin) may be considered **medically necessary** when used as an adjunct to standard wound management for the following indications:
1. Neuropathic diabetic ulcers extending into the subcutaneous tissue
Appropriate candidates for becaplermin gel for treatment of neuropathic ulcers should meet **ALL** of the following criteria:
 - a. Adequate tissue oxygenation, as measured by a transcutaneous partial pressure of oxygen of 30 mm Hg or greater on the foot dorsum or at the margin of the ulcer **AND**
 - b. Full-thickness ulcer (i.e., stage III or IV), extending through dermis into subcutaneous tissues **AND**
 - c. Participation in a wound-management program, which includes sharp debridement, pressure relief (i.e., non-weight-bearing), and infection control
 2. Pressure ulcers extending into the subcutaneous tissue
Appropriate candidates for becaplermin gel for the treatment of pressure ulcers should meet **ALL** of the following criteria:
 - a. Full-thickness ulcer (i.e., stage III or IV), extending through dermis into subcutaneous tissues **AND**
 - b. Ulcer in an anatomic location that can be offloaded for the duration of treatment **AND**
 - c. Albumin concentration >2.5 dL **AND**
 - d. Total lymphocyte count >1000/uL **AND**
 - e. Normal values of vitamins A and C
- B. Other applications of recombinant platelet-derived growth factor (i.e., becaplermin) are considered **experimental / investigational**, including, but not limited to:
1. Ischemic ulcers
 2. Venous stasis ulcers, and
 3. Ulcers not extending through the dermis into the subcutaneous tissue
- C. Use of platelet-rich plasma (i.e., autologous blood-derived preparations) is considered **experimental / investigational** for the treatment of acute or chronic wounds, including surgical wounds and nonhealing ulcers.

POLICY GUIDELINES

- A. Becaplermin
1. Individuals are typically treated once daily for up to 20 weeks or until completely healed. Application of the gel may be performed by the individual in the home.
 2. Becaplermin is available in 2-, 7.5-, and 15-g tubes and is applied in a thin continuous layer, about 1/16 of an inch thick (i.e., 1.6 mm or the thickness of a dime). The amount of the gel used will depend on the size of the ulcer, measured in square centimeters. However, an average-sized ulcer, measuring 3 cm², treated for an average length of time of 85 days, will require a little more than one 15-g tube. If the ulcer is treated for the maximum length of time of 140 days, 1.75 of the 15-g tubes would be required.

Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

RATIONALE

This evidence review has been updated regularly with searches of the PubMed database. The most recent literature update was performed through November 14, 2023.

The platelet-rich plasma (PRP) portion of this evidence review on the platelet-derived wound healing formulae was originally based on a 1992 TEC Assessment that primarily focused on the Procurement process.³ This preparation method is no longer commercially available. Currently, a large number of devices are available for the preparation of PRP or PRP gel. The amount and mixture of growth factors produced by different cell-separating systems vary, and it is unknown whether platelet activation before an injection is necessary.^{4,5,6,7,8}

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function^{3,4}including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Promotion of greater diversity and inclusion in clinical research of historically marginalized groups (e.g., People of Color [African-American, Asian, Black, Latino and Native American]; LGBTQIA (Lesbian, Gay, Bisexual, Transgender, Queer, Intersex, Asexual); Women; and People with Disabilities [Physical and Invisible]) allows policy populations to be more reflective of and findings more applicable to our diverse members. While we also strive to use inclusive language related to these groups in our policies, use of gender-specific nouns (e.g., women, men, sisters, etc.) will continue when reflective of language used in publications describing study populations.

RECOMBINANT PLATELET-DERIVED GROWTH FACTOR FOR DIABETIC LOWER-EXTREMITY ULCERS

Clinical Context and Therapy Purpose

The purpose of recombinant platelet-derived growth factor (PDGF) is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with diabetic lower-extremity ulcers.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with diabetic lower-extremity ulcers.

Interventions

The therapy being considered is recombinant PDGF.

Comparators

Comparators of interest include standard wound care.

Outcomes

The general outcomes of interest are symptoms, change in disease status, morbid events, quality of life (QOL), and treatment-related morbidity.

Follow-up at 20 weeks is of interest for recombinant PDGF to monitor relevant outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

The portion of this evidence review on the use of recombinant PDGF (becaplermin gel) was informed by a 1999 TEC Assessment, which found that the evidence supported the conclusion that becaplermin gel, in conjunction with good wound care, improves the health outcomes of patients with chronic neuropathic diabetic ulcers that met the patient selection criteria defined therein.⁹ Becaplermin gel plus good wound care resulted in a 43% complete wound closure rate, compared with 28% for patients treated with good wound care alone. Becaplermin gel also appeared to reduce the average time to complete wound closure.

Systematic Reviews

A 2014 systematic review identified 6 RCTs (N=992 patients) that compared recombinant PDGFs with placebo or standard care.¹⁰ There was a combined odds ratio of 1.53 (95% confidence interval [CI], 1.14 to 2.04; p=.004) favoring recombinant PDGF for complete healing rate.

Sridharan et al (2018) conducted a systematic review and meta-analysis of RCTs on topical growth factors compared with standard of care in patients with diabetic foot ulcers (DFUs). The

primary outcome of concern was complete healing and the second outcome of concern was the existence of adverse events. Rankogram was generated based on the surface under the cumulative ranking curve. In total, 26 studies with 2088 participants and 1018 adverse events were included. The pooled odds ratio estimates for recombinant human epidermal growth factor (rhEGF), autologous-PRP, and recombinant human platelet-derived growth factor were 5.7 [95% CI, 3.34 to 10.37], 2.65 [95% CI, 1.65 to 4.54], and 1.97 [95% CI, 1.54 to 2.55] respectively. The surface under the cumulative ranking curve for rhEGF was 0.95; sensitivity analysis did not reveal significant changes from pooled estimates and rankogram. With regard to adverse events, no differences were observed for the overall risk of adverse events between the growth factors; however, the growth factors were observed to lower the risk of lower limb amputations compared to standard of care. The results lead the authors to conclude that rhEGF, recombinant human platelet-derived growth factor, and autologous PRP significantly improved the healing rate when used as adjuvants to the standard of care. Compared to other growth factors, rhEGF performed better. The limitations of this study include the following: the strength of most of the outcomes assessed was low, and the findings may not be applicable for DFU with infection or osteomyelitis.¹¹

Table 1. Systematic Reviews of Trials Assessing Recombinant Platelet-Derived Growth Factor for Diabetic Lower-Extremity Ulcers

Study (Year)	Literature Search	Studies	Participants	N	Design	Results
Sridharan et al (2018)	Dec 2016	RCTs	Patients with diabetic lower-extremity ulcers treated with platelet-derived growth factor	2088	RCTs	Pooled analysis estimated rhEGF, PRP, rhPDGF

PRP: autologous platelet-rich plasma; RCT: Randomized Controlled Trial; rhEGF: recombinant epidermal growth factor; rhPDGF: recombinant human platelet-derived growth factor

Retrospective Studies

A 2005 industry-sponsored study assessed the effectiveness of recombinant PDGF for diabetic neuropathic foot ulcers in actual clinical practice.¹² Among a cohort of 24,898 patients in wound care centers, those subjects whose wounds did not heal over an 8-week observation period were eligible for the study and were retrospectively assessed over 20 weeks or until they healed. Any subject with an open wound who was lost to follow-up was considered unhealed. Of the nearly 25000 patients treated for foot ulcers, 2394 (9.6%) received recombinant PDGF. A propensity score method with covariates to statistically model treatment selection was used to adjust for selection bias; results were stratified by 5 propensity score groups. Overall, the rate of healing was 26.5% in the control group and 33.5% in patients treated with recombinant PDGF. The relative risk (RR), controlling for the propensity to receive PDGF, was 1.32 (95% CI, 1.22 to 1.38) for healing and 0.65 (95% CI, 0.54 to 0.78) for amputation (6.4% in controls vs. 4.9% in the PDGF group). The analysis also indicated those who received PDGF were more likely to be younger, male, and have older wounds-factors not known to affect wound healing. These results support the clinical utility of recombinant PDGF for treatment of diabetic neuropathic foot ulcers in actual clinical practice.

Section Summary: Recombinant Platelet-Derived Growth Factor for Diabetic Lower-Extremity Ulcers

Published evidence includes an industry-sponsored study and 2 systematic reviews that showed an improvement in treatment over control for tested outcome measures.

RECOMBINANT PLATELET-DERIVED GROWTH FACTOR FOR PRESSURE ULCERS

Clinical Context and Therapy Purpose

The purpose of recombinant PDGF is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with pressure ulcers.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with pressure ulcers.

Interventions

The therapy being considered is recombinant PDGF.

Comparators

Comparators of interest include standard wound care.

Outcomes

The general outcomes of interest are symptoms, change in disease status, morbid events, QOL, and treatment-related morbidity.

Though not completely standardized, follow-up for pressure ulcer symptoms would typically occur in the months after starting treatment.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

Randomized Controlled Trials

Rees et al (1999) conducted an RCT focusing on the use of becaplermin gel as a treatment for pressure ulcers.¹³ Patient selection criteria included full-thickness ulcers and an anatomic location where pressure could be offloaded during treatment. This latter patient selection criterion might have limited the number of patients with pressure ulcers who would have been considered candidates for becaplermin therapy. Patients were randomized to 1 of 4 parallel treatment groups and received either a placebo or 1 of 3 dosages of becaplermin. All patients received a

standardized program of good wound care. In the 2 groups treated with the once-daily dosage (becaplermin 0.01% or 0.03%), the incidence of complete healing was significantly improved compared with the placebo group. There was no difference in outcome between the 0.01% and 0.03% groups, suggesting there is no clinical benefit in increasing the potency above 0.01%. A third group received becaplermin 0.01% twice daily. That group did not report improved outcomes compared with placebo, a finding that is unexplained.

Section Summary: Recombinant Platelet-Derived Growth Factor for Pressure Ulcers

Published evidence includes a multicenter, double-blind RCT that showed an improvement in treatment over control for tested outcome measures.

RECOMBINANT PLATELET-DERIVED GROWTH FACTOR FOR VENOUS STASIS LEG ULCERS

Clinical Context and Therapy Purpose

The purpose of recombinant PDGF is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with venous stasis leg ulcers.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with venous stasis leg ulcers.

Interventions

The therapy being considered is recombinant PDGF.

Comparators

Comparators of interest include standard wound care.

Outcomes

The general outcomes of interest are symptoms, change in disease status, morbid events, QOL, and treatment-related morbidity.

Though not completely standardized, follow-up for venous stasis leg ulcer symptoms would typically occur in the months after starting treatment.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

Randomized Controlled Trials

Senet et al (2011) in France, published a multicenter, double-blind RCT of becaplermin gel for venous leg ulcers.¹⁴ There was no significant difference between the becaplermin (n=28) and control hydrogel (n=31) groups for any of the outcome measures, which included complete closure rates after 8 and 12 weeks, changed ulcer area, and changed ulcer-related pain and QOL.

Section Summary: Recombinant Platelet-Derived Growth Factor for Venous Stasis Leg Ulcers

Published evidence includes a multicenter, double-blind RCT that showed no difference between treatment and control for tested outcome measures.

RECOMBINANT PLATELET-DERIVED GROWTH FACTOR FOR ACUTE SURGICAL OR TRAUMATIC WOUNDS

Clinical Context and Therapy Purpose

The purpose of recombinant PDGF is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with acute surgical or traumatic wounds.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with acute surgical or traumatic wounds.

Interventions

The therapy being considered is recombinant PDGFs.

Comparators

Comparators of interest include standard wound care.

Outcomes

The general outcomes of interest are symptoms, change in disease status, morbid events, QOL, and treatment-related morbidity.

Though not completely standardized, follow-up for acute surgical or traumatic wound symptoms would typically occur in the months after starting treatment.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Topical recombinant PDGF has also been investigated for repair of work-related fingertip injuries. A 2005 prospective controlled trial alternately assigned 50 patients (fingertip wound area ≥ 1.5 cm, with or without phalangeal exposure) to daily treatment with PDGF (n=25) or surgical reconstruction (n=25).¹⁵ Statistical analysis showed that baseline characteristics of the 2 groups were similar for patient age, wound area (2.2 to 2.4 cm), and distribution of fingertip injuries across the digits. Assessment by an independent physician showed that, compared with the surgical intervention, treatment with recombinant PDGF resulted in faster return to work (10 days vs. 38 days) and wound healing (25 days vs. 35 days), less functional impairment (10% vs. 22%), and less need for physical therapy (20% vs. 56%), respectively. Fingertips treated with PDGF were also reported to have satisfactory aesthetic results, while surgically treated fingertips were shorter and often unsightly. These results, if confirmed in additional RCTs, could lead to improvement in health outcomes for patients with fingertip injuries. However, this trial was limited by its small sample size, method of randomization, and potential for investigator bias (although examining physicians were blinded to treatment allocation, actual treatment might have been obvious).

Adverse Events

Growth factors cause cells to divide more rapidly. For this reason, the manufacturer of Regranex continued to monitor studies that started before its approval (in December 1997) for any evidence of adverse events, such as increased numbers of cancers. In a long-term safety study completed in 2001, more deaths from cancer occurred among patients who used Regranex than in those who did not. A subsequent study was performed using a health insurance database that covered the period from January 1998 through June 2003. This trial identified 2 groups of patients with similar diagnoses, drug use, and use of health services: 1 group used Regranex, and the other group did not. Results showed there were more deaths from cancer among patients who were given 3 or more prescriptions for Regranex than deaths for those not treated with Regranex. No single type of cancer was identified; deaths from all types of cancer were observed. In 2008, the U.S. Food and Drug Administration concluded that the increased risk of death from cancer in patients who used 3 or more tubes of Regranex was 5 times higher compared with those who did not use Regranex, prompting the manufacturer to add a black box warning to the labeling for Regranex. The risk of new cancers among Regranex users was not increased compared with nonusers, although the duration of follow-up of patients in this study was not long enough to detect new cancers.

Section Summary: Recombinant Platelet-Derived Growth Factor for Acute Surgical or Traumatic Wounds

Published evidence includes nonrandomized controlled trials reporting satisfactory aesthetic results. Larger RCTs are required to confirm and expound on these results.

PLATELET-RICH PLASMA FOR CHRONIC WOUNDS

Clinical Context and Therapy Purpose

The purpose of PRP is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with chronic wounds

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with chronic wounds.

Interventions

The therapy being considered is PRP.

Comparators

Comparators of interest include standard wound care.

Outcomes

The general outcomes of interest are symptoms, change in disease status, morbid events, QOL, and treatment-related morbidity.

Though not completely standardized, follow-up for chronic wound symptoms would typically occur in the months after starting treatment.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

DIABETIC FOOT ULCERS

Systematic Reviews

A number of systematic reviews of the evidence on PRP have been published.^{16,17,18,19,20,21,22,23} These reviews are heterogeneous in whether they pooled data from studies reflecting a variety of wound types^{16,17,24,18,25} or focused on specific wound types, primarily diabetic foot ulcers.^{19,20,21,22} Results from the reviews that pooled data from a variety of wound types^{16,17,24,18,25} are not discussed herein as their design precludes drawing conclusions about the applicability of the review findings to specific wound types. As the majority of the RCTs included in the systematic reviews were published post-2014, herein are summarized those systematic reviews that focused on specific wound types with search dates that extend to at least 2015.^{21,22,23}

Three recent systematic reviews have evaluated studies of PRP for individuals with diabetic foot ulcers.^{21,22,23} Table 2 provides a crosswalk of the studies included in the systematic reviews.

Table 2. Comparison of Trials of Platelet-Rich Plasma in Individuals with Diabetic Foot Ulcers Included in Systematic Reviews

Primary Study (Year)	Li 2019 ^{21,}	Qu 2021 ^{22,}	Deng 2023 ^{23,}
Ahmed 2017 ^{26,}	●	●	●
Alamdari 2021 ^{27,}			●
Chen 2008 ^{a28,}	●		
Driver 2006 ^{29,}	●	●	●
Elsaid 2020 ^{30,}		●	●
Friese 2007 (conference proceeding) ^{31,}	●		●
Game 2018 ^{32,}		●	
Gude 2019 ^{33,}		●	
Habeeb 2020 ^{34,}			●
Helmy 2021 ^{35,}			●
Hossam 2021 ^{36,}			●
Jeong 2010 ^{37,}			●
Kakagia 2007 ^{38,}	●	●	●
Karimi 2016 ^{39,}		●	●
Li 2012 ^{a 40,}			●
Li 2015 ^{41,}	●	●	●
Liu 2016 ^{a42,}	●		●
Liao 2020 ^{43,}			●
Meamar 2021 ^{44,}			●
Ma 2014 ^{a45,}	●		
Milek 2017 ^{46,}		●	
Qi 2014 ^{a47,}	●		
Rainys 2019 ^{48,}			●
Saad Setta 2011 ^{49,}	●	●	●
Saldamacchia 2004 ^{50,}	●	●	●
Serra 2013 ^{51,}	●	●	
Singh 2018 ^{52,}		●	●
Steed 1992 ^{53,}			●
Steed 1996 ^{54,}			●

Primary Study (Year)	Li 2019 ²¹ ,	Qu 2021 ²² ,	Deng 2023 ²³ ,
Tofign 2022 ⁵⁵ ,			●
Xie 2020 ⁵⁶ ,		●	
Yang 2017 ⁵⁷ ,		●	
Zhang 2016 ^{a58} ,	●		
Zhou 2015 ^{a59} ,	●		
Zhu 2012 ^{a60} ,	●		

^a In Chinese

Tables 3 and 4 summarize the characteristics and results of the 3 systematic reviews that have evaluated studies of PRP for individuals with diabetic foot ulcers. ^{21,22,23}

In their meta-analysis, Li et al (2019) assessed the efficacy and safety of autologous platelet-rich gel for topical treatment of diabetic chronic cutaneous ulcers²¹. Their analysis included 15 RCTs with 829 patients. Results indicated that autologous platelet-rich gel had a significant positive effect on healing rate, shorter healing time, and lower risk of infection than conventional treatment. Autologous platelet-rich gel also had a significantly lower incidence of infection when compared with conventional treatment (odds ratio [OR]=0.34; 95% CI: 0.15 to 0.77; p=.009). This meta-analysis was limited by a high or unclear risk of bias among the trials, which may indicate the trials were underpowered. Also, some studies had small sample sizes and limited outcome information. Further, 7 of the included trials are available only in the Chinese language. Finally, most of the trials were 8 to 12 weeks long and others only 2 to 5 weeks, making it difficult to analyze the relationship of time of observation to ulcer healing.

The Agency for Healthcare Research and Quality (AHRQ) (2020) published a Technology Assessment on Platelet-Rich Plasma for Wound Care in the Medicare Population. This Technology Assessment was requested by the Centers for Medicare & Medicaid Services to inform reconsideration of a National Coverage Decision on autologous blood-derived products for chronic non-healing wounds.⁶¹ This Technology Assessment evaluates evidence in lower extremity diabetic ulcers, lower extremity venous ulcers and pressure ulcers. Separate meta-analyses were conducted for each wound type. Here the focus is on findings for lower extremity diabetic ulcers and those for the other populations are discussed below. Risk of bias of individual studies was assessed using the Cochrane Collaboration's Risk of Bias 2 tool and rated high in 8 RCTs (57.14%), moderate in 6 RCTs (42.86%) and high in the 1 observational study (100%). Strength of the body of evidence was rated based on the Evidence-based Practice Center methods guide. The findings of this Technology Assessment indicated that there is moderate-strength evidence that PRP modestly increases complete wound closure (see meta-analysis results in Table 4 below) and low-strength evidence that PRP may shorten time to wound closure (meta-analysis not feasible). However, due to risk of bias and severe imprecision, evidence is insufficient to draw conclusions about other important outcomes, including wound infection, amputation, pain reduction, and wound recurrence. Important limitations of the literature were described as "inadequate description of offloading and wound care procedures, wound characteristics, PRP formulation techniques, concentration and volume; inadequate length of follow-up, and lack of stratification by comorbidities and other patient characteristics, such as diabetes control, vascular perfusion, and under representation of older adults."

A meta-analysis by Deng et al (2023) assessed 22 RCTs (N=1559) to determine the safety and efficacy of PRP to treat diabetic foot ulcers.²³ Results indicated PRP significantly increased the overall healing rate of diabetic foot ulcers compared with standard treatment (risk ratio [RR]=1.42; 95% CI: 1.30 to 1.56; $p<.001$; $I^2=55\%$). PRP increased the complete wound healing time of diabetic foot ulcers compared to conventional treatment (mean difference [MD]=-3.13; 95% CI: -5.86 to -0.39; $p<.001$; $I^2=97.5\%$) and resulted in a greater reduction in diabetic foot ulcer area (MD=1.02; 95% CI: 0.51 to 1.53; $p<.001$; $I^2=36\%$). The rate of amputation, reported by 3 trials, significantly reduced risk for the autologous PRP group (RR=0.35; 95% CI, 0.15 to 0.83; $p<.001$; $I^2=0\%$). Four studies reported adverse events, and pooled analysis revealed a similar rate of events between the PRP and control groups (RR=0.96; 95% CI, 0.57 to 1.61; $p>0.05$; 35%). The authors reported no significant publication bias was detected by funnel plot analysis; however, a sensitivity analysis suggested that the pooled outcome assessment for time to wound healing may be affected by considerable inter-study variability. The low number of high-quality of studies available on PRP for diabetic foot ulcers and the low number of studies reporting some outcomes of interest were limitations of this meta-analysis.

Table 3. Characteristics of Key Systematic Reviews with Meta-Analyses in Individuals with Diabetic Foot Ulcers

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Li (2019) ²¹ ,	2004-2017	15	Patients with diabetic chronic cutaneous wounds/ulcers that do not show signs of healing in 4 weeks	N=829 (14-117)	RCTs	NR
Qu (2021) ²² ,	Inception-2020	14	Adults with lower extremity diabetic ulcers, lower extremity venous ulcers, or pressure ulcers in any location, or a mix of these 3 etiologies	N=1,096 (range NR)	RCTs	Median = 6 wk (range, none to 11 months)
Deng (2023) ²³ ,	Inception-2023	22	Adults with diabetic foot ulcers	N=1559	RCTs	NR

NR: not reported; RCT: randomized controlled trial; wk: week(s); y: year(s).

Table 4. Results of Key Systematic Reviews with Meta-Analyses in Individuals with Diabetic Foot Ulcers

Study	Healing Rate	Healing Time	Complete Wound Healing	Risk of Infection	Wound complications	Pain Reduction	Recurrence
Li (2019) ²¹ ,							
RR	1.39						
MD		-9.18					
OR				0.34			

Study	Healing Rate	Healing Time	Complete Wound Healing	Risk of Infection	Wound complications	Pain Reduction	Recurrence
95% CI	1.29 to 1.50	-11.32 to -7.05		0.15 to 0.77			
P-value	<.001	<.001		.009			
Qu (2021) ^{22,}							
RR			1.20	0.77			2.09
WMD						-1.10 ^a	
95% CI			1.09 to 1.32	0.54 to 1.11		-1.81 to -0.39	0.31 to 13.93
P-value							
Deng (2023) ^{23,}							
RR	1.42				.096		
MD		-3.13					
95% CI	1.30 to 1.56	-5.86 to -0.39			0.57 to 1.61		
P-value	<.001	<.001			.203		

^a Visual Analog Scale

CI: confidence interval; MD: mean difference; OR: odds ratio; RR: risk ratio; WMD: weighted mean difference; Z: indicates overall effect.

Randomized Controlled Trials

Key characteristics and results of several RCTs of diabetic foot ulcers published subsequent to the AHRQ review (2020) are summarized in Tables 5 and 6 below.

One RCT of PRP dressing with total-contact casting compared to standard saline dressing for diabetic foot ulcers (Gupta et al [2021])^{62,} did not find significant differences in rates of ulcer area reduction or absolute ulcer area reduction between groups over the 6-week study period.

Another RCT of PRP versus standard wound care found accelerated rates of ulcer area reduction and decreased incidence of wound infections with PRP treatment; however, the difference in the percentage of healed surface between groups lost statistical significance at 6, 7, or 8 weeks of follow-up and it is unclear whether complete wound healing was achieved in either group.^{36,}

Table 5. Summary of Key RCT Characteristics

Study	Countries	Sites	Dates	Participants	Intervention	Control
Gupta et al (2021) ^{62,}	India	1	2016 to 2018	Individuals with diabetes mellitus with noninfected diabetic foot ulcers with total ulcer area of 20 cm ² or less on the plantar surface	Autologous intralesional PRP therapy with total contact casting (n=30)	Saline dressing (n= 30)
Hossam et al (2022) ^{36,}	Egypt	1	2018	Individuals with type 1 or 2 diabetes with non-ischemic revascularized chronic diabetic foot ulcers of more than 6 months duration with no clinical signs of infection, Wagner grade 1 or 2, and ASA physical status class 2	Autologous intralesional CaCl ₂ -activated PRP therapy (injection and/or gel) with saline gauze (n=40)	Standard wound care with moist dressing with or without collagenase ointment (n=40)

ASA: American Society of Anesthesiologists; PRP: platelet-rich plasma; RCT: randomized controlled trial.

Table 6. Summary of Key RCT Results

Study	Complete Healing	Percentage of Healed Surface Area ^a	Complete Healing Time	Pain	Quality of Life	Infection	Recurrence
Gupta et al (2021) ^{62,}	NR	6 weeks: 85.98% vs 81.72%; p=NR	NR	NR	NR	NR	NR
Hossam et al (2022) ^{36,}	95% vs 77.8% ^b ; p<.001	1 week: 23.1% vs 0%; p=.002 5 weeks: 89.2% vs 60.1%; p<.001 8 weeks: 96.7% vs 95.5%; p=.529	NR	NR	NR	PRP: 4 (10%) Control: 18 (45%) with 4 resulting in amputation p<.001	NR

NR: not reported; PRP: platelet-rich plasma; RCT: randomized controlled trial.

^a Percentage of healed surface area in treatment vs. control groups.

^b Proportion of patients with complete healing in treatment (n=38) vs. control groups (n=28) at 6 and 9 weeks, respectively.

Study relevance, design, and conduct limitations are summarized below in Tables 9 and 10.

Other Chronic Wound Types

The AHRQ (2020) Technology Assessment on Platelet-Rich Plasma for Wound Care in the Medicare Population described above also evaluated evidence on use of PRP in individuals with lower extremity venous ulcers and individuals with pressure ulcers.^{63,}

For individuals with lower extremity venous ulcers, the evidence included 8 RCTs and 3 observational studies (total N=615). The majority compared PRP to management without PRP. Risk of bias was described as moderate due to randomization and outcome measurement limitations. There were no significant differences between PRP versus management without PRP in complete wound closure (RR=1.49; 95% CI: 0.72 to 3.06; 5 studies, N=250; $I^2=29.4\%$), wound recurrence (RR=0.38; 95% CI: 0.09 to 1.57), wound infection (RR=0.79; 95% CI: 0.22 to 2.81), or quality of life as measured by the Chronic Lower Limb Venous Insufficiency Questionnaire (weighted mean difference [WMD]=10.99; 95%CI: -50.5 to 72.5). For the outcomes time to complete wound closure and pain, meta-analysis of 2 studies was not possible due to insufficient data and findings were mixed between studies on both outcomes. The strength of evidence was rated as 'insufficient' to draw conclusions on all outcomes. Oliveira et al (2020) also conducted a meta-analysis of cost and effectiveness of studies of PRP for venous ulcers.^{64,} Based on fewer studies identified from searches only through July 2018, although their findings indicated greater reductions in wound area for PRP, findings were consistent with the ARHQ review in finding no significant difference in complete wound closure (RR=2.54; 95% CI, 0.42 to 15.30; 4 studies, N=156; $I^2=69\%$).

For individuals with pressure ulcers, the AHRQ Technology Assessment (2020)^{22,} included 1 RCT and 1 comparative observational study (total N not reported). The comparator was serum physiological dressing in the RCT and saline dressing in the observational study. Risk of bias of the primary studies was described as moderate, due to limitations in the randomization process and outcome measurement, deviations from intended interventions, and selective outcome reporting. Although both studies found that PRP significantly reduced wound size (strength of evidence=insufficient), neither study evaluated other important outcomes, such as complete wound closure.

A meta-analysis by Fang et al (2023) pooled data from 6 studies on patients treated for lower extremity venous ulcers with PRP.^{65,} A total of 294 patients were included, with 148 patients in the PRP group and 146 in the control group. PRP was found to have a greater reduction in elliptical area at the end of treatment compared to the control group (Mean difference [MD], -1.19; 95% CI, -1.8 to -.058; P=.0001) with a moderate quality of evidence. The healing rate also favored PRP over the control group (RR=5.73; 95% CI, 3.29 to 9.99; P<.00001) with a moderate quality to the evidence base. The authors suggest there may be publication bias in the calculation of these pooled estimates according to Egger's test.

Randomized Controlled Trials

Two RCTs of PRP for chronic wounds (Saha et al [2020])^{66,67}, were identified as published subsequent to the AHRQ review (2020).²² Key characteristics and results of selected RCTs are reported in Tables 7 and 8 below.

Saha et al.'s analyses included 91.5% (n=108) of randomized individuals. Participants were mostly males in their late 40s with trophic ulcer duration of 13.4 months. Reduction in ulcer surface area, the primary outcome, was significantly greater for the PRP group from the first week (38.96% vs 12.46%; $p<.001$) through the fifth (and last) week of follow-up (91.10% vs 79.77%; $p<.001$). However, healing time and recurrence were not reported and there was no significant difference in complete healing rate.

Shehab et al (2023) conducted an RCT of adjunct PRP in addition to compression therapy in individuals with post-phlebotic venous ulcers.⁶⁷ Forty patients were randomized 1:1 to either PRP and compression therapy or placebo. The median number of treatments was 6 (range 3 to 6). Both participants and outcome assessors were blinded to treatment allocation. The median ulcer surface area, the primary outcome, was significantly lower for the PRP group (4 cm² vs 10 cm²; $p=.036$) as well as the median volume of ulcers (1 cm³ vs 3 cm³; $p=.008$). This translated to individuals in the PRP group experiencing a larger drop in ulcer area (74% vs 40%; $p=.008$) and volume (81% vs 48%; $p=.013$) compared to placebo. Differences in VAS pain scores were observed in favor of the PRP group at both the 3-month and 6-month follow-ups. Nine patients in the PRP group had complete wound healing, but the authors did not report the rate of complete healing in the control group, and healing time and recurrence were not reported.

Table 7. Summary of Key RCT Characteristics

Study	Countries	Sites	Dates	Participants	Intervention	Control
Saha et al (2020) ⁶⁶ ,	Iran	1	2016 to 2018	Individuals with clinically diagnosed trophic ulcers due to leprosy	Autologous PRP therapy with total contact casting (n=59)	Only total contact casting (n=59)
Shehab et al (2023) ⁶⁷ ,	Egypt	1	2019 to 2020	Adults with chronic post-phlebotic lower limb venous ulcers	Autologous PRP therapy with compression therapy (n=20)	Placebo plus compression therapy (n=20)

PRP: platelet-rich plasma; RCT: randomized controlled trial.

Table 8. Summary of Key RCT Results

Study	Complete Healing	Healing Time	Pain	Quality of Life	Infection	Recurrence
Saha et al (2020) ⁶⁶ ,	22 (39.29%) vs 11 (21.15%); p NR	NR	NR	NR	0 vs 0; p=.773	NR
Shehab et al (2023) ⁶⁷ ,	9 (45%) vs NR	NR	BL: 6.5 vs 6.4; p=.43 3 mos: 1 vs 4.5; p<.0001 6 mos: 0.5 vs 2.2; p<.0001	NR	NR	NR

NR: not reported; RCT: randomized controlled trial.

^a Percentage of healed surface area in study and control groups at 6 weeks.

Tables 9 and 10 summarize the relevance and design and conduct limitations of selected RCTs.

Table 9. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Saha et al (2020) ⁶⁶ ,	4. Single site in Iran	4. Short duration of treatment; 8 weeks		1. Recurrence, quality of life not addressed 5. Clinical significance of difference in wound surface area not prespecified	1. 4 weeks follow-up post-treatment insufficient to assess long-term efficacy
Gupta et al (2021) ⁶² ,	4. Single site in India	4. Short duration of treatment; 6 weeks	3. Total-contact casting not used in control group	1. Complete wound healing, recurrence, quality of life not addressed 5. Clinical significance of difference in wound surface area not prespecified	1. 6 week study period insufficient to assess long-term efficacy
Hossam et al (2022) ³⁶ ,	4. Single site in Egypt	1. Frequency and type of		1. Complete wound healing,	1. 8 week study period

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
		PRP treatment (injection and/or gel) not standardized 4. Short duration of treatment; 8 weeks		recurrence, quality of life not addressed 5. Primary outcome differences and timepoints were not prespecified	insufficient to assess long-term efficacy
Shehab et al (2023) ⁶⁷ ,	4. Single site in Egypt	1. Frequency and type of PRP treatment (injection and/or gel) not standardized 4. Short duration of treatment; 6 weeks	1. Placebo treatment not clearly defined	1. Recurrence, quality of life not addressed	

PRP: platelet-rich plasma.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 10. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Saha et al (2020) ⁶⁶ ,						
Gupta et al (2021) ⁶² ,		1-3. Blinding not described			1. Power calculations not reported	3. Confidence intervals and/or p values not reported
Hossam et al (2022) ³⁶ ,		1-3. Blinding not described		1. High loss to follow-up or missing data; reasons for and extent of missingness	1. Power calculations not reported	3. Confidence intervals not reported

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
				unclear at all timepoints		
Shehab et al (2023) ⁶⁷ ,					1. Power calculations not reported	4. Complete healing rate not reported for the control group

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Section Summary: Platelet-Rich Plasma for Chronic Wounds

The evidence for autologous PRP for a variety of chronic wounds includes systematic reviews, RCTs, which have been summarized in several systematic reviews, and nonrandomized trials. In meta-analyses of individuals with lower extremity diabetic ulcers, PRP demonstrated an improvement over the control groups in complete wound closure and healing time, but moderate to high risk of bias and imprecision preclude drawing conclusions on other important outcomes such as recurrence, infection, amputation, and quality of life. In individuals with venous ulcers, PRP did not demonstrate an improvement over the control groups in complete wound closure, recurrence, wound infection or quality of life, although imprecision likely precluded identifying differences on these outcomes. In individuals with pressure ulcers, although PRP reduced wound size, other important outcomes such as complete wound closure were not measured. Overall, the studies are small and of low quality, and the results should be interpreted with caution.

PLATELET-RICH PLASMA FOR ACUTE SURGICAL OR TRAUMATIC WOUNDS

Clinical Context and Therapy Purpose

The purpose of PRP is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with acute surgical or traumatic wounds.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with acute surgical or traumatic wounds.

Interventions

The therapy being considered is PRP.

Comparators

Comparators of interest include standard wound care.

Outcomes

The general outcomes of interest are symptoms, change in disease status, morbid events, QOL, and treatment-related morbidity.

Though not completely standardized, follow-up for acute surgical or traumatic wound symptoms would typically occur in the months after starting treatment.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

SURGICAL WOUNDS

Aortic Arch Repair

Zhou et al (2015) reported on a double-blind RCT with 80 patients that assessed the effect of PRP on the amount of blood transfused in the perioperative period for elective ascending and transverse aortic arch repair.⁶⁸ An anesthesiologist prepared the PRP so that the surgeon was unaware of the treatment group. The volume of PRP transfused was 726 mL and led to a reduction in transfusion rates for red blood cells, frozen plasma, cryoprecipitate, and platelets by 34% to 70% ($p < .02$). Hospital length of stay was also reduced (9.4 days vs. 12.7 days). There was no difference in mortality between the 2 groups (1 patient in each group) and no significant differences in postoperative complications or other outcome measures. Corroboration of the effect of PRP on perioperative blood transfusion is needed.

Sternotomy Wounds

Serraino et al (2015) reported on a large series with historical controls that assessed the occurrence of deep sternal wound infections in patients who underwent cardiac surgery either with (2010 to 2012, 422 consecutive patients) or without (2007 to 2009, 671 consecutive patients) application of PRP.⁶⁹ The 2 groups were comparable at baseline. At the end of cardiac surgery, PRP gel was applied to the sternum before the closure of subcutaneous tissue. Rates of both deep and superficial wound infections were reduced in the patients treated with PRP (deep: 0.2% vs. 1.5%; superficial: 0.5% vs. 2.8%). Interpretation of these results is limited by likely differences in treatments over time. RCTs are needed to evaluate this potential use of PRP.

Zhu et al (2023) published a meta-analysis of the effect of PRP on sternal wound healing.⁷⁰ Eleven studies with a total of 8961 cardiac surgery patients were included. Patients were either treated with PRP (n=3663) or control therapies (n=5298), with sample sizes ranging from 44 to 2000 participants. PRP was found to have a significantly lower rate of sternal wound infection (Odds ratio [OR], 0.11; 95% CI, 0.03 to 0.34; $p < .001$; I^2 , 0%), deep sternal wound infection (OR, 0.29; 95% CI, 0.16 to 0.51; $p < .001$; I^2 , 32%) and superficial sternal wound infection (OR, 0.20; 95% CI, 0.13 to 0.33; $p < .001$; I^2 , 0%) compared to patients in the control cardiac surgery groups. All pooled estimates at no to low heterogeneity (0% to 32%). The poor quality of included studies, heterogeneous PRP preparations, and heterogeneous cardiac surgeries limit the interpretation of the results.

Otolaryngology

El-Anwar et al (2016) reported on an RCT that evaluated PRP in 44 children (age range, 12 to 23 months) undergoing repair of a complete cleft palate.⁷¹ Speech and velopharyngeal valve movement on follow-up were evaluated by 3 judges who “usually assessed every patient blindly,” physical examination, video nasoendoscopy, and audio recording of audio perceptual assessment. At 6 months, PRP-treated patients had better nasality grade on audio perceptual assessment ($p = .024$) and better velopharyngeal closure on endoscopy ($p = .016$).

A 2008 double-blind RCT assessed the efficacy of PRP following tonsillectomy in 70 children (age range, 4 to 15 years).⁷² PRP was placed into the tonsil beds of half of the children, where it was directly visible. To compare pain symptoms and recovery, a daily diary was completed by the patient or a family member for 10 days after surgery. A FACES Pain Scale was used for children ages 4 to 7 years, while a numeric pain rating scale was used for children older than 7 years. Diaries from 83% of patients showed no differences in pain, medication doses, activity, and days eating solid foods between the 2 conditions.

Other Surgical Wounds

A 2011 Norwegian trial of PRP applied to saphenous vein harvest sites after wound closure found no differences in the incidence of wound infection or cosmetic result.⁷³

Alamdari et al (2018) published a clinical trial evaluating the efficacy of pleurodesis with a combination of PRP and fibrin glue compared with surgical intervention. The study population consisted of 52 esophageal cancer patients with postoperative chylothorax who did not respond to conservative management. Each member of the population was consecutively and randomly allocated to either a PRP fibrin glue pleurodesis arm or a surgical thoracic duct ligation arm. Twenty-six in each arm were treated with their respective interventions. The patients were distributed into the intervention arms in a way that made each group similar in terms of tumor size and patient demographics. This distribution procedure was not described. All patients (26) in the PRP treatment arm and 20 (76.9%) in the surgery arm were successfully treated ($p = .009$). Seven patients (26.92%) of the PRP required a second application of the PRP fibrin glue after a week. The mean length of hospital stay was higher in the surgery group (53.50 ± 16.662 days) than the PRP group (36.04 ± 8.224 days; $p < .001$). The study was limited due to the fact the procedure for randomization was not described and, thus, its efficacy cannot be evaluated.²⁷

Mohamadi et al (2019) reported on an RCT of 110 participants in Tehran that evaluated the efficacy of PRP gel in wound healing time following pilonidal sinus surgery.⁷⁴ Each group included

55 participants. Follow-up duration was 9 weeks. In the treatment group, PRP was both injected into the wound weekly, as well as applied to the wound surface and covered with latex. In the control group, wound dressing was described as "classic", but no other details were provided. Little to no detail was provided about specific outcome assessment methods (*i.e.*, "pain duration was inquired from participants"). All patients completed the study and were included in the outcome assessments. PRP significantly shortened mean healing time (4.8 vs 8.7 weeks; $p < .001$), pain duration (1.3 vs 3.4 weeks; $p < .001$), and antibiotic consumption duration (0.57 vs 1.74 weeks; $p < .001$). This RCT also performed regression analyses to evaluate the correlation between different factors in wound healing activity. Significant negative associations were found between healing time and wound volume and pain duration and angiogenesis. Notable limitations of this study included unclearly defined wound dressing in the comparator group, unblinded and poorly defined outcome assessment, short-term follow-up and lack of assessment of other important health outcomes.

Slaninka et al (2020) published an RCT that evaluated PRP in 24 individuals in the Czech Republic who had undergone dermo-epidermal skin grafts taken from the thigh area.⁷⁵ Indications for skin grafts were primarily hard-to-heal lower leg wounds. PRP was applied to 1 thigh and covered with Vaseline-impregnated, open-weave gauze and gauze. The control was the other thigh, which was also covered with open-weave gauze and gauze, but without PRP. Of the 24 included individuals, 3 (12.5%) were excluded after developing infections. The infections were described as first occurring on the non-PRP wound and only subsequently occurring on the PRP wound after several days. PRP significantly shortened median healing time (14 days vs 18 days; $p = .026$). No other outcomes were reported. Notable limitations of the RCT include its small sample size and that it did not address important health outcomes and harms.

Wang et al (2023) published a meta-analysis

Traumatic Wounds

Kazakos et al (2009) reported on a prospective RCT that evaluated treatment of acute traumatic wounds (open fractures, closed fractures with skin necrosis, friction burns) with platelet gel in 59 consecutive patients (27 PRP, 32 controls).⁷⁶ Conventional treatment consisted of topical washing and cleaning of the wounds, removal of the necrotic tissue, and dressing in petroleum jelly gauze every 2 days. In all patients with open tibial fractures, an external fixation system was applied. PRP gel was applied to the wounds after surgical debridement and placement of the external fixation system. The time needed for preparation and application of the PRP gel was 52 minutes. After that, PRP gel was applied to the wounds once weekly in the outpatient clinic until there was adequate tissue regeneration (mean, 21 days) sufficient to undergo reconstructive plastic surgery. Control patients receiving conventional treatment required a mean of 41 days for adequate tissue regeneration. Pain scores were significantly lower in PRP-treated patients at 2 and 3 weeks (visual analog scale score, 58 PRP vs. 80 controls). Although these results are encouraging, additional study with a larger number of patients is needed.

Marck et al (2016) reported on a randomized, double-blind, within-patient-controlled study in patients with deep dermal to full-thickness burns undergoing split-skin graft, comparing PRP with usual care.⁷⁷ The study randomized 52 patients, 50 of whom received the allocated PRP intervention. There were no significant differences in short-term (5 to 7 days) rates in graft take in the intervention and control areas on each patient. At 3, 6, and 12 months, there were no significant differences in skin appearance or epithelialization scores.

Yeung et al (2018) performed a prospective RCT to test the efficacy of lyophilized platelet-rich plasma powder (LPRP) on the healing rate of wounds in patients with deep, second-degree burn injuries in comparison with a control group using a placebo. LPRP was dissolved in a solution and applied on deep second-degree burn wounds once per day for 4 consecutive days. Twenty-seven patients with deep second-degree burns were recruited and then those that met eligibility criteria were randomized into 2 groups. The LPRP group received the intervention (n=15) and the control group received a placebo application (n=12). A concentration of 1.0×10^7 platelets/cm² (wound area) was sprayed on the wound evenly. Function was assessed by the percentage of wound closure and bacteria picking out rate at weeks 2 and 3. The mean burn area of control for the LPRP was 75.65 ± 50.72 cm² and 99.73 ± 70.17 cm² (p=.0013), respectively. In the control group, the original wound area was 25.49 cm² at baseline, 23.79 cm² (6.67% healed) at week 2, and 4.34 cm² (86.40% healed) at week 3. In the LPRP group, the original wound area was 84.36 cm², followed by 23.96 cm² (71.59% healed) at week 2, and 0.63 cm² (99.24% healed) at week 3. The wound closure rate at week 2 in the LPRP group reached nearly 80% and was greater than 90% by week 3, showing a significant difference (p<.05). Alternatively, in the control group, the wound closure rates were 60% and 80% in 2 and 3 weeks, respectively. The postoperative infection rate in the LPRP (26.67%) was lower than the control group (33.33%). Neither was significant, statistically. One limitation of this study is that the powder is made by an independent lab and dissolved in a specified amount of water. This provides an opportunity for accidental error-this may also be the case with some liquid PRP.⁷⁸

Huang et al (2021) published a meta-analysis of 8 RCTs representing 539 patients with burn wounds.⁷⁹ The healing rate of burn wounds was improved with PRP (OR, 4.43; 95% CI, 2.13 to 9.22), yielding a significantly shorter wound healing time (OR, -4.23; 95% CI, -5.48 to -2.98) compared to conventional dressings for both superficial and deep burn groups. Incidence of adverse events, pain scores, and scar scores was also all improved in the PRP treatment group. Interpretation of results is limited by risks of bias arising from lack of blinding, small study size, heterogenous PRP preparations, and short follow-up durations.

Imam et al (2023) published a meta-analysis of 13 comparative studies, including 808 individuals with burn wounds who were treated with PRP (n=413) or standard wound therapy (n=395) with sample sizes ranging from 25 to 100 individuals.⁸⁰ PRP had a shorter healing time than compared to standard therapy (Mean difference [MD], -5.80; 95% CI, -7.73 to -3.88; p<.001) as well as a higher healing rate (OR, 3.14; 95% CI, 2.05 to 4.8; p<.001) although these pooled estimates had substantial ($I^2=93\%$) and moderate heterogeneity ($I^2=42\%$), respectively. Individuals treated with PRP also had a higher percentage of graft take area (MD, 4.39; 95% CI, 1.51 to 7.26; p<.001) and higher percent of area healed (MD, 12.67; 95% CI, 9.79 to 15.55, p<.001) compared to standard therapy for burn wounds with a low level of heterogeneity. No differences were observed in the graft take ratio or infection rates which showed low heterogeneity across studies in the pooled estimates. Interpretation of results is limited by risks of bias arising from low overall study quality, small study sizes, heterogenous PRP preparations, limited number of studies included for some comparisons, and short follow-up durations.

Section Summary: Platelet-Rich Plasma for Acute Surgical or Traumatic Wounds

The evidence for autologous PRP for a variety of acute surgical or traumatic wounds includes systematic reviews and RCTs. For a variety of other conditions, studies have either not

demonstrated a benefit or have demonstrated small benefits in studies with methodologic limitations.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Physicians

In 2015, the American College of Physicians (ACP) published guidelines on treatment of pressure ulcers.⁸¹ The guidelines noted that "although low-quality evidence suggests that dressings containing PDGF [platelet-derived growth factors] promote healing, ACP supports the use of other dressings such as hydrocolloid and foam dressings, which are effective at promoting healing and cost less than PDGF dressings." A search of the ACP website on December 1, 2020 found that this 2015 guideline is now listed as inactive.

Association for the Advancement of Wound Care

The Association for the Advancement of Wound Care developed guideline recommendations for the management of pressure ulcers (2010)⁸² and venous ulcers (2015)⁸³:

- Pressure ulcer: "Growth factors are not indicated for PU [pressure ulcers] at this time." (level C evidence - no randomized controlled trials (RCTs) available comparing growth factors with A-level dressings)⁸²,
- Venous ulcer: "Platelet-derived growth factor has shown no significant effects on VU [venous ulcer healing or recurrence]." (level A evidence)⁸³,

National Institute for Health and Care Excellence

In 2019, the National Institute for Health and Care Excellence updated its guidance on the prevention and management of diabetic foot problems.⁸⁴ The guidance stated that neither autologous platelet-rich plasma gel nor platelet-derived growth factors should be offered in the treatment of diabetic foot ulcers.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Ongoing and Unpublished Clinical Trials

Some larger studies that might influence this review are listed in Table 9.

Table 9. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT05979584	Platelet Rich Plasma VS Platelet Fibrin Plasma in Treatment of Diabetes Foot Ulcer: a Randomized Controlled Trial	56	Dec 2024
NCT02312596 ^a	A Prospective, Randomized Clinical Trial of PRP Concepts Fibrin Bio-Matrix in Non-Healing Diabetic Foot Ulcers	200	Dec 2021
NCT02312570 ^a	A Prospective, Randomized Clinical Trial of PRP Concepts Fibrin Bio-Matrix in Chronic Non-Healing Pressure Ulcers	200	Dec 2021
NCT02307448 ^a	Effectiveness of Autologous Platelet Rich Plasma in the Treatment of Chronic Non-Healing Wounds	80	Dec 2022
NCT02402374 ^a	Randomized, Placebo-controlled, Blind-assessor Study to Evaluate the Safety and Efficacy of Autologous Platelet Rich Plasma Gel Prepared With the RegenKit-BCT Plus Family of Kits for the Treatment of Diabetic Foot Ulcer	192	Dec 2020 (unknown)
<i>Unpublished</i>			
NCT02071979 ^a	Registry Trial of the Effectiveness of Platelet Rich Plasma for Chronic Non-Healing Wounds (CMS)	1500	Jan 2018 (terminated; updated 01/18)

NCT: national clinical trial; PRP: autologous platelet-rich plasma.

^a Denotes industry-sponsored or cosponsored trial.

CODING

The following codes for treatment and procedures applicable to this policy are included below for informational purposes. This may not be a comprehensive list of procedure codes applicable to this policy.

Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

The code(s) listed below are medically necessary ONLY if the procedure is performed according to the "Policy" section of this document.

CPT/HCPCS	
86999	Unlisted transfusion medicine procedure
0232T	Injection(s), platelet rich plasma, any site, including image guidance, harvesting and preparation when performed
G0460	Autologous platelet rich plasma for non-diabetic chronic wounds/ulcers, including phlebotomy, centrifugation, and all other preparatory procedures, administration and dressings, per treatment
G0465	Autologous platelet rich plasma (prp) for diabetic chronic wounds/ulcers, using an fda-cleared device (includes administration, dressings, phlebotomy, centrifugation, and all other preparatory procedures, per treatment) (eff 01/01/2022)
P9020	Platelet rich plasma, each unit
S0157	Becaplermin gel 0.01%, 0.5 gm
S9055	Procuren or other growth factor preparation to promote wound healing

REVISIONS	
06-05-2012	<p>Policy added to the bcbsks.com web site. A stand alone policy was developed based on policy language previously contained in the Wound Care: Skin Substitutes and Growth Factors medical policy.</p> <p>In Policy section:</p> <ul style="list-style-type: none"> ▪ The new stand-alone policy adds the following: <p>"C. Use of autologous blood-derived preparations (i.e., platelet-rich plasma) is considered experimental / investigational. This includes, but is not limited to, use in the following situations:</p> <ol style="list-style-type: none"> 1. Treatment of acute or chronic wounds including nonhealing ulcers 2. Adjunctive use in surgical procedures 3. Primary use (injection) for other conditions such as epicondylitis (i.e., tennis elbow), plantar fasciitis, or Dupuytren's contracture"
02-05-2014	<p>Description section updated</p> <p>Policy section reformatted – no policy statement changes made.</p> <p>Rationale section updated</p> <p>In Coding section:</p> <ul style="list-style-type: none"> ▪ HCPCS Code added: G0460 ▪ Coding information bullets updated ▪ ICD-10 Diagnoses Codes added <p>References updated</p>

REVISIONS	
10-29-2015	Description section updated
	In Policy section: <ul style="list-style-type: none"> ▪ In Item C added "surgical sounds and" and removed "This includes, but is not limited to, use in the following situations:", "Adjunctive use in surgical procedures", and "Primary use (injection) for other condition such as epicondylitis(i.e. tennis elbow), plantar fasciitis, or Dupuytren's contracture" to read, "Use of autologous blood-derived preparations (i.e., platelet-rich plasma) is considered experimental / investigational for the treatment of acute or chronic wounds, including surgical wounds and nonhealing ulcers."
	Rationale section updated
	In Coding section: <ul style="list-style-type: none"> ▪ Updated coding notations.
	References updated
04-25-2016	Description section updated
	Rationale section updated
	In Coding section: <ul style="list-style-type: none"> ▪ Coding notations updated
	References updated
03-01-2017	Title changed to "Recombinant and Autologous Platelet-Derived Growth Factors for Wound Healing and Other Non-Orthopedic Conditions" from "Recombinant and Autologous Platelet-Derived Growth Factors as a Treatment of Wound Healing and Other Non-Orthopedic Conditions"
	Description section updated
	In Policy section: <ul style="list-style-type: none"> ▪ In Item A 2 d added "/uL" to correctly read "Total lymphocyte count >1000/uL" – no change in policy intent.
	Rationale section updated
	In Coding section: <ul style="list-style-type: none"> ▪ Removed ICD-10 Codes: E10.610, E10.618, E10.69, E11.610, E11.618, E11.69, I70.331, I70.332, I70.333, I70.334, I70.335, I70.338, I70.341, I70.342, I70.343, I70.344, I70.345, I70.348, I70.35, I70.731, I70.732, I70.733, I70.734, I70.735, I70.738, I70.741, I70.742, I70.743, I70.744, I70.745, I70.748, I70.75 ▪ Added ICD-10 Codes: L97.121, L97.122, L97.123, L97.124, L97.211, L97.212, L97.213, L97.214, L97.221, L97.222, L97.223, L97.224, L97.311, L97.312, L97.313, L97.314, L97.321, L97.322, L97.323, L97.324, L97.411, L97.412, L97.413, L97.414, L97.421, L97.422, L97.423, L97.424, L97.511, L97.512, L97.513, L97.514, L97.521, L97.522, L97.523, L97.524, L97.811, L97.812, L97.813, L97.814, L97.821, L97.822, L97.823, L97.824, L98.491, L98.492, L98.493, L98.494
	References updated
03-01-2018	Description section updated
	In Policy section: <ul style="list-style-type: none"> ▪ In Item B added "recombinant platelet-derived growth factor" to read "Other applications of recombinant platelet-derived growth factor (i.e., becaplermin) are considered experimental / investigational, including, but not limited to:" ▪ Updated Policy Guidelines
	Rationale section updated
	References updated
04-10-2019	Description section updated
	Rationale section updated
	References updated
03-23-2021	Description section updated
	Rationale section updated

REVISIONS	
	References updated
01-01-2022	In Coding section: Revised nomenclature G0460 Autologous platelet rich plasma (prp) for diabetic chronic wounds/ulcers, using an fda-cleared device (includes administration, dressings, phlebotomy, centrifugation, and all other preparatory procedures, per treatment) effective 01-01-22
03-08-2022	Updated Description Section
	Updated Rationale Section
	Updated Coding Section <ul style="list-style-type: none"> ▪ Added CPT code G0465 ▪ Changed ICD-10 Diagnoses to code ranges ▪ Removed ICD-10 codes E10.41, E10.42, E10.43, E10.44, E10.49, E10.621, E10.622, E10.628, E11.44, E11.621, E11.622, E11.628, I70.231, I70.232, I70.233, I70.234, I70.235, I70.238, I70.241, I70.242, I70.243, I70.244, I70.245, I70.248, I70.25, I70.431, I70.432, I70.433, I70.434, I70.435, I70.438, I70.441, I70.442, I70.443, I70.444, I70.445, I70.448, I70.45, I70.531, I70.532, I70.533, I70.534, I70.535, I70.538, I70.541, I70.542, I70.543, I70.544, I70.545, I70.548, I70.55, I70.631, I70.632, I70.633, I70.634, I70.635, I70.638, I70.641, I70.642, I70.643, I70.644, I70.645, I70.648, I70.65 ▪ Removed Coding bullets <ul style="list-style-type: none"> ○ There is a CPT category III code for injections of platelet-rich plasma: 0232T. ○ The instructions issued with the code state that it is not to be reported with codes 20550, 20551, 20600-20610, 20926, 76942, 77002, 77012, 77021 and 86965. ○ Code 0232T includes the harvesting and preparation of the platelet-rich plasma. ○ For situations other than injection (when 0232T would be reported), no specific CPT codes describe the preparation of autologous blood-derived products but CPT code 86999 can be used. It has been reported that providers have used CPT code 20926 (tissue graft, other) to describe the overall procedure. It is questionable whether platelet-rich plasma is appropriately considered a tissue graft. ○ The American Medical Association's Department of Coding instructs that placement of PRP into an operative site is an inclusive component of the operative procedure performed and not separately reported. ○ There is also a HCPCS code for this treatment: G0460.
	Updated References Section
02-28-2023	Updated Description Section
	Updated Rationale Section
	Updated References Section
03-12-2024	Updated Description Section
	Updated Rationale Section
	Updated Coding Section <ul style="list-style-type: none"> ▪ Removed ICD-10 Codes
	Updated References Section

REFERENCES

1. U.S. Food and Drug Administration. Guidance for Industry: Chronic Cutaneous Ulcer and Burn Wounds -- Developing Products for Treatment. Rockville, MD: Food and Drug Administration; 2006 June.
2. U.S. Food and Drug Administration (FDA). Tissue and Tissue Products. 2016; <http://www.fda.gov/BiologicsBloodVaccines/TissueTissueProducts/>. Accessed December 13, 2023.
3. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Becaplermin for wound healing. TEC Assessments. 1999;Volume 14:Tab 5.
4. Crovetti G, Martinelli G, Issi M, et al. Platelet gel for healing cutaneous chronic wounds. *Transfus Apher Sci.* Apr 2004; 30(2): 145-51. PMID 15062754
5. Eppley BL, Woodell JE, Higgins J. Platelet quantification and growth factor analysis from platelet-rich plasma: implications for wound healing. *Plast Reconstr Surg.* Nov 2004; 114(6): 1502-8. PMID 15509939
6. Kevy SV, Jacobson MS. Comparison of methods for point of care preparation of autologous platelet gel. *J Extra Corpor Technol.* Mar 2004; 36(1): 28-35. PMID 15095838
7. Castillo TN, Pouliot MA, Kim HJ, et al. Comparison of growth factor and platelet concentration from commercial platelet-rich plasma separation systems. *Am J Sports Med.* Feb 2011; 39(2): 266-71. PMID 21051428
8. Mazzucco L, Balbo V, Cattana E, et al. Not every PRP-gel is born equal. Evaluation of growth factor availability for tissues through four PRP-gel preparations: Fibrinet, RegenPRP-Kit, Plateltex and one manual procedure. *Vox Sang.* Aug 2009; 97(2): 110-8. PMID 19392780
9. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Growth factors for wound healing. TEC Evaluations. 1992;7:352-377.
10. Zhao XH, Gu HF, Xu ZR, et al. Efficacy of topical recombinant human platelet-derived growth factor for treatment of diabetic lower-extremity ulcers: Systematic review and meta-analysis. *Metabolism.* Oct 2014; 63(10): 1304-13. PMID 25060693
11. Sridharan K, Sivaramakrishnan G. Growth factors for diabetic foot ulcers: mixed treatment comparison analysis of randomized clinical trials. *Br J Clin Pharmacol.* Mar 2018; 84(3): 434-444. PMID 29148070
12. Margolis DJ, Bartus C, Hoffstad O, et al. Effectiveness of recombinant human platelet-derived growth factor for the treatment of diabetic neuropathic foot ulcers. *Wound Repair Regen.* 2005; 13(6): 531-6. PMID 16283867
13. Rees RS, Robson MC, Smiell JM, et al. Becaplermin gel in the treatment of pressure ulcers: a phase II randomized, double-blind, placebo-controlled study. *Wound Repair Regen.* 1999; 7(3): 141-7. PMID 10417749
14. Senet P, Vicaut E, Beneton N, et al. Topical treatment of hypertensive leg ulcers with platelet-derived growth factor-BB: a randomized controlled trial. *Arch Dermatol.* Aug 2011; 147(8): 926-30. PMID 21482863
15. Freedman BM, Oplinger EH, Freedman IS. Topical becaplermin improves outcomes in work related fingertip injuries. *J Trauma.* Oct 2005; 59(4): 965-8. PMID 16374289
16. Martinez-Zapata MJ, Martí-Carvajal AJ, Solà I, et al. Autologous platelet-rich plasma for treating chronic wounds. *Cochrane Database Syst Rev.* May 25 2016; 2016(5): CD006899. PMID 27223580
17. Martinez-Zapata MJ, Martí-Carvajal AJ, Solà I, et al. Autologous platelet-rich plasma for treating chronic wounds. *Cochrane Database Syst Rev.* Oct 17 2012; 10: CD006899. PMID 23076929

18. Carter MJ, Fylling CP, Parnell LK. Use of platelet rich plasma gel on wound healing: a systematic review and meta-analysis. *Eplasty*. 2011; 11: e38. PMID 22028946
19. Picard F, Hersant B, Bosc R, et al. The growing evidence for the use of platelet-rich plasma on diabetic chronic wounds: A review and a proposal for a new standard care. *Wound Repair Regen*. Sep 2015; 23(5): 638-43. PMID 26019054
20. Del Pino-Sedeño T, Trujillo-Martín MM, Andia I, et al. Platelet-rich plasma for the treatment of diabetic foot ulcers: A meta-analysis. *Wound Repair Regen*. Mar 2019; 27(2): 170-182. PMID 30575212
21. Li Y, Gao Y, Gao Y, et al. Autologous platelet-rich gel treatment for diabetic chronic cutaneous ulcers: A meta-analysis of randomized controlled trials. *J Diabetes*. May 2019; 11(5): 359-369. PMID 30182534
22. Qu W, Wang Z, Hunt C, Morrow AS, Urtecho M, Amin M, Shah S, Hasan B, Abd-Rabu R, Ashmore Z, Kubrova E, Prokop LJ, Murad MH. Platelet-Rich Plasma for Wound Care in the Medicare Population. Technology Assessment Program Project ID 040-353-492. (Prepared by the Mayo Clinic Evidence-based Practice Center under Contract No. HHS2902015000131.) Rockville, MD: Agency for Healthcare Research and Quality. <https://www.ahrq.gov/sites/default/files/wysiwyg/research/findings/ta/prp/prp-wound-care.pdf>. Accessed December 13, 2023.
23. Deng J, Yang M, Zhang X, et al. Efficacy and safety of autologous platelet-rich plasma for diabetic foot ulcer healing: a systematic review and meta-analysis of randomized controlled trials. *J Orthop Surg Res*. May 19 2023; 18(1): 370. PMID 37202812
24. Martínez-Zapata MJ, Martí-Carvajal A, Solà I, et al. Efficacy and safety of the use of autologous plasma rich in platelets for tissue regeneration: a systematic review. *Transfusion*. Jan 2009; 49(1): 44-56. PMID 18954394
25. Qu S, Hu Z, Zhang Y, et al. Clinical Studies on Platelet-Rich Plasma Therapy for Chronic Cutaneous Ulcers: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Adv Wound Care (New Rochelle)*. Feb 2022; 11(2): 56-69. PMID 33607926
26. Ahmed M, Reffat SA, Hassan A, et al. Platelet-Rich Plasma for the Treatment of Clean Diabetic Foot Ulcers. *Ann Vasc Surg*. Jan 2017; 38: 206-211. PMID 27522981
27. Alamdari DH, Asadi M, Rahim AN, et al. Efficacy and Safety of Pleurodesis Using Platelet-Rich Plasma and Fibrin Glue in Management of Postoperative Chylothorax After Esophagectomy. *World J Surg*. Apr 2018; 42(4): 1046-1055. PMID 28986682
28. Chen HY, Chen CX, Liang Y, Wang J. Efficacy of autologous platelet rich gel in the treatment of refractory diabetic foot. *Chin J New Clin Med*. 2008; 17:1-2.
29. Driver VR, Hanft J, Fylling CP, et al. A prospective, randomized, controlled trial of autologous platelet-rich plasma gel for the treatment of diabetic foot ulcers. *Ostomy Wound Manage*. Jun 2006; 52(6): 68-70, 72, 74 passim. PMID 16799184
30. Elsaid A, El-Said M, Emile S, et al. Randomized Controlled Trial on Autologous Platelet-Rich Plasma Versus Saline Dressing in Treatment of Non-healing Diabetic Foot Ulcers. *World J Surg*. Apr 2020; 44(4): 1294-1301. PMID 31811339
31. Friese G, Herten M, Scherbaum WA. The use of autologous platelet concentrate activated by autologous thrombin (APC+) is effective and safe in the treatment of chronic diabetic foot ulcers-a randomized controlled trial. In: eds. *Proceedings of the Fifth International Symposium on the Diabetic Foot*, May September 12, 2007, Noordwijkerhout, The Netherlands. 2007.
32. Game F, Jeffcoate W, Tarnow L, et al. LeucoPatch system for the management of hard-to-heal diabetic foot ulcers in the UK, Denmark, and Sweden: an observer-masked,

- randomised controlled trial. *Lancet Diabetes Endocrinol.* Nov 2018; 6(11): 870-878. PMID 30243803
33. Gude W, Hagan D, Abood F, et al. Aurix Gel Is an Effective Intervention for Chronic Diabetic Foot Ulcers: A Pragmatic Randomized Controlled Trial. *Adv Skin Wound Care.* Sep 2019; 32(9): 416-426. PMID 31436621
 34. Habeeb T, AA E, H M. Platelet-rich plasma (PRP) bio-stimulant gel dressing in treating chronic non-healing leg and foot ulcers; cost and effectiveness. *Randomized Controlled Clinical Trial.* 2021.
 35. Helmy Y, Farouk N, Ali Dahy A, et al. Objective assessment of Platelet-Rich Plasma (PRP) potentiality in the treatment of Chronic leg Ulcer: RCT on 80 patients with Venous ulcer. *J Cosmet Dermatol.* Oct 2021; 20(10): 3257-3263. PMID 33880860
 36. Hossam EM, Alserr AHK, Antonopoulos CN, et al. Autologous Platelet Rich Plasma Promotes the Healing of Non-Ischemic Diabetic Foot Ulcers. A Randomized Controlled Trial. *Ann Vasc Surg.* May 2022; 82: 165-171. PMID 34896242
 37. Jeong SH, Han SK, Kim WK. Treatment of diabetic foot ulcers using a blood bank platelet concentrate. *Plast Reconstr Surg.* Mar 2010; 125(3): 944-52. PMID 20195121
 38. Kakagia DD, Kazakos KJ, Xarchas KC, et al. Synergistic action of protease-modulating matrix and autologous growth factors in healing of diabetic foot ulcers. A prospective randomized trial. *J Diabetes Complications.* 2007; 21(6): 387-91. PMID 17967712
 39. Karimi R, Afshar M, Salimian M, et al. The effect of platelet rich plasma dressing on healing diabetic foot ulcers. *Nurs Midwifery Stud.* 2016;5(3):e30314.
 40. Li L, Wang C, Wang Y, He LP, Yang YZ, Chen LH, et al. Impact of topical application of autologous platelet-rich gel on medical expenditure and length of stay in hospitals in diabetic patients with refractory cutaneous ulcers. *Sichuan Da Xue Xue Bao Yi Xue Ban.* 2012;43(5):7625
 41. Li L, Chen D, Wang C, et al. Autologous platelet-rich gel for treatment of diabetic chronic refractory cutaneous ulcers: A prospective, randomized clinical trial. *Wound Repair Regen.* 2015; 23(4): 495-505. PMID 25847503
 42. Liu GY, Deng XL, Sun Y, Wang MZ, Gao J, Gou J. Effect of autologous platelet-rich gel on the treatment of diabetic foot ulcers. *J Xi'an Jiaotong Univ (Med Sci).* 2016;37:264-267.
 43. Liao X, Liang JX, Li SH, et al. Allogeneic Platelet-Rich Plasma Therapy as an Effective and Safe Adjuvant Method for Chronic Wounds. *J Surg Res.* Feb 2020; 246: 284-291. PMID 31622885
 44. Meamar R, Ghasemi-Mobarakeh L, Norouzi MR, et al. Improved wound healing of diabetic foot ulcers using human placenta-derived mesenchymal stem cells in gelatin electrospun nanofibrous scaffolds plus a platelet-rich plasma gel: A randomized clinical trial. *Int Immunopharmacol.* Dec 2021; 101(Pt B): 108282. PMID 34737130
 45. Ma L. Clinical efficacy of autologous platelet rich gel in the treatment of diabetic foot and diabetic chronic cutaneous ulcer. *Chin J Mod Drug Appl.* 2014;8:86-88
 46. Miłek T, Baranowski K, Zydlewski P, et al. Role of plasma growth factor in the healing of chronic ulcers of the lower legs and foot due to ischaemia in diabetic patients. *Postepy Dermatol Alergol.* Dec 2017; 34(6): 601-606. PMID 29422826
 47. Qi KQ, Chen TJ, Shang XL. The application of autologous platelet-rich gel in the treatment of diabetic foot ulcers. *Chin J Diabetes.* 2014;22: 1102-1105.
 48. Rainys D, Cepas A, Dambrauskaite K, et al. Effectiveness of autologous platelet-rich plasma gel in the treatment of hard-to-heal leg ulcers: a randomised control trial. *J Wound Care.* Oct 02 2019; 28(10): 658-667. PMID 31600109

49. Saad Setta H, Elshahat A, Elsherbiny K, et al. Platelet-rich plasma versus platelet-poor plasma in the management of chronic diabetic foot ulcers: a comparative study. *Int Wound J*. Jun 2011; 8(3): 307-12. PMID 21470370
50. Saldalamacchia G, Lapice E, Cuomo V, et al. A controlled study of the use of autologous platelet gel for the treatment of diabetic foot ulcers. *Nutr Metab Cardiovasc Dis*. Dec 2004; 14(6): 395-6. PMID 15853123
51. Serra R, Grande R, Butrico L, et al. Skin grafting and topical application of platelet gel in the treatment of vascular lower extremity ulcers. *Acta Phlebologica*. 2014 01 Dec;15(3):129-36.
52. Singh SP, Kumar V, Pandey A, et al. Role of platelet-rich plasma in healing diabetic foot ulcers: a prospective study. *J Wound Care*. Sep 02 2018; 27(9): 550-556. PMID 30204574
53. Steed DL, Goslen JB, Holloway GA, et al. Randomized prospective double-blind trial in healing chronic diabetic foot ulcers. CT-102 activated platelet supernatant, topical versus placebo. *Diabetes Care*. Nov 1992; 15(11): 1598-604. PMID 1468291
54. Steed DL, Edington HD, Webster MW. Recurrence rate of diabetic neurotrophic foot ulcers healed using topical application of growth factors released from platelets. *Wound Repair Regen*. 1996; 4(2): 230-3. PMID 17177818
55. Mohammadi Tofigh A, Tajik M. Comparing the standard surgical dressing with dehydrated amnion and platelet-derived growth factor dressings in the healing rate of diabetic foot ulcer: A randomized clinical trial. *Diabetes Res Clin Pract*. Mar 2022; 185: 109775. PMID 35149167
56. Xie J, Fang Y, Zhao Y, et al. Autologous Platelet-Rich Gel for the Treatment of Diabetic Sinus Tract Wounds: A Clinical Study. *J Surg Res*. Mar 2020; 247: 271-279. PMID 31706541
57. Yang L, Gao L, Lv Y, et al. Autologous platelet-rich gel for lower-extremity ischemic ulcers in patients with type 2 diabetes. *International Journal of Clinical and Experimental Medicine*. 2017 30 Sep;10(9):13796-801.
58. Zhang L Qiang D, Sun YH. Clinical observation of autologous platelet rich gel in the treatment of diabetic foot ulcers. *Ningxia Med J*. 2016;38:809-811.
59. Zhou XP, Gong YX, Yang ZD, Wang W. Application value analysis of autologous platelet gel in refractory skin ulcer of diabetic patients. *World Lat Med Inform*. 2015;15:19-20
60. Zhu SF, Liu H, Li L, Wang XF. Preliminary application of autologous platelet rich gel in diabetic neuropathic ulcers. *Med Innov China*. 2012;9:18-19.
61. Centers for Medicare & Medicaid Services (CMS). National Coverage Analysis (NCA) Tracking Sheet for Autologous Blood-Derived Products for Chronic Non-Healing Wounds (CAG-00190R4). 2020; <https://www.cms.gov/medicare-coverage-database/details/nca-tracking-sheet.aspx?NCAId=300&NCDId=217>. Accessed December 10, 2023.
62. Gupta A, Channaveera C, Sethi S, et al. Efficacy of Intralesional Platelet-Rich Plasma in Diabetic Foot Ulcer. *J Am Podiatr Med Assoc*. May 01 2021; 111(3). PMID 33231614
63. Qu W, Wang Z, Hunt C, et al. The Effectiveness and Safety of Platelet-Rich Plasma for Chronic Wounds: A Systematic Review and Meta-analysis. *Mayo Clin Proc*. Sep 2021; 96(9): 2407-2417. PMID 34226023
64. Oliveira BGRB, Carvalho MR, Ribeiro APL. Cost and effectiveness of Platelet Rich Plasma in the healing of varicose ulcer: Meta-analysis. *Rev Bras Enferm*. 2020; 73(4): e20180981. PMID 32609173
65. Fang Q, Zhang Y, Tang L, et al. Clinical Study of Platelet-Rich Plasma (PRP) for Lower Extremity Venous Ulcers: A Meta-Analysis and Systematic Review. *Int J Low Extrem Wounds*. Dec 2023; 22(4): 641-653. PMID 34665051

66. Saha S, Patra AC, Gowda SP, et al. Effectiveness and safety of autologous platelet-rich plasma therapy with total contact casting versus total contact casting alone in treatment of trophic ulcer in leprosy: An observer-blind, randomized controlled trial. *Indian J Dermatol Venereol Leprol*. 2020; 86(3): 262-271. PMID 31997794
67. Shehab AW, Eleshra A, Fouda E, et al. Randomized prospective comparative study of platelet-rich plasma versus conventional compression in treatment of post-phlebotic venous ulcer. *Vascular*. Dec 2023; 31(6): 1222-1229. PMID 35603798
68. Zhou SF, Estrera AL, Loubser P, et al. Autologous platelet-rich plasma reduces transfusions during ascending aortic arch repair: a prospective, randomized, controlled trial. *Ann Thorac Surg*. Apr 2015; 99(4): 1282-90. PMID 25661906
69. Serraino GF, Dominijanni A, Jiritano F, et al. Platelet-rich plasma inside the sternotomy wound reduces the incidence of sternal wound infections. *Int Wound J*. Jun 2015; 12(3): 260-4. PMID 23692143
70. Zhu S, Gao J, Yu W, et al. Platelet-rich plasma influence on the sternal wounds healing: A meta-analysis. *Int Wound J*. Nov 2023; 20(9): 3794-3801. PMID 37350616
71. El-Anwar MW, Nofal AA, Khalifa M, et al. Use of autologous platelet-rich plasma in complete cleft palate repair. *Laryngoscope*. Jul 2016; 126(7): 1524-8. PMID 27075516
72. Sidman JD, Lander TA, Finkelstein M. Platelet-rich plasma for pediatric tonsillectomy patients. *Laryngoscope*. Oct 2008; 118(10): 1765-7. PMID 18622315
73. Almdahl SM, Veel T, Halvorsen P, et al. Randomized prospective trial of saphenous vein harvest site infection after wound closure with and without topical application of autologous platelet-rich plasma. *Eur J Cardiothorac Surg*. Jan 2011; 39(1): 44-8. PMID 20634084
74. Mohamadi S, Norooznezhad AH, Mostafaei S, et al. A randomized controlled trial of effectiveness of platelet-rich plasma gel and regular dressing on wound healing time in pilonidal sinus surgery: Role of different affecting factors. *Biomed J*. Dec 2019; 42(6): 403-410. PMID 31948604
75. Slaninka I, Fibír A, Kaška M, et al. Use of autologous platelet-rich plasma in healing skin graft donor sites. *J Wound Care*. Jan 02 2020; 29(1): 36-41. PMID 31930949
76. Kazakos K, Lyras DN, Verettas D, et al. The use of autologous PRP gel as an aid in the management of acute trauma wounds. *Injury*. Aug 2009; 40(8): 801-5. PMID 18703188
77. Marck RE, Gardien KL, Stekelenburg CM, et al. The application of platelet-rich plasma in the treatment of deep dermal burns: A randomized, double-blind, intra-patient controlled study. *Wound Repair Regen*. Jul 2016; 24(4): 712-20. PMID 27169627
78. Yeung CY, Hsieh PS, Wei LG, et al. Efficacy of Lyophilised Platelet-Rich Plasma Powder on Healing Rate in Patients With Deep Second Degree Burn Injury: A Prospective Double-Blind Randomized Clinical Trial. *Ann Plast Surg*. Feb 2018; 80(2S Suppl 1): S66-S69. PMID 29369904
79. Huang H, Sun X, Zhao Y. Platelet-rich plasma for the treatment of burn wounds: A meta-analysis of randomized controlled trials. *Transfus Apher Sci*. Feb 2021; 60(1): 102964. PMID 33127309
80. Imam MS, Alotaibi AAS, Alotaibi NOM, et al. Efficiency of platelet-rich plasma in the management of burn wounds: A meta-analysis. *Int Wound J*. Sep 30 2023. PMID 37776166
81. Qaseem A, Humphrey LL, Forciea MA, et al. Treatment of pressure ulcers: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. Mar 03 2015; 162(5): 370-9. PMID 25732279

82. Association for the Advancement of Wound Care (AAWC). Guideline of Pressure Ulcer Guidelines. Malvern, PA: AAWC; 2010.
83. Association for the Advancement of Wound Care (AAWC). International Consolidated Venous Ulcer Guideline (ICVUG). 2015; <https://aawconline.memberclicks.net/assets/appendix%20c%20guideline%20icvug-textformatrecommendations-final%20v42%20changessaved18aug17.pdf>. Accessed December 13, 2023.
84. National Institute for Health and Clinical Excellence (NICE). Diabetic foot problems: prevention and management [NG19]. 2019; <https://www.nice.org.uk/guidance/ng19/resources/diabetic-foot-problems-prevention-and-management-pdf-1837279828933>. Accessed December 13, 2023.
85. Centers for Medicare and Medicaid Services (CMS). National coverage determination (NCD) for blood-derived products for chronic non-healing wounds (270.3). Effective date of version August 2, 2012. <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=217&ncdver=5&NCAId=260>. Accessed December 7, 2023.
86. Centers for Medicare & Medicaid Services (CMS). Decision Memo for Autologous Blood-Derived Products for Chronic Non-Healing Wounds (CAG-00190R3). 2012; <https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=260>. Accessed December 11, 2023.
87. Centers for Medicare & Medicaid Services (CMS). CMS Manual System: Pub 100-3 Medicare National Coverage Determinations (Transmittal 127). 2010 Oct; <https://www.cms.gov/Regulations-and-Guidance/Guidance/Transmittals/downloads/R127NCD.pdf>. Accessed December 13, 2023.
88. Centers for Medicare & Medicaid Services (CMS). Decision Memo for Autologous Blood Derived Products for Chronic Non-Healing Wounds (CAG-00190R2). 2008; <https://www.cms.gov/medicare-coverage-database/details/nca-decision-memo.aspx?NCAId=208>. Accessed December 12, 2023.
89. Qu W, Wang Z, Hunt C, Morrow AS, Urtecho M, Amin M, Shah S, Hasan B, Abd-Rabu R, Ashmore Z, Kubrova E, Prokop LJ, Murad MH. Platelet-Rich Plasma for Wound Care in the Medicare Population. Technology Assessment Program Project ID 040-353-492. (Prepared by the Mayo Clinic Evidence-based Practice Center under Contract No. HHS290201500013I.) Rockville, MD: Agency for Healthcare Research and Quality. <https://www.ahrq.gov/sites/default/files/wysiwyg/research/findings/ta/prp/prp-wound-care.pdf>. Accessed December 13, 2023.
90. Centers for Medicare & Medicaid Services (CMS). National Coverage Analysis (NCA) for Autologous Blood-Derived Products for Chronic Non-Healing Wounds (CAG-00190R4). 2021; <https://www.cms.gov/medicare-coverage-database/view/ncacal-decision-memo.aspx?proposed=N&ncaid=300>. Accessed December 9, 2023.