

Medical Policy



An independent licensee of the
Blue Cross Blue Shield Association

Title: Miscellaneous Genetic and Molecular Diagnostic Tests

Professional

Original Effective Date: January 7, 2016
Revision Date(s): January 7, 2016;
August 17, 2016; January 1, 2017;
October 25, 2017; December 20, 2017;
October 12, 2018; January 1, 2019
Current Effective Date: October 12, 2018

Institutional

Original Effective Date: January 7, 2016
Revision Date(s): January 7, 2016;
August 17, 2016; January 1, 2017;
October 25, 2017; December 20, 2017;
October 12, 2018; January 1, 2019
Current Effective Date: October 12, 2018

State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).

The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.

The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.

If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.

Populations	Interventions	Comparators	Outcomes
Individuals: • With symptoms of various conditions thought to be hereditary or with a known genetic component	Interventions of interest are: • Diagnostic testing with a miscellaneous genetic or molecular test	Comparators of interest are: • Standard care without genetic or molecular diagnostic testing	Relevant outcomes include: • Overall survival • Disease-specific survival • Test accuracy • Test validity • Change in disease status • Morbid events
Individuals: • Who are being screened for colorectal cancer	Interventions of interest are: • Testing with a <i>SEPT9</i> methylated DNA test	Comparators of interest are: • Standard care without <i>SEPT9</i> methylated DNA testing	Relevant outcomes include: • Overall survival • Disease-specific survival • Test accuracy • Test validity • Change in disease status • Morbid events

Populations	Interventions	Comparators	Outcomes
Individuals: • Who are diagnosed with various conditions (eg, Crohn's disease, thymomas and thymic carcinomas, celiac disease)	Interventions of interest are: • Prognostic testing with a miscellaneous genetic or molecular test	Comparators of interest are: • Standard care without genetic or molecular prognostic testing	Relevant outcomes include: • Overall survival • Disease-specific survival • Test accuracy • Test validity • Change in disease status • Morbid events
Individuals: • Who are diagnosed with stage I or II melanoma	Interventions of interest are: • Prognostic testing with DecisionDx-Melanoma	Comparators of interest are: • Standard care without DecisionDx-Melanoma testing	Relevant outcomes include: • Overall survival • Disease-specific survival • Test accuracy • Test validity • Change in disease status • Morbid events
Individuals: • Who are diagnosed with various conditions (eg, Crohn's disease, thymomas and thymic carcinomas, rheumatoid arthritis)	Interventions of interest are: • Prognostic testing with a miscellaneous genetic or molecular test	Comparators of interest are: • Standard care without genetic or molecular therapeutic testing	Relevant outcomes include: • Overall survival • Disease-specific survival • Test accuracy • Test validity • Change in disease status • Morbid events
Individuals: • Who are diagnosed with various conditions (eg, colon cancer, non-Hodgkin lymphoma)	Interventions of interest are: • Therapeutic testing with a miscellaneous genetic or molecular test	Comparators of interest are: • Standard care without genetic or molecular therapeutic testing	Relevant outcomes include: • Overall survival • Disease-specific survival • Test accuracy • Test validity • Change in disease status • Morbid events
Individuals: • With a family history of various conditions thought to be hereditary or with a known genetic component	Interventions of interest are: • Testing for future risk of disease with a miscellaneous genetic or molecular test	Comparators of interest are: • Standard care without genetic or molecular diagnostic testing for future risk	Relevant outcomes include: • Overall survival • Disease-specific survival • Test accuracy • Test validity • Change in disease status • Morbid events

DESCRIPTION

There are numerous commercially available genetic and molecular diagnostic, prognostic, and therapeutic tests for individuals with certain diseases or asymptomatic individuals with a future risk. This policy evaluates miscellaneous genetic and molecular diagnostic tests not addressed in a separate review. If a separate policy exists, then conclusions reached there supersede conclusions in this review. The main criterion for inclusion in this review is that there is limited evidence on the clinical validity for the test. As a result, these tests do not have clinical utility, and the evidence is insufficient to determine the effect on health outcomes.

OBJECTIVE

The objective of this policy is to determine whether diagnostic, prognostic, therapeutic, or future risk assessment testing using one of several miscellaneous genetic or molecular diagnostic tests improves the net health outcome in individuals with, or with a risk of, one of various genetic conditions.

BACKGROUND

Tests Addressed in This Policy

Tests assessed in this policy are listed in Table 1. Three types of tests are related to testing of an affected (symptomatic) individual's germline to benefit the individual (excluding reproductive testing): diagnostic testing, prognostic testing, and therapeutic testing. The fourth type of test reviewed is testing of an asymptomatic individual to determine future risk of disease.

Table 1: Genetic and Molecular Diagnostic Tests in This Policy

Test Name	Manufacturer	Date Added	Diagnostic	Prognostic	Therapeutic	Future Risk
Celiac PLUS	Prometheus	Oct 2014	•			•
ColonSentry®	GeneNews ^a	Aug 2015				•
Crohn's Prognostic	Prometheus	Oct 2014		•		
DecisionDx-Thymoma	Castle	Jan 2015		•		
DNA Methylation Pathway Profile	Great Plains Laboratory	Jan 2015	•			
GI Effects® (Stool)	Genova Dxcs	Jan 2015	•			
IBD sgi Diagnostic™	Prometheus	Oct 2014	•			
ImmunoGenomic® Profile	Genova Dxcs	Aug 2015				•
Know Error™	Strand Dxcs	July 2016	•			
ResponseDX®: Colon	Response Gxcs	Jan 2015			•	
SEPT9 methylated DNA ^b	Severalc	Oct 2014	•			
TransPredict Fc gamma 3A	Transgenomic	Oct 2014			•	

Castle: Castle Biosciences; Dxcs: Diagnostics; Gxcs: Genetics.

^a In a joint venture with Innovative Diagnostic Laboratory.

^b For example, ColoVantage®, Epi proColon®.

^c ARUP, Quest, Clinical Genomics, Epigenomics.

Diagnostic Tests

Multiple Conditions

Single-nucleotide polymorphisms (SNPs) are the most common type of genetic variation, and each SNP represents a difference in a single nucleotide in the DNA sequence. Most commonly, SNPs are found in the DNA between genes and can act as biological markers of genes and disease association. When SNPs occur within a gene or a gene regulatory region, they can play a more direct role in disease by affecting the gene's function. SNPs may predict an individual's response to certain drugs, susceptibility to environmental factors, and the risk of developing certain diseases.

DNA specimen provenance assays can be used to confirm that tissue specimens are correctly matched to the patient of origin. Specimen provenance errors may occur in up to 1% to 2% of pathology tissue specimens,¹ and have serious negative implications for patient care if the error is not corrected.² Analysis of DNA microsatellites from tissue specimens can be performed by analyzing long tandem repeats (LTR), and comparing the LTRs of the tissue specimen to LTRs from a patient sample.

Test Description: DNA Methylation Pathway Profile

The DNA Methylation Pathway Profile (Great Plains Laboratory, Lenexa, KS) analyzes SNPs associated with certain biochemical processes, including methionine metabolism,

detoxification, hormone imbalances, and vitamin D function. Intended uses for the test include clarification of a diagnosis suggested by other testing and as an indication for supplements and diet modifications.

Test Description: Know Error DNA Specimen Provenance Assay

The Know Error test (Strand Diagnostics, Indianapolis, IN) compares the LTRs of tissue samples with LTRs from a buccal swab of the patient. The intended use of the test is to confirm tissue of origin and avoid specimen provenance errors due to switching of patient samples, mislabeling, or sample contamination.

Celiac Disease

Celiac disease (previously called sprue, celiac sprue, gluten-sensitive enteropathy, gluten intolerance, nontropical sprue, idiopathic steatorrhea) is an immune-based reaction to gluten (water insoluble proteins in wheat, barley, rye) that primarily affects the small intestine. Celiac disease occurs almost exclusively in patients who carry at least 1 human leukocyte antigen (HLA) DQ2 or DQ8 allele; negative predictive value (NPV) of having neither allele exceeds 98%.³ Serum antibodies to tissue transglutaminase (TTG), endomysium, and deamidated gliadin peptide (DGP) support a diagnosis of celiac disease, but diagnostic confirmation requires duodenal biopsy taken when patients are on a gluten-containing diet.⁴

Test Description: Celiac PLUS

Celiac PLUS (Prometheus Therapeutics & Diagnostics, San Diego) is a panel of 2 genetic and 5 serologic markers associated with celiac disease. Per the manufacturer, Celiac PLUS is a diagnostic test that also stratifies future risk of celiac disease.⁵ Genetic markers, HLA DQ2 and DQ8, are considered predictive of the risk of developing celiac disease⁶; serologic markers—immunoglobulin A (IgA) anti-TTG antibody, IgA anti-endomysial antibodies, IgA anti-DGP antibodies, IgG anti-DGP, and total IgA—are considered diagnostic for celiac disease. Celiac PLUS is intended for patients at risk for disease (eg, with an affected first-degree relative) or with symptoms suggestive of disease.

Irritable Bowel Syndrome

Irritable bowel syndrome (IBS) is a functional gastrointestinal (GI) disorder that affects 10% to 20% of the general population in the United States and worldwide. Symptoms include abdominal pain and/or bloating associated with disordered bowel habit (constipation, diarrhea, or both). Pathophysiology is poorly understood but may be related to chronic low-grade mucosal inflammation and disturbances in GI flora.⁷ Recommended treatments include dietary restriction and pharmacologic symptom control.⁸⁻¹⁰ Probiotics—living microorganisms that promote health when administered to a host in therapeutic doses¹¹—are being investigated as a treatment for IBS. Several systematic reviews of randomized controlled trials (RCTs) have found evidence to support efficacy,^{7,12-15} but results from recent RCTs have been mixed.¹⁶⁻²¹ This

discrepancy may be due in part to differential effects of different probiotic strains and doses.

Test Description: GI Effects Comprehensive Stool Profile

The GI Effects Comprehensive Stool Profile (Genova Diagnostics, Asheville, NC) is a multianalyte stool assay.²² The test uses polymerase chain reaction (PCR) to quantify 26 commensal gut bacteria, and standard biochemical and culture methods to measure levels of other stool components (eg, lipids, fecal occult blood) and potential pathogens (ova and parasites, opportunistic bacteria, yeast). The test is purported to optimize management of gut health and to differentiate IBS from inflammatory bowel disease (IBD).

Inflammatory Bowel Disease

IBD is an autoimmune condition characterized by inflammation of the bowel wall, and clinical symptoms of abdominal pain, diarrhea and associated symptoms. Crohn's disease (CD) and ulcerative colitis (UC) are the 2 main entities under the category of IBD. The diagnosis is typically made by endoscopy or colonoscopy with biopsy and histologic analysis. This requires a semi-invasive procedure; as a result, a blood test to diagnose IBD could avoid the need for the procedures.

Test Description: IBD sgi Diagnostic

IBD sgi Diagnostic (Prometheus Therapeutics & Diagnostics, San Diego, CA) is a panel of 17 serologic (n=8), genetic (n=4), and inflammatory biomarkers (n=5). A proprietary algorithm produces an IBD score; results are reported as consistent with IBD (consistent with UC, consistent with CD, or inconclusive for UC vs CD) or not consistent with IBD. The test is intended for use in patients with clinical suspicion of IBD.

Colon Cancer

Early detection of colorectal cancer (CRC) reduces disease-related mortality, yet many individuals do not undergo recommended screening with fecal occult blood test or colonoscopy. It is thought that a simpler screening blood test may encourage screening and decrease mortality, although this has not been proved. Serum biomarkers that are shed from colorectal tumors have been identified and include Septin 9 hypermethylated DNA (*SEPT9*). Septin 9 protein is involved in cell division, migration, and apoptosis, and acts as a tumor suppressor; when hypermethylated, expression of *SEPT9* is reduced.

A cofounder of the biotechnology firm GeneNews developed a patented platform technology based on the sentinel principle.²³ The sentinel principle posits that because blood interacts with all bodily tissues, "subtle changes occurring in association with injury or disease, within the cells and tissues of the body, may trigger specific changes in gene expression in blood cells reflective of the initiating stimulus."²³ In this way, blood cells (specifically, leukocytes) may act as sentinels of disease. In studies that led to the formulation of this principle, investigators compared gene expression (total RNA levels) in blood samples with cataloged genes from 9 different organs (brain, colon,

heart, kidney, liver, lung, prostate, spleen, stomach) and estimated that 66% to 82% of genes encoded in the human genome are expressed in human leukocytes.²³

Test Descriptions: SEPT9 Methylated DNA

ColoVantage (various manufacturers) blood tests for serum *SEPT9* methylated DNA are offered by several laboratories (ARUP Laboratories, Quest Diagnostics, Clinical Genomics). Epi proColon (Epigenomics, Berlin) received FDA approval in the United States in April 2016. Epigenomics has licensed its Septin 9 DNA biomarker technology to Polymedco and LabCorp. ColoVantage and Epi proColon are both PCR assays; however, performance characteristics vary across tests, presumably due to differences in methodology (eg, DNA preparation, PCR primers, probes).

Test Description: ColonSentry

ColonSentry (GeneNews; Innovative Diagnostic Laboratory) is a PCR assay that uses a blood sample to detect expression of 7 genes found to be differentially expressed in CRC patients compared with controls²⁴: *ANXA3*, *CLEC4D*, *TNFAIP6*, *LMNB1*, *PRRG4*, *VNN1*, and *IL2RB*. Per the company website, these genes are early-warning signs of colon cancer, and test results can indicate the odds of having CRC compared with an average-risk person.²⁵ An average-risk person is defined as one who is "at least 50 years old, is asymptomatic for CRC, has no personal history of benign colorectal polyps, colorectal adenomas, CRC, or inflammatory bowel disease, and does not have a first-degree relative with CRC."²⁵ The test is intended for use in adults who are averse to colonoscopy and/or fecal occult blood testing. "Because of its narrow focus, the test is not expected to alter clinical practice for patients who comply with recommended screening schedules."²⁶

Prognostic Tests

Crohn's Disease

Recent studies have identified serologic²⁷ and genetic^{28,29} correlates of aggressive CD that is characterized by fistula formation, fibrostenosis, and the need for surgical intervention. Prometheus has developed a blood test that aims to identify patients with CD who are likely to experience an aggressive disease course.

Test Description: Crohn's Prognostic

Crohn's Prognostic (Prometheus Therapeutics & Diagnostics, San Diego, CA) is a panel of 6 serologic (n=3) and genetic (n=3) biomarkers. Limited information about the test is available on the manufacturer's website.

Thymomas and Thymic Carcinomas

Thymomas and thymic carcinomas are rare epithelial tumors of the thymus. Most are diagnosed in individuals between 40 and 60 years of age. Thymic epithelial tumors range from histologically benign tumors to microscopically or macroscopically invasive low- or high-grade malignant tumors. However, even tumors that are histologically benign can behave aggressively.

Test Description: DecisionDx-Thymoma

DecisionDx-Thymoma (Castle Biosciences, Friendswood, TX) is a gene expression profile test that measures the activity of 23 genes within the thymic tumor. Its intended use is to distinguish between thymic carcinoma and thymoma, and to predict tumor aggressiveness by likelihood that the tumor will metastasize.

Therapeutic Tests

Test Description: ResponseDX: Colon

Response Genetics currently markets 2 colon cancer genetic panels to guide treatment selection, as well as separate tests for 11 genes associated with colon cancer prognosis and/or treatment response. The Driver Profile panel comprises PCR mutation testing in *KRAS*, *BRAF*, and mismatch repair genes (microsatellite instability), plus *NRAS* exon 2 and 3 sequencing. These gene tests are reviewed elsewhere, and this panel is not considered here. The ResponseDX: Colon test comprises the 4 tests in the Driver Profile plus: *EGFR* expression; *PI3K* exon 1, 9, and 20 sequencing; *TS* expression; *ERCC1* expression; *UGT1A1* SNP testing (rs8175347, rs4148323); *VEGFR2* expression; and *MET* amplification by fluorescence in situ hybridization.

Non-Hodgkin Lymphoma

Rituximab is a humanized IgG monoclonal antibody against the CD20 antigen, which is commonly expressed on B lymphocytes. It is FDA-approved for treatment of non-Hodgkin lymphoma, chronic lymphocytic leukemia, and nononcologic uses (eg, rheumatoid arthritis).³⁰ Although rituximab has demonstrated improved response and survival rates in combination chemotherapy regimens in patients with follicular lymphoma, chronic lymphocytic leukemia, and diffuse large B-cell lymphoma than chemotherapy alone (not all patients responded). Altered binding to lymphocyte-bound rituximab by cytotoxic effector cells (eg, natural killer cells, macrophages) has been identified as a mechanism of reduced rituximab efficacy. Effector cells with a Val158Phe substitution mutation in their surface receptors for IgG molecules (eg, rituximab) have impaired binding affinity, and cellular cytotoxicity is reduced. A genetic test for the Val158Phe mutation of the gene that encodes the IgG receptor on effector cells (*FCGR3A*) has been developed and investigated as a means of predicting response to rituximab.

Future Risk in Asymptomatic Individuals

Immunologic Disorders

Test Description: ImmunoGenomic Profile

The ImmunoGenomic Profile (Genova Diagnostics) is a buccal swab test that evaluates SNVs in 6 genes associated with immune function and inflammation: interleukin (IL)-10, IL-13, IL-1 β , IL-4, IL-6, and tumor necrosis factor α .³¹ According to the company website, variations in these genes "can affect balance between cell (TH-1) and humoral (TH-2) immunity, trigger potential defects in immune system defense, and stimulate mechanisms underlying chronic, overactive inflammatory responses... The test uncovers potential genetic susceptibility to: Asthma, Autoimmune Disorders, Certain

Cancers, Allergy, Infectious Diseases, Bone Inflammation, Arthritis, Inflammatory Bowel Disease, Heart Disease, Osteopenia, and Helicobacter pylori infection (cause of ulcers).”

REGULATORY STATUS

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests (LDTs) must meet the general standards of the Clinical Improvement Amendments (CLIA). Genetic tests reviewed in this policy are available under the auspices of CLIA. Laboratories that offer LDTs must be licensed by CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of these tests.

POLICY

- A. All of the tests listed in this policy are considered **experimental / investigational** and grouped according to the categories of genetic testing:
1. Testing of an affected (symptomatic) individual's germline to benefit the individual (excluding reproductive testing)
 2. Diagnostic testing
 3. Prognostic testing
 4. Therapeutic testing
 5. Testing an asymptomatic individual to determine future risk of disease

Policy Guidelines

Genetic Counseling

Genetic counseling is primarily aimed at patients who are at risk for inherited disorders, and experts recommend formal genetic counseling in most cases when genetic testing for an inherited condition is considered. The interpretation of the results of genetic tests and the understanding of risk factors can be very difficult and complex. Therefore, genetic counseling will assist individuals in understanding the possible benefits and harms of genetic testing, including the possible impact of the information on the individual's family. Genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing. Genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

RATIONALE

The literature review was performed through May 7, 2018.

Evidence reviews assess whether a medical test is clinically useful. A useful test provides information to make a clinical management decision that improves the net health outcome. That is, the balance of benefits and harms is better when the test is used to manage the condition than when another test or no test is used to manage the condition.

The first step in assessing a medical test is to formulate the clinical context and purpose of the test. The test must be technically reliable, clinically valid, and clinically useful for that purpose. Evidence reviews assess the evidence on whether a test is clinically valid and clinically useful. Technical reliability is outside the scope of these reviews, and credible information on technical reliability is available from other sources.

Diagnostic Testing for Multiple Conditions

Clinical Context and Test Purpose

The purpose of diagnostic testing in patients for heritable or genetic pathogenic variants in a symptomatic individual is to establish a molecular diagnosis defined by the presence of known pathologic variant(s). For genetic testing, a symptomatic individual is defined as an individual with a clinical phenotype that correlates with a known pathologic variant.

The question addressed in this evidence review is: Does diagnostic testing for heritable or genetic pathogenic variants using the tests described below in symptomatic individuals improve the net health outcome?

The specific clinical context of each test is described briefly in the following sections. The following PICOTS were used to select literature to inform this review.

Patients

The relevant population of interest is patients with symptoms of a particular disease for which a definitive diagnosis cannot be made using other diagnostic methods.

Interventions

The interventions of interest are miscellaneous genetic or molecular diagnostic tests, specifically: DNA Methylation Pathway Profile, Know Error, Celiac PLUS, GI Effects (Stool), and IBD sgi Diagnostic.

Comparators

The comparator of interest is standard care without genetic or molecular diagnostic testing.

Outcomes

The outcomes of interest are overall survival, disease-specific survival, test accuracy and validity, change in disease status, and morbid events.

Timing

The timing of follow-up for irritable bowel syndrome, inflammatory bowel disease, and celiac disease ranges from weeks for the diagnosis to years for assessment of health outcomes.

Setting

These tests are offered commercially through various manufacturers.

Technically Reliable

Assessment of technical reliability focuses on specific tests and operators and requires review of unpublished and often proprietary information. Review of specific tests, operators, and unpublished data are outside the scope of this evidence review and alternative sources exist. This evidence review focuses on the clinical validity and clinical utility.

Diagnostic Testing for Multiple Conditions: DNA Methylation Pathway Profile

Clinically Valid: A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

No full-length, peer-reviewed studies of the DNA Methylation Pathway Profile were identified.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence: Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

Direct evidence for clinical utility is lacking.

Chain of Evidence: Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

It is not possible to construct a chain of evidence for clinical utility due to the lack of clinical validity.

Section Summary: DNA Methylation Pathway Profile

No studies were identified that evaluated this test.

Diagnostic Testing for Multiple Conditions: Know Error Specimen Provenance AssayClinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Evidence for the clinical validity of the Know Error Specimen Provenance Assay is lacking. There is some evidence on the application of short tandem repeat testing for specimen provenance assays in general,³² but these data are not specific to the Know Error test.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence: Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

Direct evidence for clinical utility is lacking.

Chain of Evidence: Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

It is not possible to construct a chain of evidence for clinical utility due to the lack of clinical validity.

Section Summary: Know Error Specimen Provenance Assays

There is a lack of published evidence on the use of the Know Error test to confirm the tissue of origin. Studies are needed that compare the use of Know Error with standard laboratory quality measures and that demonstrate a reduction in specimen provenance errors associated with the use of Know Error.

Diagnostic Testing for Celiac Disease: Celiac PLUS

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Celiac PLUS tests for genetic and serologic factors known to be associated with celiac disease. All 7 test components are included in an evidence-based diagnostic algorithm developed by the American College of Gastroenterology.³³ However, algorithmic testing is individualized according to baseline risk of disease and is done sequentially, rather than simultaneously as in Celiac PLUS.

No studies of the combined serologic and genetic Celiac PLUS test were identified. Information about clinical validity of obtaining several serologic and genetic tests at once (ie, Celiac PLUS) is lacking; improved sensitivity and reduced specificity may be expected.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence: Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

No studies examining the clinical utility of Celiac PLUS were identified.

Chain of Evidence: Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Factors that support a chain of evidence for clinical utility are lacking. A comparison of clinical and/or histopathologic outcomes using either Celiac PLUS or ACG's published diagnostic algorithm would be required to demonstrate improved health outcomes with Celiac PLUS.

Section Summary: Celiac Disease

No studies examining the clinical utility of Celiac PLUS were identified. Factors that support a chain of evidence for prognostic or diagnostic utility are lacking.

Diagnostic Testing for Irritable Bowel Syndrome: GI Effects Comprehensive Stool Profile*Clinically Valid*

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

No studies were identified that assessed the accuracy of the GI Effects fecal panel for diagnosing IBS or for documenting "gut health," a concept that may be difficult to define given large interindividual variability in gut flora.³⁴

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence: Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

Clinical trials demonstrating a net health benefit with the GI Effects fecal panel were not identified.

Chain of Evidence: Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because probiotics are not currently a standard treatment of IBS, the impact of test results on disease management is uncertain; ie, a chain of evidence for clinical utility of the test cannot be established.

Section Summary: Irritable Bowel Syndrome

Evidence for the clinical validity and utility of the GI Effects Comprehensive Stool Profile is lacking.

Diagnostic Testing for Inflammatory Bowel Disease: IBD sgi Diagnostic

The IBD sgi Diagnostic product monograph includes an extensive bibliography that documents associations of the 18 component markers, individually and in combination, with ulcerative colitis and/or Crohn's disease (CD).³⁵

In a review of the monograph, Shirts et al (2012)³⁶ observed that serologic tests for ASCA-IgA, ASCA-IgG, and atypical perinuclear anti-neutrophil cytoplasmic antibody are standard of care in the diagnostic workup of IBD,^{37,38} although not all investigators include these tests in recommended diagnostic strategies.³⁹⁻⁴² These 3 markers are included in the 18-marker panel. Based on a 2006 meta-analysis of 60 studies (total N=11,608 patients), Reese et al (2006) reported that pooled sensitivity and specificity of the 3-test panel were 63% and 93%,

respectively, for diagnosing IBD.⁴³ Because the product monograph did not compare the 18-marker panel with the 3-marker panel, incremental improvement in diagnosis with the 18-marker panel is unknown. Shirts et al (2012) calculated an area under the curve (AUC) for the 3-marker panel of 0.899.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Published evidence supports associations of each marker in the 18-marker panel, alone and in combination, with IBD diagnosis. Based on manufacturer data, the accuracy for IBD diagnosis of the 18-marker panel exceeds that of each component marker, but the relevant comparison—with a panel of 3 markers that has good discrimination for IBD—was not included; subsequent analysis has suggested that the panels may perform similarly. Performance characteristics for the 18-marker panel to distinguish ulcerative colitis from CD were not provided.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence: Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

No studies examining the clinical utility of IBD sgi Diagnostic were identified.

Chain of Evidence: Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility. It is not possible to construct a chain of evidence for clinical utility due to the lack of clinical validity.

Section Summary: Inflammatory Bowel Disease

No studies examining the clinical utility of IBD sgi Diagnostic were identified. Although manufacturer data supported the clinical validity of the test for diagnosing IBD, this evidence is insufficient to support a chain of evidence for clinical utility. For distinguishing ulcerative colitis from CD, clinical validity has not been established; therefore, a chain of evidence for clinical utility for this purpose cannot be established.

Diagnostic Testing for Colorectal Cancer Screening

The U.S. Preventive Services Task Force has recommended screening for colorectal cancer (CRC) starting at age 50 years and continuing until age 75 years, but many adults do not receive screening for CRC.⁴⁴ It is thought that less burdensome methods of screening could increase the number of adults screened and thereby improve outcomes.

Clinical Context and Test Purpose

The purpose of diagnostic testing in patients with potential CRC is to establish a molecular diagnosis defined by the presence of a known pathologic variant(s).

The question addressed in this evidence review is: Does CRC screening using the tests described below in individuals diagnosed with a disease improve the net health outcome?

The specific clinical context of each test is described briefly in the following sections. The following PICOTS were used to select literature to inform this review.

Patients

The relevant population of interest is patients who are being screened for CRC.

Intervention

The intervention of interest is *SEPT9* methylated DNA testing (eg, ColoVantage, Epi proColon, ColonSentry).

Comparators

The comparator of interest is standard of care without genetic screening.

Outcomes

The outcomes of interest are overall survival, disease-specific survival, test accuracy and validity, change in disease status, and morbid events.

Timing

The timing of follow-up for CRC screening is weeks for the diagnosis of CRC to years for survival outcomes.

Setting

These tests are offered commercially through various manufacturers.

Technically Reliable

Assessment of technical reliability focuses on specific tests and operators and requires review of unpublished and often proprietary information. Review of specific tests, operators, and unpublished data are outside the scope of this evidence review and alternative sources exist. This evidence review focuses on the clinical validity and clinical utility.

Diagnostic Testing for Colorectal Cancer Screening: *SEPT9* Methylated DNA With ColoVantage and Epi proColon

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

The diagnostic performance of *SEPT9* methylation for colon cancer has been reported in meta-analyses. The systematic reviews identified from 2016 and 2017 included from 14 to 39 studies (see Table 2). Pooled sensitivity ranged from 62% to 71% and pooled specificity ranged from 91% to 93% (see Table 3). The systematic review by Nian et al (2017) found that study designs (case-control vs cross-sectional), assays or kits used (Epi proColon vs other), country

(Asia or other), sample sizes (>300 or <300), and risk of bias of included studies all contributed to heterogeneity.⁴⁵ Most included studies were case-control with the exclusion of difficult to diagnose patients, which may lead to a spectrum bias and overestimation of diagnostic accuracy. Reviewers included 20 studies of Epi proColon test 1.0, 2.0, or a combination of the two. When only looking at studies of Epi ProColon 2.0, sensitivity was 75% compared with 71% in the overall analysis, with a specificity of 93% (see Table 3). Sensitivity and specificity may be additionally affected by the specific algorithm used, with the 1/3 algorithm resulting in higher sensitivity and the 2/3 algorithm resulting in higher specificity.⁴⁶

Table 2. Systematic Review Characteristics

Study	Studies Included	N	Study Designs Included	Study Reference Standards Included	11-Item QUADAS Quality Assessment		
					No Domains	1-2 Domains	>2 Domains
Nian et al (2017) ⁴⁵	25	9927	CC and CS	Colonoscopy	3	14	1
Li et al (2016) ⁴⁷	39			Colonoscopy	6	12	21
Yan et al (2016) ⁴⁸	14	9870	CC and CS	Colonoscopy	0	13	1

CC: case-control; CS: cross-sectional.

Table 3. Systematic Review Results

Study	Test	Sensitivity (95% CI), %	Specificity (95% CI), %
Nian et al (2017) ⁴⁵	Various	71 (67 to 75)	92 (89 to 94)
Nian et al (2017) ⁴⁵	Epi Procolon 2.0	75 (67 to 77)	93 (88 to 96)
Li et al (2016) ⁴⁷	Various	62 (56 to 67)	91 (89 to 93)
Yan et al (2016) ⁴⁸	Various	66 (64 to 69)	91 (90 to 91)
Yan et al (2016) ⁴⁸	Epi Procolon	63 (58 to 67)	91 (90 to 92)

CI: confidence interval.

The Epi proColon test is the only *SEPT9* DNA test that has received U.S. Food and Drug Administration approval. It was approved in 2016 for use in average-risk patients who decline other screening methods.

The evidence review for the 2016 U.S. Preventive Services Task Force update on CRC screening included studies on blood tests for methylated *SEPT9* DNA. The inclusion criteria were fair- or good-quality English-language studies, asymptomatic screening populations, age of 40 years or older, and at average risk for CRC or not selected for inclusion based on CRC risk factors. The only study on *SEPT9* found to meet these inclusion criteria was PRESEPT (described below).

PRESEPT (Church et al [2014]) was an international prospective screening study of the first-generation Epi proColon test (see Table 4).⁴⁹ Of 1516 patients selected for laboratory analysis, colonoscopy identified 53 (3%) patients with invasive adenocarcinoma, 315 (21%) with advanced adenoma, and 210 (14%) with nonadvanced adenoma. The overall sensitivity, specificity, positive predictive value, and negative predictive value for the detection of invasive adenocarcinoma are shown in Table 5. Sensitivity for any adenoma was 48% and advanced adenoma was 11%.

Table 4. Study Characteristics

Study	Study Population	Design	Reference Standard	Timing of Reference and Index Tests	Blinding of Assessors
Church et al (2014) ⁴⁹	Patients ≥50 y at average risk and scheduled for colonoscopy	Prospective random sampling from 7941 patients at 32 sites	Colonoscopy	6-16 d before colonoscopy	Yes

Table 5. Study Results

Study	Initial N	Final N	Excluded Samples	Clinical Validity (95% Confidence Interval), %			
				Sensitivity	Specificity	PPV	NPV
Church et al (2014) ⁴⁹	1516	1510	6	48.2 (32.4 to 63.6)	91.5 (89.7 to 93.1)	5	100

NPV: negative predictive value; PPV: positive predictive value.

The purpose of the gaps tables (see Tables 6 and 7) is to display notable gaps identified in each study. This information is synthesized as a summary of the body of evidence following each table and provides the conclusions on the sufficiency of evidence supporting the position statement.

Table 6. Relevance Gaps

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Church et al (2014) ⁴⁹		3. First-generation test			

The evidence gaps stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.
^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest.

^c Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

^d Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

^e Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true positives, true negatives, false positives, false negatives cannot be determined).

Table 7. Study Design and Conduct Gaps

Study	Selection ^a	Blinding ^b	Delivery of Test ^c	Selective Reporting ^d	Data Completeness ^e	Statistical ^f
Church et al (2014) ⁴⁹	2. Not randomly sampled					

The evidence gaps stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Selection key: 1. Selection not described; 2. Selection not random or consecutive (ie, convenience).

^b Blinding key: 1. Not blinded to results of reference or other comparator tests.

^c Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

^d Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^e Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

^f Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison to other tests not reported

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence: Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

Studies comparing survival outcomes in patients who undergo CRC screening with *SEPT9* methylated DNA testing or with standard screening were not identified. Such comparative studies with clinically meaningful outcomes (eg, survival) are necessary to demonstrate incremental improvement in the net health outcome compared with current standard screening approaches (fecal immunochemical test, colonoscopy) and to address lead-time bias for cancers identified through screening.

Chain of Evidence: Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility. Because the sensitivity of *SEPT9* methylated DNA is low, a chain of evidence establishing the clinical utility of *SEPT9* methylated DNA cannot be established.

Subsection Summary: Colorectal Cancer Screening with SEPT9 Methylated DNA Testing

The evidence for clinical validity of CRC screening includes case-control studies and prospective screening studies. Systematic reviews have reported that the sensitivity of testing ranges from 62% to 75% and the specificity from 91% to 93%. Studies were generally of low to fair quality. The prospective PRESEPT study with average-risk patients scheduled for colonoscopy estimated the sensitivity and specificity of Epi proColon for detection of invasive adenocarcinoma to be 48% and for an advanced adenoma to be 11%. Based on results from these studies, the clinical validity of *SEPT9* methylated DNA screening is limited by low sensitivity and low positive predictive value of the test.

Detection of only half of preclinical cancers and a small proportion of advanced adenomas limits clinical utility of the test. There is a need for further studies comparing survival outcomes in patients screened with *SEPT9* methylated DNA testing (ColoVantage, Epi proColon) and with other screening methods. Such comparative studies with clinically meaningful outcomes (eg, survival) are necessary to demonstrate improvement in the net health outcome compared with current standard screening approaches (fecal immunochemical test, colonoscopy) and to address lead-time bias for cancers identified through screening. Because the evidence on clinical validity has reported that the test has a lower sensitivity than other screening methods, the clinical utility is uncertain. If the test is restricted only to patients who would otherwise not be screened, outcomes might be improved. However, if the test is used as a substitute for other screening tests that have higher sensitivity, outcomes may be worse.

Diagnostic Testing for Colorectal Cancer Screening with ColonSentry

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Two case-control studies have been identified. Marshall et al (2010) conducted a genome-wide association study in 189 whole blood samples (98 controls, 91 patients with CRC) and identified 45 differentially expressed gene biomarker candidates using microarray hybridization.⁵⁰ Through logistic regression and bootstrapping (subsampling with replacement) in a training set of 232 samples, 7 genes were selected for further development. In a subsequent test set of 410 samples (208 controls, 202 patients with CRC), sensitivity, specificity, PPV, and NPV were determined (see Tables 8 and 9). Yip et al (2010) conducted a similar cross-sectional study of 210 blood samples from patients in Malaysia.²⁴ The Malaysian population has different ethnic groups with different CRC incidences and CRC in Asian populations is more likely to be nonpolypoid (ie, flat or depressed) compared with Western populations in whom the test was developed.

Sensitivity for the 2 studies ranged from 61% to 72% and specificity for detecting CRC were 70% to 77%. AUC was 0.76 (95% CI, 0.70 to 0.82).

Table 8. Study Characteristics

Study	Study Population	Design	Reference Standard	Timing of Reference and Index Tests
Marshall et al (2010) ⁵⁰	202 patients with CRC and 208 controls	Case-control	NA	NA
Yip et al (2010) ²⁴	99 patients with CRC and 111 controls	Case-control	NA	NA

NA: not applicable

Table 9. Study Results

Study	Initial N	Final N	Excluded Samples	AUC (95% CI)	Clinical Validity (95% Confidence Interval), %			
					Sensitivity	Specificity	PPV	NPV
Marshall et al (2010) ⁵⁰	410			0.80 (0.76 to 0.84)	72	70	70	72
Yip et al (2010) ²⁴	200				61	77		

AUC: area under the curve; NPV: negative predictive value; PPV: positive predictive value.

Gaps in relevance and design and conduct are shown in Tables 10 and 11. Because of its cross-sectional design, follow-up of controls to determine which strata developed CRC was not reported, limiting conclusions drawn about the accuracy of the test for risk prediction.

Table 10. Relevance Gaps

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Marshall et al (2010) ⁵⁰	4. Included patients with CRC and healthy controls				
Yip et al (2010) ²⁴	4. Included patients with CRC and healthy controls				

The evidence gaps stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. CRC: colorectal cancer.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest.

^c Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

^d Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity and predictive values); 4. Reclassification of diagnostic or risk

categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

^e Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true positives, true negatives, false positives, false negatives cannot be determined).

Table 11. Study Design and Conduct Gaps

Study	Selection ^a	Blinding ^b	Delivery of Test ^c	Selective Reporting ^d	Data Completeness ^e	Statistical ^f
Marshall et al (2010) ⁵⁰	2. Selection not random					
Yip et al (2010) ²⁴	2. Selection not random					

The evidence gaps stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Selection key: 1. Selection not described; 2. Selection not random or consecutive (ie, convenience).

^b Blinding key: 1. Not blinded to results of reference or other comparator tests.

^c Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

^d Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^e Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

^f Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison to other tests not reported.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence: Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

No studies examining the clinical utility of ColonSentry were identified.

Chain of Evidence: Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

A chain of evidence supporting the use of ColonSentry for predicting CRC risk cannot be constructed due to lack of clinical validity.

Section Summary: Colorectal Cancer Screening with ColonSentry

ColonSentry is intended to stratify patients with average CRC risk who are averse to current screening approaches to identify those at increased risk and therefore choose a less-invasive screening method. However, 2 cross-sectional studies are insufficient to demonstrate the risk predictive ability of the test; ie, clinical validity has not been established. Direct and indirect evidence of clinical utility is currently lacking.

Prognostic Testing

Clinical Context and Test Purpose

The purpose of prognostic testing of diagnosed disease is to predict natural disease course (eg, aggressiveness, the risk of recurrence, death). This type of testing uses gene expression of affected tissue to predict the course of the disease.

The question addressed in this evidence review is: Does prognostic testing using the tests described below in individuals diagnosed with a disease improve the net health outcome?

The specific clinical context of each test is described briefly in the following sections. The following PICOTS were used to select literature to inform this review.

Patients

The relevant population of interest is patients diagnosed with a disease (eg, CD, thymomas, and thymic carcinomas).

Interventions

The interventions of interest are miscellaneous prognostic tests, specifically Crohn's Prognostic for CD and DecisionDx-Thymoma for thymomas and thymic carcinomas.

Comparators

The comparator of interest is standard care without prognostic testing.

Outcomes

The outcomes of interest are overall survival, disease-specific survival, test accuracy and validity, change in disease status, and morbid events.

Timing

The timing of follow-up ranges from months for aggressiveness of the disease to years for risk of recurrence or death.

Setting

These tests are offered commercially through various manufacturers.

Technically Reliable

Assessment of technical reliability focuses on specific tests and operators and requires review of unpublished and often proprietary information. Review of specific tests, operators, and unpublished data are outside the scope of this evidence review and alternative sources exist. This evidence review focuses on the clinical validity and clinical utility.

Prognostic Testing for Crohn's Disease with Crohn's Prognostic

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

No studies of the 6-marker Crohn's Prognostic test were identified.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

Direct evidence for clinical utility is lacking.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

It is not possible to construct a chain of evidence for clinical utility due to the lack of clinical validity.

Section Summary: Crohn's Disease

Direct and indirect evidence for clinical utility of the Crohn's Prognostic test to identify individuals likely to have an aggressive disease course are currently lacking.

Prognostic Testing for Thymomas and Thymic Carcinomas with DecisionDx-Thymoma*Clinically Valid*

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

No full-length, peer-reviewed studies assessing DecisionDx-Thymoma were identified.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence: Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

Direct evidence for clinical utility is lacking.

Chain of Evidence: Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

It is not possible to construct a chain of evidence for clinical utility due to the lack of clinical validity.

Section Summary: Thymomas and Thymic Carcinomas

Evidence for the clinical validity and utility of the DecisionDx-Thymoma test to identify individuals likely to have an aggressive disease course is currently lacking.

Therapeutic Testing

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function—including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice

Clinical Context and Test Purpose

The purpose of therapeutic testing in patients who have been diagnosed with conditions like colon cancer and non-Hodgkin lymphoma is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does therapeutic testing using ResponseDX: Colon and TransPredict Fc gamma 3A in individuals diagnosed with colon cancer or non-Hodgkin lymphoma improve the net health outcome?

The specific clinical context of each test is described briefly in the following sections. The following PICOTS were used to select literature to inform this review.

Patients

The relevant population of interest is patients diagnosed with colon cancer or non-Hodgkin lymphoma.

Interventions

The interventions of interest are miscellaneous tests for variants that affect response to treatment or environmental exposure, specifically ResponseDX: Colon and TransPredict Fc gamma 3A.

Comparators

The comparator of interest is standard care without therapeutic testing.

Outcomes

The outcomes of interest are overall survival, disease-specific survival, test accuracy and validity, change in disease status, and morbid events.

Timing

The timing of follow-up ranges from weeks for treatment selection to years for survival outcomes.

Setting

These tests are offered commercially through various manufacturers.

Therapeutic Testing for Colon Cancer with ResponseDX: Colon

No full-length, peer-reviewed studies of the ResponseDX: Colon test were identified.

Section Summary: Colon Cancer

Evidence supporting the use of the ResponseDX Colon test to guide treatment selection in patients with colon cancer is currently lacking.

Therapeutic Testing for Non-Hodgkin Lymphoma and Rheumatoid Arthritis With TransPredict Fc Gamma 3A

Systematic Reviews

Two meta-analyses were identified, which came to different conclusions about the association between *FCGR2A* and *FCGR3A* single nucleotide variants and response to rituximab.

Ghesquières et al (2017) published a patient-level meta-analysis from 2 cohorts of patients with B-cell lymphoma (see Table 12).⁵¹ There was a marginally significant trend toward worse event-free survival for patients with *FCGR3A* (see Table 13). In a meta-analysis of patients with rheumatoid arthritis, Lee et al (2014) reported no significant association between *FCGR3A* genotype and response to rituximab or TNF blockers (see Table 13).⁵² However, stratification by biologic type indicated an association between the *FCGR3A* VV+VF genotype and nonresponders to rituximab. Statistical heterogeneity was high ($I^2=82\%$).

Table 12. Systematic Review Characteristics

Study	Trials	Dates	Participants	N	Design
Ghesquières et al (2017) ⁵¹	2		Patients with B-cell lymphoma	1034	MA of patient-level data from cohort studies
Lee et al (2014) ⁵²	7	Through Jan 2014	Patients with rheumatoid arthritis	500	

MA: meta-analysis

Table 13. Systematic Review Results

Study	Event-Free Survival	Nonresponse to Rituximab	Nonresponse to TNF Blocker
Ghesquières et al (2017) ⁵¹			
HR (95% CI)	0.87 (0.76 to 0.99)		
p	0.04		
Lee et al (2014) ⁵²			
OR (95% CI)		0.88 (0.51 to 1.54)	1.337 (0.87 to 2.06)
p		0.66	0.19

CI: confidence interval; HR: hazard ratio; OR: Odds ratio; TNF: tumor necrosis factor

Small studies in patients with non-Hodgkin lymphoma have suggested that the Val158Phe variant of the *FCGR3A* gene might predict response to rituximab therapy, although survival outcomes do not differ by genotype. In subsequent, larger studies in rituximab-treated patients with follicular lymphoma and chronic lymphocytic leukemia, this finding was not replicated. Studies in other types of non-Hodgkin lymphoma have also reported no association between *FCGR3A* genotype and outcomes. Meta-analysis of studies in rheumatoid arthritis did not find an association between *FCGR3* genotype and response to rituximab.

Section Summary: Non-Hodgkin Lymphoma

There is mixed evidence on the TransPredict Fc gamma 3A test. Some studies have reported an association with response to rituximab while others have not. Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials. No studies examining the clinical utility of TransPredict Fc gamma 3A were identified. Factors supporting a chain of evidence for predicting response to rituximab are lacking primarily because the evidence for clinical validity of the test is lacking.

Future Risk Disease Testing

Clinical Context and Test Purpose

The purpose of testing for future risk of disease in asymptomatic patients is that predictive and presymptomatic types of testing can be used to detect gene variants associated with disorders that appear after birth, usually later in life. These tests can be used in individuals with a family history of a genetic disorder, but who themselves have no features of the disorder at the time of testing. Predictive testing can identify variants that increase an individual's risk of developing disorders with a genetic basis (eg, certain types of cancer or cardiovascular disease). Presymptomatic testing can determine whether a person will develop a genetic disorder, before any signs or symptoms appear, by determining whether an individual has a genetic variant that may lead to the development of the disease.

The question addressed in this evidence review is: Does testing of asymptomatic individuals for future risk of disease using the tests described below in asymptomatic individuals improve the net health outcome?

The specific clinical context of each test is described briefly in the following sections. The following PICOTS were used to select literature to inform this review.

Patients

The relevant population of interest is patients with a family history of a genetic disorder that might develop later in life but who are currently without symptoms of the disorder.

Interventions

The interventions of interest are miscellaneous genetic or molecular risk assessment tests, specifically ImmunoGenomic Profile.

Comparators

The comparator of interest is standard care without genetic testing for future risk.

Outcomes

The outcomes of interest are overall survival, disease-specific survival, test accuracy and validity, change in disease status, and morbid events.

Timing

The timing of follow-up varies by test and is discussed in the following sections.

Setting

These tests are offered commercially through various manufacturers.

Future Risk Disease Testing for Immunologic Disorders with ImmunoGenomic Profile

Technically Reliable

Assessment of technical reliability focuses on specific tests and operators and requires review of unpublished and often proprietary information. Review of specific tests, operators, and unpublished data are outside the scope of this evidence review and alternative sources exist. This evidence review focuses on the clinical validity and clinical utility.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

No full-length, peer-reviewed studies of the ImmunoGenomic Profile were identified.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence: Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

Direct evidence for clinical utility is lacking.

Direct Evidence: Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test.

It is not possible to construct a chain of evidence for clinical utility due to the lack of clinical validity.

Section Summary: Immunologic Disorders

Evidence for the clinical validity and utility of the ImmunoGenomic Profile to predict the risk of developing arthritis, asthma, allergies, or other chronic inflammatory disorders is currently lacking.

SUMMARY OF EVIDENCE

Diagnostic Testing

For individuals with symptoms of various conditions thought to be hereditary or with a known genetic component who receive diagnostic testing with a miscellaneous genetic or molecular test (eg, DNA Methylation Pathway Profile, Know Error, Celiac PLUS, GI Effects [Stool], IBD sgi Diagnostic), the evidence includes case series, cross-sectional studies, diagnostic accuracy studies, and cohort studies. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, change in disease status, and morbid events. The lack of demonstrated clinical utility of these tests is based on the following factors: (1) there is no or extremely limited published data addressing the test; and/or (2) there is insufficient evidence demonstrating the clinical validity of the test. For each test addressed, a literature review was conducted. The literature review was not comprehensive, but sufficient to establish lack of clinical utility. A test will be removed from this evidence review and addressed separately if it is determined that enough evidence has accumulated to reevaluate its potential clinical utility. The evidence is insufficient to determine the effects of the technologies on health outcomes.

For individuals who are being screened for colorectal cancer who receive *SEPT9* methylated DNA testing (eg, ColoVantage, Epi proColon, ColonSentry), the evidence includes case-control, cross-sectional, and prospective diagnostic accuracy studies along with systematic reviews of those studies. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, change in disease status, and morbid events. The PRESEPT prospective study estimated the sensitivity and specificity of Epi proColon detection of invasive adenocarcinoma at 48% and 92%, respectively. Other studies were generally low to fair quality. Based on results from these studies, the clinical validity of *SEPT9* methylated DNA screening is limited by the low sensitivity of the test given that the sensitivity of the test is lower than imaging screening strategies. Optimal intervals for retesting are not known. The evidence is insufficient to determine the effects of the technologies on health outcomes.

Prognostic Testing

For individuals who are diagnosed with various conditions (eg, Crohn's disease, thymomas and thymic carcinomas, rheumatoid arthritis) who receive therapeutic testing with a miscellaneous genetic or molecular test (eg, Crohn's Prognostic, DecisionDx-Thymoma), there are no published studies. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, change in disease status, and morbid events. The evidence is insufficient to determine the effects of the technologies on health outcomes.

Therapeutic Testing

For individuals who are diagnosed with various conditions (eg, colon cancer, non-Hodgkin lymphoma) who receive therapeutic testing with a miscellaneous genetic or molecular test (eg, ResponseDX: Colon, TransPredict Fc gamma 3A), the evidence includes case series, cross-sectional studies, diagnostic accuracy studies, and cohort studies. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, change in disease status, and morbid events. The lack of demonstrated clinical utility of these tests is based on the following factors: (1) there is no or extremely limited published data addressing the test; and/or (2) there is insufficient evidence demonstrating the clinical validity of the test. For each test addressed, a literature review was conducted. The literature review was not comprehensive, but sufficient to establish lack of clinical utility. A test will be removed from this evidence review and addressed separately if it is determined that enough evidence has accumulated to

reevaluate its potential clinical utility. The evidence is insufficient to determine the effects of the technologies on health outcomes.

Testing for Future Risk of Disease

For individuals with a family history of various conditions thought to be hereditary or with a known genetic component who receive testing for future risk of disease with a miscellaneous genetic or molecular test (eg, ImmunoGenomic Profile), the evidence includes diagnostic accuracy studies. Relevant outcomes are overall survival, disease-specific survival, test accuracy and validity, change in disease status, and morbid events. The lack of demonstrated clinical utility of these tests is based on the following factors: (1) there is no or extremely limited published data addressing the test; and/or (2) there is insufficient evidence demonstrating the clinical validity of the test. For each test addressed, a literature review is conducted. The literature review was not comprehensive, but sufficient to establish lack of clinical utility. A test will be removed from this evidence review and addressed separately if it is determined that enough evidence has accumulated to reevaluate its potential clinical utility. The evidence is insufficient to determine the effects of the technologies on health outcomes.

PRACTICE GUIDELINES AND POSITION STATEMENTS

Diagnostic Tests

Multiple Conditions

No guidelines or statements were identified.

Celiac Disease

The American College of Gastroenterology (2013) published an evidence-based consensus algorithm for the diagnosis and management of celiac disease.³³ A recommendation for genetic testing using a multigene panel test (eg, Celiac PLUS) was not included.

Irritable Bowel Syndrome

American College of Gastroenterology practice guidelines on ulcerative colitis (2010)⁴⁰ and Crohn's disease (2009)⁴² did not contain recommendations for multimarker panels that include genetic tests to facilitate diagnosis or prognosis.

Colorectal Cancer

National Comprehensive Cancer Network

Current National Comprehensive Cancer Network (NCCN) guidelines on colorectal cancer (CRC) screening (v.1.2018) state that tests for methylated *SEPT9* DNA "may provide an alternative for individuals who refuse other screening modalities. However, the NCCN panel notes that its ability to detect CRC and advanced adenomas is inferior to other recommended screening modalities. The interval for repeated testing is unknown."⁵³

American Cancer Society

The American Cancer Society (2018) has recommended that "adults aged 45 y and older with an average risk of CRC undergo regular screening with either a high-sensitivity stool-based test or a structural (visual) examination, depending on patient preference and test availability. As a part of the screening process, all positive results on noncolonoscopy screening tests should be followed up with timely colonoscopy."⁵⁴ The stool-based tests listed as options are a fecal immunochemical test, fecal occult blood test, and multi-target stool DNA test. The College

noted that "...at this time, mSept9 is not included in this guideline as an option for routine CRC screening for average-risk adults."

American College of Physicians

Based on its review of U.S. guidelines, the American College of Physicians issued a guidance statement in 2012 on screening for CRC.⁵⁵ For average-risk adults ages 50 to 75 years, the College recommended using a stool-based test, flexible sigmoidoscopy, or optical colonoscopy for screening. For high-risk patients, it recommended using optical colonoscopy. No recommendation for genetic or molecular testing of average-risk individuals was included.

U.S. Multi-Society Task Force of Colorectal Cancer

The U.S. Multi-Society Task Force of Colorectal Cancer represents the American College of Gastroenterology, the American Gastroenterological Association, and the American Society for Gastrointestinal Endoscopy.⁵⁶ The Task Force's 2017 clinical guidelines stated that the advantage of *SEPT9* assays for CRC screening is convenience. The disadvantage is "markedly inferior performance characteristics compared with FIT [fecal immunochemical test]." The guidelines also stated that the best frequency for performing the test is unknown and that the task force recommended not using *SEPT9* assays for CRC screening.

Prognostic Tests

Crohn's Disease

No guidelines or statements were identified.

Thymomas and Thymic Carcinomas

NCCN guidelines for thymomas and thymic carcinomas (v.2.2018) do not address the use of gene expression profiling of tumors of the thymus.⁵⁷

Therapeutic Tests

Colon Cancer

NCCN guidelines for colon cancer (v.2.2018) state that it has "not been established if molecular markers are useful in treatment determination (predictive markers) and prognosis."⁵⁸

Non-Hodgkin Lymphoma

American College of Rheumatology (2016) recommendations for the use of disease-modifying antirheumatic drugs and biologic agents in the treatment of rheumatoid arthritis do not address *FCGR3* testing.⁵⁹

U.S. PREVENTIVE SERVICES TASK FORCE RECOMMENDATIONS

Unless otherwise indicated for the diagnostic, prognostic, therapeutic, and future risk testing, no U.S. Preventive Services Task Force recommendations for genetic or molecular tests have been identified.

The U.S. Preventive Services Task Force updated its recommendations for CRC screening in adults in 2016.^{44,60} It recommended screening for CRC starting at age 50 years and continuing until age 75 years. The 2016 recommendations differ from the 2008 recommendations in that current guidance does not emphasize specific screening approaches but highlights evidence that CRC screening substantially reduces deaths from the disease among adults ages 50 to 75 years and not enough adults in the United States are using effective preventive interventions. The

evidence review supporting the recommendations included a search for studies of blood tests for methylated *SEPT9* DNA but concluded that the test “currently has limited evidence evaluating its use.”

ONGOING AND UNPUBLISHED CLINICAL TRIALS

Some currently unpublished trials that might influence this review are listed in Table 14.

Table 14. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT03218423 ^a	Performance of Epi proColon in Repeated Testing in the Intended Use Population	4500	Jan 2022

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

CODING

The following codes for treatment and procedures applicable to this policy are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

CPT/HCPCS

81327	SEPT9 (Septin9) (eg, colorectal cancer) promoter methylation analysis
81382	HLA Class II typing, high resolution (ie, alleles or allele groups); one locus (eg, HLA-DRB1, -DRB3/4/5, -DQB1, -DQA1, -DPB1, or -DPA1), each
81479	Unlisted molecular pathology procedure
82397	Chemiluminescent assay
82784	Gammaglobulin (immunoglobulin); IgA, IgD, IgG, IgM, each
83520	Immunoassay for analyte other than infectious agent antibody or infectious agent antigen; quantitative, not otherwise specified
84999	Unlisted chemistry procedure
86021	Antibody identification; leukocyte antibodies
86140	C-reactive protein;
86255	Fluorescent noninfectious agent antibody; screen, each antibody
87045	Culture, bacterial; stool, aerobic, with isolation and preliminary examination (eg, KIA, LIA), Salmonella and Shigella species
87046	Culture, bacterial; stool, aerobic, additional pathogens, isolation and presumptive identification of isolates, each plate
87075	Culture, bacterial; any source, except blood, anaerobic with isolation and presumptive identification of isolates
87102	Culture, fungi (mold or yeast) isolation, with presumptive identification of isolates; other source (except blood)
87177	Ova and parasites, direct smears, concentration and identification
87209	Smear, primary source with interpretation; complex special stain (eg, trichrome, iron hemotoxylin) for ova and parasites

87328	Infectious agent antigen detection by immunoassay technique, (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative, multiple-step method; cryptosporidium
87329	Infectious agent antigen detection by immunoassay technique, (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative, multiple-step method; giardia
87336	Infectious agent antigen detection by immunoassay technique, (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative, multiple-step method; Entamoeba histolytica dispar group
87798	Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; amplified probe technique, each organism
88346	Immunofluorescence, per specimen; initial single antibody stain procedure
88350	Immunofluorescence, per specimen; each additional single antibody stain procedure (List separately in addition to code for primary procedure)

Diagnoses

Experimental / Investigational for all diagnoses related to this medical policy.

REVISIONS

01-07-2016	Policy added to the bcbsks.com web site on 12-08-2015 with an effective date of 01-07-2016.
08-17-2016	Updated Description section.
	Updated Rationale section.
	In Coding section: <ul style="list-style-type: none"> ▪ Removed CPT code: 88347.
	Updated References section.
01-01-2017	In Coding section: <ul style="list-style-type: none"> ▪ Added CPT code: 81327 (<i>New code, effective January 1, 2017</i>).
10-25-2017	Updated Description section.
	In Policy section: <ul style="list-style-type: none"> ▪ Added new Item A 1, "Testing of an affected (symptomatic) individual's germline to benefit the individual (excluding reproductive testing)". ▪ Previous Item A 1 is new Item A 2. ▪ Removed previous Item A 2, "Risk assessment". ▪ Removed previous Item A 4, "Genetic variants that alter response to treatment or to an environmental factor". ▪ Added new Item A 4, "Therapeutic testing". ▪ Added new Item A 5, "Testing an asymptomatic individual to determine future risk of disease".
	Updated Rationale section.
	In Coding section: <ul style="list-style-type: none"> ▪ Removed CPT code: 81401.
	Updated References section.
Updated Description section.	
12-20-2017	Updated Rationale section.
	Updated Description section.
10-12-2018	Updated Description section.

	Updated Rationale section.
	In Coding section: ▪ Added CPT codes: 88346, 88349.
	Updated References section.
	Removed Appendix section.
01-01-2019	In Coding section: ▪ Revised nomenclature to CPT code: 81327.

REFERENCES

1. Pfeifer JD, Zehnbauser B, Payton J. The changing spectrum of DNA-based specimen provenance testing in surgical pathology. *Am J Clin Pathol*. Jan 2011;135(1):132-138. PMID 21173135
2. Beauvais W, Fournie G, Jones BA, et al. Modelling the expected rate of laboratory biosafety breakdowns involving rinderpest virus in the post-eradication era. *Prev Vet Med*. Nov 1 2013;112(3-4):248-256. PMID 24029703
3. Pallav K, Kabbani T, Tariq S, et al. Clinical utility of celiac disease-associated HLA testing. *Dig Dis Sci*. Sep 2014;59(9):2199-2206. PMID 24705698
4. Ludvigsson JF, Leffler DA, Bai JC, et al. The Oslo definitions for coeliac disease and related terms. *Gut*. Jan 2013;62(1):43-52. PMID 22345659
5. Prometheus® Therapeutics and Diagnostics. IBD sgi Diagnostic <https://www.prometheuslabs.com/Products/Default.aspx>. Accessed June 25, 2018.
6. Pietzak MM, Schofield TC, McGinniss MJ, et al. Stratifying risk for celiac disease in a large at-risk United States population by using HLA alleles. *Clin Gastroenterol Hepatol*. Sep 2009;7(9):966-971. PMID 19500688
7. Ford AC, Quigley EM, Lacy BE, et al. Efficacy of prebiotics, probiotics, and synbiotics in irritable bowel syndrome and chronic idiopathic constipation: systematic review and meta-analysis. *Am J Gastroenterol*. Oct 2014;109(10):1547-1561; quiz 1546, 1562. PMID 25070051
8. National Institute for Health and Care Excellence (NICE). Irritable bowel syndrome in adults: diagnosis and management [CG61]. 2017; <https://www.nice.org.uk/guidance/cg61>. Accessed June 29, 2018.
9. McKenzie YA, Alder A, Anderson W, et al. British Dietetic Association evidence-based guidelines for the dietary management of irritable bowel syndrome in adults. *J Hum Nutr Diet*. Jun 2012;25(3):260-274. PMID 22489905
10. Weinberg DS, Smalley W, Heidelbaugh JJ, et al. American Gastroenterological Association Institute guideline on the pharmacological management of irritable bowel syndrome. *Gastroenterology*. Nov 2014;147(5):1146-1148. PMID 25224526
11. Hill C, Guarner F, Reid G, et al. Expert consensus document. The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol*. Aug 2014;11(8):506-514. PMID 24912386
12. Trinkley KE, Nahata MC. Treatment of irritable bowel syndrome. *J Clin Pharm Ther*. Jun 2011;36(3):275-282. PMID 21545610
13. Hungin AP, Mulligan C, Pot B, et al. Systematic review: probiotics in the management of lower gastrointestinal symptoms in clinical practice -- an evidence-based international guide. *Aliment Pharmacol Ther*. Oct 2013;38(8):864-886. PMID 23981066
14. Ortiz-Lucas M, Tobias A, Saz P, et al. Effect of probiotic species on irritable bowel syndrome symptoms: A bring up to date meta-analysis. *Rev Esp Enferm Dig*. Jan 2013;105(1):19-36. PMID 23548007
15. Whelan K. Probiotics and prebiotics in the management of irritable bowel syndrome: a review of recent clinical trials and systematic reviews. *Curr Opin Clin Nutr Metab Care*. Nov 2011;14(6):581-587. PMID 21892075

16. Stevenson C, Blaauw R, Fredericks E, et al. Randomized clinical trial: effect of *Lactobacillus plantarum* 299 v on symptoms of irritable bowel syndrome. *Nutrition*. Oct 2014;30(10):1151-1157. PMID 25194614
17. Shavakhi A, Minakari M, Farzamnia S, et al. The effects of multi-strain probiotic compound on symptoms and quality-of-life in patients with irritable bowel syndrome: A randomized placebo-controlled trial. *Adv Biomed Res*. 2014;3:140. PMID 25161987
18. Ludidi S, Jonkers DM, Koning CJ, et al. Randomized clinical trial on the effect of a multispecies probiotic on visceroperception in hypersensitive IBS patients. *Neurogastroenterol Motil*. May 2014;26(5):705-714. PMID 24588932
19. Rogha M, Esfahani MZ, Zargarzadeh AH. The efficacy of a synbiotic containing *Bacillus Coagulans* in treatment of irritable bowel syndrome: a randomized placebo-controlled trial. *Gastroenterol Hepatol Bed Bench*. Summer 2014;7(3):156-163. PMID 25120896
20. Urgesi R, Casale C, Pistelli R, et al. A randomized double-blind placebo-controlled clinical trial on efficacy and safety of association of simethicone and *Bacillus coagulans* (Colinox(R)) in patients with irritable bowel syndrome. *Eur Rev Med Pharmacol Sci*. 2014;18(9):1344-1353. PMID 24867512
21. Sisson G, Ayis S, Sherwood RA, et al. Randomised clinical trial: A liquid multi-strain probiotic vs. placebo in the irritable bowel syndrome--a 12 week double-blind study. *Aliment Pharmacol Ther*. Jul 2014;40(1):51-62. PMID 24815298
22. Genova Diagnostics. GI Effects® Comprehensive Profile - Stool. n.d.; <https://www.gdx.net/product/gi-effects-comprehensive-stool-test>. Accessed June 29, 2018.
23. Liew CC, Ma J, Tang HC, et al. The peripheral blood transcriptome dynamically reflects system wide biology: a potential diagnostic tool. *J Lab Clin Med*. Mar 2006;147(3):126-132. PMID 16503242
24. Yip KT, Das PK, Suria D, et al. A case-controlled validation study of a blood-based seven-gene biomarker panel for colorectal cancer in Malaysia. *J Exp Clin Cancer Res*. Sep 16 2010;29:128. PMID 20846378
25. Innovative Diagnostic Laboratory. ColonSentry®. n.d.; <https://www.myinnovativelab.com/colonsentry/>. Accessed June 29, 2018.
26. Chao S, Ying J, Liew G, et al. Blood RNA biomarker panel detects both left- and right-sided colorectal neoplasms: a case-control study. *J Exp Clin Cancer Res*. Jul 23 2013;32:44. PMID 23876008
27. Targan SR, Landers CJ, Yang H, et al. Antibodies to CBir1 flagellin define a unique response that is associated independently with complicated Crohn's disease. *Gastroenterology*. Jun 2005;128(7):2020-2028. PMID 15940634
28. Ippoliti A, Devlin S, Mei L, et al. Combination of innate and adaptive immune alterations increased the likelihood of fibrostenosis in Crohn's disease. *Inflamm Bowel Dis*. Aug 2010;16(8):1279-1285. PMID 20027650
29. Abreu MT, Taylor KD, Lin YC, et al. Mutations in NOD2 are associated with fibrostenosing disease in patients with Crohn's disease. *Gastroenterology*. Sep 2002;123(3):679-688. PMID 12198692
30. Genentech Inc. Rituxan / Rituximab. n.d.; <http://www.rituxan.com/>. Accessed June 29, 2018.
31. Genova Diagnostics. ImmunoGenomic® Profile. n.d.; <https://www.gdx.net/product/immunogenomic-profile-saliva>. Accessed June 29, 2018.
32. Pfeifer JD, Singleton MN, Gregory MH, et al. Development of a decision-analytic model for the application of STR-based provenance testing of transrectal prostate biopsy specimens. *Value Health*. Sep-Oct 2012;15(6):860-867. PMID 22999136
33. Rubio-Tapia A, Hill ID, Kelly CP, et al. ACG clinical guidelines: diagnosis and management of celiac disease. *Am J Gastroenterol*. May 2013;108(5):656-676; quiz 677. PMID 23609613
34. Hanaway P. Ask the experts. *Explore (NY)*. May 2006;2(3):284. PMID 16781657
35. Prometheus® Therapeutics & Diagnostics. IBD sgi monograph: the synergistic role of serology, genetics, and inflammation in the diagnosis of inflammatory bowel disease, 2011. <http://www.prometheuslabs.com/Products/Default.aspx?section=GIDiagnostics>. Accessed June 17, 2015.

36. Shirts B, von Roon AC, Tebo AE. The entire predictive value of the Prometheus IBD sigmoid diagnostic product may be due to the three least expensive and most available components. *Am J Gastroenterol*. Nov 2012;107(11):1760-1761. PMID 23160303
37. Conrad K, Roggenbuck D, Laass MW. Diagnosis and classification of ulcerative colitis. *Autoimmun Rev*. Apr-May 2014;13(4-5):463-466. PMID 24424198
38. Laass MW, Roggenbuck D, Conrad K. Diagnosis and classification of Crohn's disease. *Autoimmun Rev*. Apr-May 2014;13(4-5):467-471. PMID 24424189
39. Ordas I, Eckmann L, Talamini M, et al. Ulcerative colitis. *Lancet*. Nov 3 2012;380(9853):1606-1619. PMID 22914296
40. Kornbluth A, Sachar DB. Ulcerative colitis practice guidelines in adults: American College of Gastroenterology, Practice Parameters Committee. *Am J Gastroenterol*. Mar 2010;105(3):501-523; quiz 524. PMID 20068560
41. Baumgart DC, Sandborn WJ. Crohn's disease. *Lancet*. Nov 3 2012;380(9853):1590-1605. PMID 22914295
42. Lichtenstein GR, Hanauer SB, Sandborn WJ. Management of Crohn's disease in adults. *Am J Gastroenterol*. Feb 2009;104(2):465-483; quiz 464, 484. PMID 19174807
43. Reese GE, Constantinides VA, Simillis C, et al. Diagnostic precision of anti-Saccharomyces cerevisiae antibodies and perinuclear antineutrophil cytoplasmic antibodies in inflammatory bowel disease. *Am J Gastroenterol*. Oct 2006;101(10):2410-2422. PMID 16952282
44. U.S. Preventive Services Task Force. Colorectal cancer: screening. 2016; <https://www.uspreventiveservicestaskforce.org/Page/Document/UpdateSummaryFinal/colorectal-cancer-screening2>. Accessed June 25, 2018.
45. Nian J, Sun X, Ming S, et al. Diagnostic accuracy of methylated SEPT9 for blood-based colorectal cancer detection: a systematic review and meta-analysis. *Clin Transl Gastroenterol*. Jan 19 2017;8(1):e216. PMID 28102859
46. Song L, Jia J, Peng X, et al. The performance of the SEPT9 gene methylation assay and a comparison with other CRC screening tests: A meta-analysis. *Sci Rep*. Jun 8 2017;7(1):3032. PMID 28596563
47. Li B, Gan A, Chen X, et al. Diagnostic performance of DNA hypermethylation markers in peripheral blood for the detection of colorectal cancer: a meta-analysis and systematic review. *PLoS One*. May 2016;11(5):e0155095. PMID 27158984
48. Yan S, Liu Z, Yu S, et al. Diagnostic value of methylated septin9 for colorectal cancer screening: a meta-analysis. *Med Sci Monit*. Sep 25 2016;22:3409-3418. PMID 27665580
49. Church TR, Wandell M, Lofton-Day C, et al. Prospective evaluation of methylated SEPT9 in plasma for detection of asymptomatic colorectal cancer. *Gut*. Feb 2014;63(2):317-325. PMID 23408352
50. Marshall KW, Mohr S, Khattabi FE, et al. A blood-based biomarker panel for stratifying current risk for colorectal cancer. *Int J Cancer*. Mar 1 2010;126(5):1177-1186. PMID 19795455
51. Ghesquières H, Larrabee BR, Haioun C, et al. FCGR3A/2A polymorphisms and diffuse large B-cell lymphoma outcome treated with immunochemotherapy: a meta-analysis on 1134 patients from two prospective cohorts. *Hematol Oncol*. Dec 2017;35(4):447-455. PMID 27282998
52. Lee YH, Bae SC, Song GG. Functional FCGR3A 158 V/F and IL-6 -174 C/G polymorphisms predict response to biologic therapy in patients with rheumatoid arthritis: a meta-analysis. *Rheumatol Int*. Apr 12 2014;34(10):1409-1415. PMID 24728031
53. National Comprehensive Cancer Network (NCCN). NCCN Clinical practice guidelines in oncology: colorectal cancer screening. Version 1.2018. https://www.nccn.org/professionals/physician_gls/pdf/colorectal_screening.pdf. Accessed June 25, 2018.
54. Wolf AMD, Fontham ETH, Church TR, et al. Colorectal cancer screening for average-risk adults: 2018 guideline update from the American Cancer Society. *CA Cancer J Clin*. May 30 2018. PMID 29846947
55. Qaseem A, Denberg TD, Hopkins RH, Jr., et al. Screening for colorectal cancer: a guidance statement from the American College of Physicians. *Ann Intern Med*. Mar 06 2012;156(5):378-386. PMID 22393133

56. Rex DK, Boland CR, Dominitz JA, et al. Colorectal cancer screening: recommendations for physicians and patients from the U.S. Multi-Society Task Force on Colorectal Cancer. *Am J Gastroenterol*. Jul 2017;112(7):1016-1030. PMID 28555630
57. National Comprehensive Cancer Network (NCCN). NCCN Clinical practice guidelines in oncology: thymomas and thymic carcinoma. Version 2.2018. https://www.nccn.org/professionals/physician_gls/pdf/thymic.pdf. Accessed June 25, 2018.
58. National Comprehensive Cancer Network (NCCN). NCCN Clinical practice guidelines in oncology: colon cancer. Version 2.2018. https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf. Accessed June 25, 2018.
59. Singh JA, Saag KG, Bridges SL, Jr., et al. 2015 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Rheumatol*. Jan 2016;68(1):1-26. PMID 26545940
60. Bibbins-Domingo K, Grossman DC, Curry SJ, et al. Screening for colorectal cancer: US Preventive Services Task Force Recommendation Statement. *Jama*. Jun 21 2016;315(23):2564-2575. PMID 27304597

Other References

1. Blue Cross and Blue Shield of Kansas Pathology Liaison Committee, July 2016.
2. Blue Cross and Blue Shield of Kansas Surgery Liaison Committee, February 2016; May 2017.
3. Blue Cross and Blue Shield of Kansas Oncology Liaison Committee, August 2017.