

Medical Policy



Title: Balloon Sinuplasty for Treatment of Chronic Sinusitis

Professional

Original Effective Date: November 10, 2006
 Revision Date(s): February 16, 2011;
 August 1, 2011; September 21, 2011;
 September 24, 2012; November 11, 2016;
 July 1, 2017; January 1, 2018;
 March 14, 2018; March 27, 2019;
 January 1, 2020; April 19, 2021
 Current Effective Date: November 11, 2016

Institutional

Original Effective Date: January 14, 2009
 Revision Date(s): February 16, 2011,
 August 1, 2011; September 21, 2011;
 September 24, 2012; November 11, 2016;
 July 1, 2017; January 1, 2018;
 March 14, 2018; March 27, 2019;
 January 1, 2020; April 19, 2021
 Current Effective Date: November 11, 2016

State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).

The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.

The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.

If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.

Populations	Interventions	Comparators	Outcomes
Individuals: • With chronic rhinosinusitis	Interventions of interest are: • Balloon ostial dilation as a stand-alone procedure	Comparators of interest are: • Medical management • Functional endoscopic sinus surgery	Relevant outcomes include: • Symptoms • Change in disease status • Quality of life • Treatment-related morbidity
Individuals: • With recurrent acute rhinosinusitis	Interventions of interest are: • Balloon ostial dilation a stand-alone procedure.	Comparators of interest are: • Medical management • Functional endoscopic sinus surgery alone	Relevant outcomes include: • Symptoms • Change in disease status • Quality of life • Treatment-related morbidity

DESCRIPTION

Balloon ostial dilation (also known as balloon sinuplasty) is proposed as an alternative to functional endoscopic sinus surgery (FESS) for patients with chronic rhinosinusitis (CRS) or recurrent acute rhinosinusitis (RARS) who fail medical management. The procedure involves placing a balloon in

the sinus ostium and inflating the balloon to stretch the opening. It can be performed as a stand-alone procedure or as an adjunctive procedure to functional endoscopic sinus surgery (FESS). This evidence review addresses BOD as a standalone procedure.

OBJECTIVE

The objective of this evidence review is to evaluate whether balloon ostial dilation improves the net health outcome for patients with CRS or RARS.

BACKGROUND

Chronic and Recurrent Acute Rhinosinusitis

Chronic rhinosinusitis (CRS) is characterized by purulent nasal discharge, usually without fever, that persists for weeks to months. Symptoms of congestion often accompany the nasal discharge. There also may be mild pain and/or a headache. Thickening of mucosa may restrict or close natural openings between sinus cavities and the nasal fossae, although symptoms vary considerably because of the location and shape of these sinus ostia.

Recurrent acute rhinosinusitis (RARS) is defined as 4 or more episodes per year of acute bacterial rhinosinusitis without signs or symptoms of rhinosinusitis between episodes.

Medical Treatment

Most cases of CRS and RARS are treated with medical therapy (e.g., antihistamines, steroids, nasal lavage, and antibiotics).¹

Functional Endoscopic Sinus Surgery

FESS involves the insertion of an endoscope into the nose for a direct visual examination of the openings into the sinuses. Using the endoscope and a combination of surgical tools (e.g., curettes, forceps, powered micro-debriders, powered shavers, and/or sinus balloon catheters), surgeons enlarge the patient's sinus openings to clear passageways in order to restore normal sinus ventilation and drainage. The goal of surgery is to improve sinus ventilation and drainage by enlarging the openings of the sinuses, removing any polyps and correcting significant structural problems that may be hindering drainage.

The maxillary sinus creates a unique challenge. The maxillary ostia, located within the ethmoid infundibulum, often cannot be accessed transnasally without excising a portion of the uncinat process. An alternative approach to the maxillary ostia is through the sinus, via the canine fossa. A guidewire can be advanced from within the maxillary sinus to the nasal fossa. The dilating balloon can enlarge the ostia while deflecting the uncinat process.

Approximately 350,000 FESS procedures are done each year in the United States for CRS.

Balloon Ostial Dilation

A newer procedure, balloon ostial dilatation can be used as an alternative or as an adjunct to FESS for those with CRS or RARS. The goal of this technique, when used as an alternative to FESS, is to improve sinus drainage using a less invasive approach. The procedure involves placing a guidewire in the sinus ostium, advancing a balloon over the guidewire, and then stretching the opening by inflating the balloon. The guidewire location is confirmed with fluoroscopy or with direct transillumination of the targeted sinus cavity. General anesthesia may be needed for this procedure to minimize patient movement. According to the manufacturer, the RELIEVA SPINPLUS® Balloon Sinuplasty System is intended to: provide a means to access the sinus space

and illuminate within and transilluminate across nasal and sinus structures; dilate the sinus ostia and spaces associated with the paranasal sinus cavities for diagnostic and therapeutic procedures; and irrigate from within a target sinus for therapeutic procedures and to facilitate diagnostic procedures.<https://www.jnjmedicaldevices.com/en-US/product/relieva-spinplus-balloon-sinuplasty-system>

This evidence review is limited to BOD when used as a standalone procedure. BOD may also be used in combination with FESS.^{2,3} When used as an adjunct to FESS, it is intended to facilitate and/or increase access to the sinuses. BOD may also be used on 1 sinus and FESS on another sinus in the same patient during the same operation.

REGULATORY STATUS

In 2008, the Relieva™ Sinus Balloon Catheter (Acclarent, Menlo Park, CA) was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. FDA determined that this device was substantially equivalent to existing devices for use in dilating the sinus ostia and paranasal spaces in adults and maxillary sinus spaces in children. Subsequent devices developed by Acclarent have also been cleared by FDA through the 510(k) process. They include the Relieva Spin Sinus Dilation System® (cleared in 2011) and the Relieva Seeker Balloon Sinuplasty System® (cleared in 2012).

In 2008, the FinESS™ Sinus Treatment (Entellus Medical, Maple Grove, MN) was cleared for marketing by FDA through the 510(k) process. The indication noted is to access and treat the maxillary ostia/ethmoid infundibulum in adults using a transantral approach (FDA product code: EOB). The bony sinus outflow tracts are remodeled by balloon displacement of adjacent bone and paranasal sinus structures. Two other balloon sinus ostial dilation devices, the ENTrigue® Sinus Dilation System (ENTrigue Surgical, acquired by more recently by Smith & Nephew), and the XprESS™ Multi-Sinus Dilation Tool, also received 510(k) clearance in 2012.

In 2013, a sinus dilation system (Medtronic Xomed, Jacksonville, FL), later named the NuVent™ EM Balloon Sinus Dilation System, was cleared for marketing by FDA through the 510(k) process for use in conjunction with a Medtronic computer-assisted surgery system when surgical navigation or image-guided surgery may be necessary to locate and move tissue, bone, or cartilaginous tissue surrounding the drainage pathways of the frontal, maxillary, or sphenoid sinuses.

Also, in 2013, a sinus dilation system (Smith & Nephew), later named the Ventera™ Sinus Dilation System, was cleared for marketing through the 510(k) process to access and treat the frontal recesses, sphenoid sinus ostia, and maxillary ostia/ethmoid infundibula in adults using a transnasal approach. Ventera™ Sinus Dilation System does not require a guide wire or an illumination system as it is intended for use as a tool in combination with endoscopic sinus surgery.²

Table 1 summarizes the currently FDA cleared balloon sinus dilation devices.

FDA product code: LRC.

Table 1. Balloon Ostial Dilation Devices Cleared by the U.S. Food and Drug Administration

Device	Manufacturer	510(k) No.	Date Cleared	Indication
MESIRE - Balloon Sinus Dilatation System	Meril Life Sciences	K172737	12/12/2017	Sinus Ostia Dilation
Relieva UltirraNav Sinus Balloon Catheter	Acclarent Inc.	K161698	10/24/2016	Sinus Ostia Dilation
Vent-Os Sinus Dilation Family	Sinusys Corp.	K160770	6/29/2016	Sinus Ostia Dilation
Relieva Scout Multi-Sinus Dilation System	Acclarent Inc.	K153341	2/12/2016	Sinus Ostia Dilation
XprESS Multi-Sinus Dilation System	Entellus Medical Inc.	K152434	11/20/2015	Sinus Ostia Dilation
DSS Sinusplasty Balloon Catheter	Intuit Medical Products LLC	K143738	8/27/2015	Sinus Ostia Dilation
Relieva SpinPlus Balloon Sinuplasty System	Acclarent Inc.	K143541	4/22/2015	Sinus Ostia Dilation
XprESS Multi-Sinus Dilation Tool	Entellus Medical Inc.	K142252	10/17/2014	Sinus Ostia Dilation
Relieva Scout Multi-Sinus Dilation System	Acclarent Inc.	K140160	2/20/2014	Sinus Ostia Dilation

POLICY

- A. Balloon sinus ostial dilation is considered **medically necessary** for treating rhinosinusitis lasting longer than 12 weeks or 4 or more documented episodes of acute rhinosinusitis (in one year) when **ALL** of the following are met:
1. Confirmation with computed tomography (CT) scan and/or nasal endoscopy of findings of rhinosinusitis that includes one or more of the following:
 - a) Nasal endoscopy showing purulent discharge or persistent inflammatory sinus pathology, **OR**
 - b) CT evidence of mucosal thickening, bony remodeling, bony thickening, or obstruction of the ostiomeatal complex; **AND**
 2. Balloon sinus ostial dilation is limited to the frontal, maxillary or sphenoid sinuses; **AND**
 3. Balloon sinus ostial dilation is performed either as a stand-alone procedure or as part of functional endoscopic sinus surgery (FESS); **AND**
 4. Balloon sinus ostial dilation is performed on patients whose symptoms persist despite maximal medical therapy (see Policy Guidelines).
- B. Balloon ostial dilation is considered **not medically necessary** in the following situations:
1. Nasal polyposis (Grade 2 or greater);
 2. Samter's triad (aspirin sensitivity);
 3. Severe sinusitis secondary to autoimmune or connective tissue disorders (i.e., including, but not limited to, sarcoidosis, Wegener's granulomatosis);
 4. Severe sinusitis secondary to ciliary dysfunction, including, but not limited to, cystic fibrosis;
 5. Contraindication to, or inability to tolerate local and/or topical anesthetic;
 6. History of failed balloon procedure in the sinus to be treated;
 7. Sinusitis with extensive fungal disease;
 8. Isolated ethmoid sinus disease;
 9. Significant neo-osteogenesis.
- C. Balloon sinus ostial dilation for treatment of chronic sinusitis is considered **experimental / investigational** for any other indication.

Policy Guidelines

1. There are specific category I CPT codes for these procedures (31295-31297). These codes may be used to describe balloon sinuplasty when no other surgical intervention has been performed on the same sinus site.
2. When a balloon sinuplasty is used as an adjunct in performance of a medically necessary functional endoscopic sinus surgery (FESS) in the treatment of chronic sinusitis, the use of the balloon catheter is considered included in the allowance for the FESS procedure. There is no additional reimbursement for use of the balloon catheter.
3. If balloon sinuplasty is performed in conjunction with cutting tools such as curettes and forceps, the procedure might be coded using the CPT codes for nasal/sinus endoscopy with maxillary antrostomy (31256), nasal/sinus endoscopy with frontal sinus exploration (31276),

or nasal/sinus endoscopy with sphenoidectomy (31287). In this instance, the balloon dilation would be considered inclusive/incidental to the procedure.

4. Maximal medical therapy:

- a) Oral antibiotics of 2-4 weeks' duration for patients with CRS (chronic rhinosinusitis) (culture-directed if possible).
- b) Oral antibiotics with multiple 1-3 week courses for patients with RARS (recurrent acute rhinosinusitis).
- c) Systemic or topical steroids (discretion of physician).
- d) Saline irrigation (optional).
- e) Topical and/or systemic decongestants (optional).

RATIONALE

This evidence review has been updated regularly with searches of the PubMed database. The most recent literature update was performed through November 17, 2020.

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function^{3/4}including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

BALLOON OSTIAL DILATION AS A STAND-ALONE PROCEDURE FOR PATIENTS WITH CHRONIC RHINOSINUSITIS

Clinical Context and Test Purpose

The purpose of balloon ostial dilation (BOD) as a stand-alone procedure in patients with chronic rhinosinusitis (CRS) is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as medical management and functional endoscopic sinus surgery (FESS).

The question addressed in this evidence review is: Does balloon ostial dilation as a stand-alone procedure improve the net health outcome for patients with chronic rhinosinusitis?

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals 18 years of age and older with CRS, defined as an inflammatory condition involving the paranasal sinuses and linings of the nasal passages characterized by purulent nasal discharge, nasal obstruction, facial pain or pressure, and reduction in sense of smell, usually without fever, that persists for 12 weeks or longer.

Interventions

The treatment being considered is BOD (also known as balloon sinuplasty). The procedure involves placing a balloon in the sinus ostium and inflating it to stretch the opening.

BOD can be performed in the operating room or in an office setting under local anesthesia.

Comparators

Comparators of interest include medical management (steroids, antibiotics, or decongestants) and FESS.

Outcomes

The general outcomes of interest are symptoms, change in disease status, quality of life, and treatment-related morbidity.

To quantify the severity of CRS and to assess treatment response, various outcomes measures can be used, including radiologic scores, endoscopic grading, and patient-reported quality of life measures. The primary outcome measures relevant for the treatment of chronic rhinosinusitis (CRS) are patient-reported symptoms and quality of life. Examiner evaluation of the nasal and sinus appearance and polyp size may provide some information about treatment outcomes, but these evaluations are limited by the lack of universally accepted standards.

Disease-specific patient-reported quality of life scores include the commonly used Sino-Nasal Outcome Test-20 (SNOT-20), which is a validated questionnaire for which patients complete 20 symptom questions on a categorical scale (0 [no bother] to 5 [worst symptoms can be]). Average rankings can be reported over all 20 symptoms, as well as by 4 subclassified symptom domains. The possible range of SNOT-20 scores is 0 to 5, with a higher score indicating a greater rhinosinusitis-related health burden. The impact of treatment is measured by calculating the difference between SNOT-20 scores before and after treatment. A SNOT-20 change score of 0.8 or greater is believed to be clinically meaningful. The SNOT-22, a variation of the SNOT-20, includes 2 additional questions (on "nasal obstruction" and "loss of smell and taste"). The minimally important difference in SNOT-22 is considered to be 8.9 points.⁴

The Lund-Mackay scoring system uses radiologist-rated information derived from computed tomography scans to assess opacification of the sinus cavities, generating a score from 0 to 24.⁵ Although CT scans can provide an objective measure, often they do not correlate well with symptoms.⁶

Six months to 1 year of follow-up is considered necessary to demonstrate efficacy.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;

- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded

Review of Evidence
Systematic Review

Levy et al (2016) conducted a systematic review and meta-analysis of BOD for CRS (Table 2).⁷ Studies of balloon ostial dilation in combination with FESS were included if they reported data on subgroups of patients undergoing BOD as a standalone procedure. Reviewers included 17 studies; 11 of these provided data for meta-analysis. Two RCTs were included. The other studies were prospective or retrospective observational studies.

Results of the meta-analyses conducted by Levy et al are summarized in Table 3. Change from baseline in quality of life, as measured by SNOT-20 scores was clinically and statistically significant in patients who received BOD. Secondary outcome measures of postoperative complications, debridements, and revision surgery were heterogeneously reported without the consistency or power needed to make statistically valid comparisons. The reviewers concluded that BOD for the treatment of CRS in the reported study population had positive impact on patient quality of life as assessed by a validated measurement. Improvements exceeded the threshold of 0.8 and could be considered clinically significant. The reviewers also concluded that additional information was needed to determine the role of BOD in specific patient populations such as those with moderate to advanced sinus disease, to compare the incidence of postoperative complications and debridements in patients who receive BOD compared with FESS, and additional study of patients outcomes following BOD in the operating room versus the office setting.

Table 2. Systematic Review of Balloon Ostial Dilation for Chronic Rhinosinusitis-Characteristics

Study	Search Dates	Studies	Participants	N (Range)	Design	Duration
Levy et al (2016) ⁷ .	1996-2014	17 (11 provided data for meta-analysis)	Adults >18 years undergoing transnasal paranasal sinus BOD for CRS	1032 (6-328)	<ul style="list-style-type: none"> • RCT (n=2) • Prospective cohort (n=9) • Retrospective cohort (n=6) 	Varied (<6 months to >1 year)

BOD: balloon ostial dilation; CRS: chronic rhinosinusitis; RCT: randomized controlled trial; N: sample size

Table 3. Systematic Review of Balloon Ostial Dilation for Chronic Rhinosinusitis-Results

Study	Quality of Life (SNOT-20)			CT Findings(Lund-McKay Score)	Recovery Time
Levy et al (2016) ⁷ .	Change from baseline <6 months	Change from baseline >1 year	BOD vs FESS	Improvement from baseline	<ul style="list-style-type: none"> • BOD vs FESS • Number days to return of regular activity following intervention

Study	Quality of Life (SNOT-20)			CT Findings(Lund-McKay Score)	Recovery Time
	N analyzed	242	214		
Pooled effect (95% CI)	1.45 (0.99, 1.91)	1.41 (1.07, 1.74)	-0.42 (-1.39, 1.55)	1.15 (0.87-1.43)	Weighted mean 1.72 days vs 4.84 days (P <.001)
\bar{P} (P-value)	78% (.001)	59% (.04)	76% (.04)	30% (.22)	NA

SNOT-20: Sino-Nasal Outcome Test-20; CT: computed tomography; BOD: balloon ostial dilation; FESS: functional endoscopic sinus surgery; N: sample size; CI: confidence interval

Randomized Controlled Trials

BOD as a standalone procedure for patients with CRS has been evaluated in 4 RCTs reported in 6 publications (Tables 2 and 3). Two studies were published after the systematic review conducted by Levy et al.^{8,9}

The largest RCT is the REMODEL (randomized evaluation of maxillary antrostomy versus ostial dilation efficacy through long-term follow-up) trial. REMODEL results at 6, 12, and 24 months have been reported in 3 publications.^{10,11,8} This was an industry-sponsored RCT that compared BOD as a stand-alone procedure with FESS. A total of 105 patients with CRS or RARS and failure of medical therapy were randomized to BOD or FESS. Patients with gross sinonasal polyposis were excluded. Balloon ostial dilation was performed with the Entellus device, which is labeled for a transantral approach. FESS consisted of maxillary antrostomy and uncinectomy with or without anterior ethmoidectomy. Thirteen patients withdrew consent before treatment, 11 (21%) in the FESS group and 2 (4%) in the BOD group. The primary outcomes were the change in SNOT-20 scores at 6-month follow-up and mean number of postoperative debridements. Secondary outcomes included recovery time, complication rates, and rates of revision surgery. Noninferiority analysis was performed for the primary outcome of change in symptom score and superiority analyses was performed on the debridement outcome.

Ninety-one patients who were enrolled in REMODEL were available at 6-month follow-up.¹⁰ The improvement in the mean SNOT-20 score was 1.67 (1.10) in the balloon dilation group and 1.60 (0.96) in the FESS arm ($P = .001$) for noninferiority. Postoperative debridements were more likely in the FESS group with a mean of 1.2 (1.0) compared to a mean of 0.1(0.6) in the balloon dilation group ($P < .001$) for superiority in the balloon arm). Patients in the BOD arm returned to normal daily activities faster (1.6 days vs 4.8 days, $P = .002$ for superiority) and required fewer days of prescription pain medications (0.9 days vs 2.8 days, $P = .002$ for superiority) with balloon dilation. There were no major complications in either group, and 1 patient in each group required revision surgery.

Bikhazi et al (2014) reported 1-year follow-up from the REMODEL trial.¹¹ Eighty-nine (96.7%) subjects were available at 1 year. Improvement in the mean SNOT-20 score was 1.64 in the balloon dilation arm and 1.65 in the FESS arm ($P < .001$ for noninferiority). During the year post procedure, both groups had fewer self-reported rhinosinusitis episodes (mean reduction in episodes, 4.2 in the balloon arm vs 3.5 in the FESS arm; $P < .001$).

Final REMODEL results were reported in Chandra et al (2016).⁸ This publication included results up to 2 years post procedure for subjects in the REMODEL trial, along with an additional 30 subjects treated with FESS or in-office balloon sinus dilation, for a reported total of 61 FESS patients and 74

BOD patients. Follow-up data were available for 130, 66, and 25 patients at 12, 18, and 24 months, respectively. Details about group-specific treatment received and loss to follow-up were not reported for the additional 30 patients not included in the REMODEL trial. The BOD group required 0.2 debridements per patient compared with 1.0 per patient in the FESS group ($P < .001$). Mean change in SNOT-20 score from baseline to 12-month follow-up was -1.59 ($P < .001$) and -1.60 ($P < .001$) for the BOD and FESS groups, respectively, which was considered clinically significant. These changes were maintained at 24 months. At 18 months, overall revision rates were 2.7% in the balloon dilation group and 6.9% in the FESS group.

In addition to REMODEL, 3 smaller RCTs provide evidence on the comparison of BOD to FESS in patients with CRS.

Minni et al (2018) published a prospective, randomized study comparing BOD and traditional endoscopic sinus surgery (ESS) for CRS of the frontal sinuses.⁹ At 3 Italian hospitals, 102 individuals (148 sinuses) were enrolled with mild involvement of the frontal sinus, the average post-procedure SNOT-20 scores for the BOD and ESS groups were 24.6 and 27.54 ($P = .42$), respectively; for patients with moderate/severe involvement, the scores were 23.47 and 30.71 ($P < .05$), respectively. Post-procedure Lund-Mackay scores were 0.58 (BOD) and 0.54 (ESS; $P = .30$) in the mild group and 0.53 (BOD) and 0.78 (ESS; $P = .38$) in the moderate/severe group.

Bizaki et al (2014) reported on results from a RCT that compared balloon ostial dilation with FESS among patients with symptomatic chronic or recurrent acute rhinosinusitis.⁴ Results were not reported separately for patients with CRS and RARS, and the study authors stated, "For this study, both CRS and RARS were considered to be 1 disease." The trial enrolled 46 subjects, 4 of whom withdrew; the analysis included 42 patients (n=21 in each group; statistical power calculations not reported). Both treatment groups demonstrated significant improvements in SNOT-22 scores from baseline to post procedure. There were no differences in change in total SNOT-22 scores between groups at 3 months post procedure.

Achar et al (2012) was an open-label pilot study of 24 patients with CRS who had failed medical therapy and were scheduled for surgery.¹² Patients were randomized to BOD or to FESS and followed for 24 weeks. The primary outcome measures were changes in SNOT-20 scores and clearance time using the saccharin test. Both groups improved significantly on both measures. The degree of improvement was greater for the balloon dilatation group than for the FESS group on both the SNOT-20 score (43.8 vs 29.7, $P < .03$) Patients who received BOD were able to return to normal activities sooner than those who received FESS (2.2 days vs 5.0 days; P NR). Adverse events were not reported.

Table 4. RCTs of BOD compared to FESS in CRS: Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
REMODEL ^{10,11,8} • NCT01525849 • (6 month data) • (12-month data)	US	10	2011-2014	135 adults with medically refractory chronic (68%) or recurrent acute (32%) rhinosinusitis according to AAO-HNS clinical practice guidelines; all	• BOD (office setting) • N=74	• FESS (operating room) • N=61

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
<ul style="list-style-type: none"> (24-month data) 				met criteria for medically necessary FESS. Patients with nasal polyps were excluded.		
Minni et al (2018) ⁹ ,	Italy	3	NR	102 adults (148 sinuses) with non-polypoid CRS according to European Position Paper on Rhinosinusitis (EPOS) (2012) criteria	<ul style="list-style-type: none"> BOD N=69 sinuses 	<ul style="list-style-type: none"> FESS (DRAF I) N=79 sinuses
Bizaki 2014 ⁴ ,	Finland	1	NR	42 adults with CRS or RARS who fulfilled indications for surgical treatment. Patients with visible polyps in nasal direct endoscopy were excluded.	<ul style="list-style-type: none"> BOD N=21 	<ul style="list-style-type: none"> FESS N=21
Achar 2012 ¹² ,	UK	2	NR	24 adults with CRS diagnosed as per EPOS guidelines who failed medical treatment (topical steroids for 12 weeks with or without antibiotics) and were proceeding to surgery. Patients with extensive nasal polyps were excluded.	<ul style="list-style-type: none"> BOD N=12 	<ul style="list-style-type: none"> FESS N=12

REMODEL: randomized evaluation of maxillary antrostomy versus ostial dilation efficacy through long-term follow-up; RCT: randomized controlled trial; BOD: balloon ostial dilation; FESS: functional endoscopic sinus surgery; CRS: chronic rhinosinusitis; NCT: National Clinical Trial; AAO-HNS: American Academy of Otolaryngology – Head and Neck Surgery; N: sample size; RARS: recurrent acute rhinosinusitis

Table 5. RCTs of BOD Compared to FESS in CRS: Results

Study	Quality of Life	Symptoms	CT Scan Results	Adverse Events
Outcome measure Number analyzed	Mean change from baseline in SNOT-20 score N=91 at 6 months, 89 at 12 months	Time to return to normal daily activities	Overall Ostial Patency N=89 patients, 169 ostia	
REMODEL ^{10,11,8} • NCT01525849 • (6 month data) • (12-month data) • (24-month data)				
BOD	6 months: 1.67 (1.10) 12 months: 1.64 (1.06) 24 months: -1.65	1.6 days	6 months: NR 12 months: 96.7% (88/91)	No complications 28.0% nasal bleeding 1 (2.1%) revision surgery through 1 year
FESS	6 months: 1.60 (0.96) 12 months: 1.65 (0.94) 24 months: -1.45	4.8 days	6 months: NR 12 months: 98.7% (77/78)	No complications 54.8% nasal bleeding 1 (2.4%) revision surgery through 1 year
Between-group p-value	6 months: $P < 0.001$ 12 months): 0.01 (95% CI -0.43 to 0.44); BOD noninferior to FESS ($P < .0001$) 24 months:	0.002	12 months: $P = NS$	Nasal bleeding: $P = .011$
Minni et al 2018 ⁹ ,				
Outcome measure Number analyzed	Mean decrease in SNOT-20 at 12 months mild: 105 sinuses severe: 33 sinuses		Mean decrease in Lund-McKay score at 12 months mild: 105 sinuses severe: 33 sinuses	102 patients
BOD	mild: 36.34 severe: 41.32		mild: 1.1 severe: 2.57	No major complications
FESS	mild: 38.0 severe: 36.57		mild: 1.03 severe: 2.29	No major complications

Study	Quality of Life	Symptoms	CT Scan Results	Adverse Events
Between group difference p-value	mild: $P = .42$ severe: $P < .05$		mild: $P = .30$ severe: $P = .38$	
Bizaki et al (2014) ⁴ ,				
Outcome measure Number analyzed	Mean decrease in SNOT-22 from baseline to 3 months N=42		NR	N=42
BOD	21.47			No major complications 7 infection, 2 crusting, 2 synechia, 1 anosmia, 1 bleeding
FESS	20.95			No major complications 4 infection, 3 crusting, 6synechia, 4 anosmia
Between group difference p-value	$P = .587$			$P > .05$
Achar et al (2012) ¹² ,				
Outcome measure Number analyzed	Mean decrease in SNOT-22 from baseline to 6 months N=24	Mean time to get back to routine activities	NR	NR
BOD	43.83 (SD 15.17)	2.2 days		
FESS	29.66 (SD 12.33)	5.0 days		
Between group difference p-value	$P = .026$	NR		

REMODEL: randomized evaluation of maxillary antrostomy versus ostial dilation efficacy through long-term follow-up; RCT: randomized controlled trial; BOD: balloon ostial dilation; FESS: functional endoscopic sinus surgery; SNOT-20: Sino-Nasal Outcome Test-20; NR: not reported

Tables 6 and 7 summarize the limitations of the RCTs of BOD in patients with CRS. A major limitation of these trials was a lack of blinding, combined with the use of subjective outcome measures, and small sample sizes. However, objective measures (CT findings), additional evidence from observational studies, and consistency and magnitude of effects across studies make these limitations less concerning.

Table 6. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
REMODEL	3. Source and characteristics of patients added to the study for final results was unclear	1. Randomization of added patients occurred outside of key study			1. Differential loss post-randomization between study arms
Minni et al 2018 ⁹ ,					
Achar et al ¹² ,					
Bikazi et al 2014 ⁴ ,	3. Combined patients with CRS and RARS; results not reported separately by diagnosis				1,2. 3 month followup may be insufficient to assess benefits and harms

REMODEL: randomized evaluation of maxillary antrostomy versus ostial dilation efficacy through long-term follow-up. The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 7. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
REMODEL		1, 2. Not blinded				
Minni et al ⁹ ,	3. Method not described	1,2, 3. No information on blinding	1. Not registered		1. Power calculation not reported	Results reported by sinuses (N=148), not by patient (N=102)
Achar et al ¹² ,		1, 2. Not blinded	1. Not registered		1. Power calculation not reported; small sample size (N=24)	

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Bizaki et al 2014 ⁴ .	3. Method not described	1,2, 3. No information on blinding	1. Not registered		1. Power calculation not reported; small sample size (N=42)	

REMODEL: randomized evaluation of maxillary antrostomy versus ostial dilation efficacy through long-term follow-up. The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Observational Study of Adverse Events

A retrospective cohort study used data from a large commercial insurance database to examine adverse events reported in patients who underwent balloon dilation (n=2851), FESS (n=11,955), or a hybrid procedure (n=1234) between 2011 and 2014.¹³ The primary outcomes were surgical complication and revision rates within 6 months of the initial surgery. The overall complication rate was 7.35% with FESS and 5.26% with balloon dilation. The 6-month revision rates for balloon dilation, FESS, and hybrid surgeries were 7.89%, 16.85%, and 15.15%, respectively. Almost all revisions occurred with FESS regardless of primary procedure. However, differences in revision rates could have been due to differences in disease severity in patients who received FESS versus balloon dilation. Major complications included orbital complications, cerebrospinal fluid leak, severe epistaxis, and requirement for revision.

Section Summary: Balloon Ostial Dilation as a Stand-Alone Procedure for Patients with Chronic Rhinosinusitis

Four RCTs have compared BOD to FESS for patients with CRS. The best evidence is from the REMODEL trial, which showed statistically and clinically significant improvements in quality of life for up to 24 months, as measured by the validated SNOT-20 scale. REMODEL results are supported by smaller RCTs, multiple comparative observational studies, and a systematic review showing improvements in quality of life, CT outcomes, and shorter recovery time with BOD than FESS. In a retrospective cohort study that used data from a large commercial insurance database to examine adverse events in patients who underwent BOD (n=2851 or FESS (n=11,955), the overall complication rate 5.26% with BOD and 7.35% with FESS.

BALLOON OSTIAL DILATION AS A STAND-ALONE PROCEDURE FOR PATIENTS WITH RECURRENT ACUTE RHINOSINUSITIS

Clinical Context and Test Purpose

The purpose of balloon ostial dilation (BOD) as a stand-alone procedure in patients with recurrent acute rhinosinusitis (RARS) is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as medical management and functional endoscopic sinus surgery,

The question addressed in this evidence review is: Does BOD improve the net health outcome for patients with RARS?

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals 18 years of age and older with RARS. The American Academy of Otolaryngology-Head and Neck Surgery defines RARS as 4 or more episodes per year of acute bacterial rhinosinusitis without signs or symptoms of rhinosinusitis between episodes.¹ Each episode of acute bacterial rhinosinusitis should meet the following diagnostic criteria:

- Acute rhinosinusitis that is caused by, or is presumed to be caused by, bacterial infection. A clinician should diagnose ABRS when: symptoms or signs of acute rhinosinusitis fail to improve within 10 days or more beyond the onset of upper respiratory symptoms, or symptoms or signs of acute rhinosinusitis worsen within 10 days after an initial improvement (double worsening)
- Confirming a true bacterial episode of rhinosinusitis is desirable, but not essential, for substantiating an underlying diagnosis of RARS

Interventions

The therapy being considered is balloon ostial dilation as a stand-alone procedure. The procedure involves placing a balloon in the sinus ostium and inflating it to stretch the opening.

Balloon ostial dilation can be performed in the operating room or in an office setting under local anesthesia.

Comparators

Comparators of interest include medical management and functional endoscopic sinus surgery.

Outcomes

The general outcomes of interest are symptoms, change in disease status, quality of life, and treatment-related morbidity.

To quantify the severity of RARS and to assess treatment response, various outcomes measures can be used, including radiologic scores, endoscopic grading, and patient-reported quality of life measures. The primary outcome measures relevant for the treatment of RARS are patient-reported symptoms and quality of life. Examiner evaluation of the nasal and sinus appearance and polyp size may provide some information about treatment outcomes, but these evaluations are limited by the lack of universally accepted standards.

Disease-specific patient-reported quality of life scores include the commonly used Sino-Nasal Outcome Test-20 (SNOT-20), which is a validated questionnaire for which patients complete 20 symptom questions on a categorical scale (0 [no bother] to 5 [worst symptoms can be]). Average rankings can be reported over all 20 symptoms, as well as by 4 subclassified symptom domains. The possible range of SNOT-20 scores is 0 to 5, with a higher score indicating a greater rhinosinusitis-related health burden. The impact of treatment is measured by calculating the difference between SNOT-20 scores before and after treatment. A SNOT-20 change score of 0.8 or greater is believed to be clinically meaningful.

The Chronic Sinusitis Survey (CSS) is a measure of symptoms and medication usage over an 8-week recall period.¹⁴ The CSS includes 3 questions regarding symptoms and 3 regarding medication usage, yielding a total score as well as symptom and medication subscores evaluated as secondary endpoints. CSS total score ranges from 0 to 100 in which a low CSS score represents greater symptoms and/or medication usage. The minimally clinically significant difference on the CSS has not been established.

A decrease in the number of acute infections occurring over a specified time period is used as an outcome measure in some studies.

Six months to 1 year of follow-up is considered necessary to demonstrate efficacy.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.

Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Randomized Controlled Trials

Two RCTs of BOD reported results separately for patients with RARS (Table 8). A third RCT, reported by Bizaki et al (2014) compared BOD with FESS among patients with CRS or RARS, but results were not reported separately by diagnosis.¹⁵ The study authors stated, "For this study, both CRS and RARS were considered to be 1 disease." This trial is discussed in the previous section on BOD for CRS.

In the REMODEL trial, 32% (N=29) of the patients enrolled had a diagnosis of RARS. The CABERNET (Comparison of Balloon Sinuplasty In-Office Versus Medical Management for Recurrent Acute Sinusitis Patients) trial compared BOD plus medical therapy to medical therapy alone in 59 patients with RARS. Both trials used the AAO-HNS diagnosis of RARS to select eligible patients: 4 or more episodes of acute rhinosinusitis in the past 12 months. In CABERNET, evidence of sinus or osteomeatal complex disease during an acute episode from a CT scan was also required for enrollment. In REMODEL, all patients met criteria for medically necessary FESS, but explicit CT requirements for patients with RARS were not specified.

Results of the RCTs of patients with RARS are summarized in Table 9. Among the 29 patients diagnosed with RARS in the REMODEL trial, there was a significant improvement in quality of life for those who received either BOD or FESS, and the difference between treatment arms was not significant ($P = .838$). Twelve-month results from REMODEL were reported in Bikhazi et al (2014).¹¹ Data were not reported separately by diagnosis, but the publication states, "At 1 year, symptom improvement in each of the 4 subgroups [including based on diagnosis] remained statistically significant ($P < .001$) in both treatment arms and there was no difference ($P = NS$) in improvement between patients who underwent balloon dilation or FESS." REMODEL results were not reported separately by diagnosis for secondary outcomes, or for the primary outcome (SNOT-20) at 24 months.

In Sikand et al (2019), the primary outcome was the difference between arms in change in Chronic Sinusitis Survey (CSS) score from baseline to 24 weeks. The change in CSS was significantly greater in the BOD group compared to the control group (mean change 37.3 vs 21.8; $P = .0424$). The study authors did not specify whether this was considered clinically significant. Patients in the BOD group had a lower mean number of sinus infections through the 24-week followup period (0.2 vs 0.9; $P = .0015$). Durability of the outcome measure differences was demonstrated up to 48 weeks. After the 24-week followup period, 18 of 30 patients who were randomized to the control arm elected to receive BOD. Of those who crossed over at 24 weeks, 0 reported no change or worsening of symptoms, 3 reported improved symptoms but still used nasal sprays at high rates, 4 had improved symptoms to varying degrees but were not eliminated, and 1 reported a sinus infection just before their 24-week visit. There was 1 procedure-related serious adverse event in the BOD group (the patient sought treatment for a headache in the emergency department the evening after the procedure), 2 possibly procedure-related nonserious adverse events, and no device-related adverse events.

Table 8. Summary of Key RCT Characteristics- Balloon Ostial Dilation for Recurrent Acute Rhinosinusitis

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
REMODEL ^{10,11,8} <ul style="list-style-type: none"> • NCT01525849 • (6 month data) • (12-month data) • (24-month data) 	US	10	2011-2014	Adults with medically refractory chronic (68%) or recurrent acute (32%) rhinosinusitis according to AAO-HNS clinical practice guidelines; all met criteria for medically necessary FESS	<ul style="list-style-type: none"> • BOD (office setting) • N=16 	<ul style="list-style-type: none"> • FESS (operating room) • N=13
Sikand et al (2019) ¹⁶ <ul style="list-style-type: none"> • CABERNET • NCT01714687 	US	3	2013-2015	Adults with a diagnosis of recurrent acute rhinosinusitis, defined as having 4 or more episodes of acute bacterial rhinosinusitis within the previous 12 months, characterized	<ul style="list-style-type: none"> • BOD plus medical management • N=30 	<ul style="list-style-type: none"> • Sham procedure plus medical management • N=29

Study; Trial	Countries	Sites	Dates	Participants	Interventions
				by signs or symptoms of acute rhinosinusitis 10 or more days beyond the onset of upper respiratory symptoms, or within 10 days after initial improvement (double worsening)	

RCT: randomized controlled trial.

Table 9. Summary of Key RCT Results- Balloon Ostial Dilation for Recurrent Acute Rhinosinusitis

Study	Quality of Life	Acute Exacerbations	Adverse Events
REMODEL • NCT01525849			
Outcome measure • Number analyzed	<ul style="list-style-type: none"> • Mean change from baseline in SNOT-20 score • N=29 	Mean number per year, year before to year after treatment	NR separately for patients with RARS
BOD	<ul style="list-style-type: none"> • 6 months: (RARS subgroup): -1.57 (+1.08); $P < .0001$ • 12 months: Data not reported separately for patients with RARS. "At 1 year, symptom improvement in each of the 4 subgroups [including based on diagnosis] remained statistically significant ($P < .001$) in both treatment arms and there was no difference ($P = NS$) in improvement between patients who underwent balloon dilation or FESS." • 24 months: NR separately for patients with RARS 	<ul style="list-style-type: none"> • 5.1 to 0.9 • $P < 0.0001$ 	•
FESS	<ul style="list-style-type: none"> • 6 months (RARS subgroup): -1.64 (+0.90); $P < .0001$ • 24 months: NR separately for patients with RARS 	<ul style="list-style-type: none"> • 4.5 to 0.8 • $P < 0.0001$ 	•
Between-group p-value	<ul style="list-style-type: none"> • 6 months: .838 	<ul style="list-style-type: none"> • .258 	•
Sikand et al (2019) ¹⁶ , • CABERNET • NCT01714687	•		
Outcome measure	<ul style="list-style-type: none"> • Mean change in CSS Score at 24 weeks 	<ul style="list-style-type: none"> • Mean number of post-enrollment 	<ul style="list-style-type: none"> • N=59

Study	Quality of Life	Acute Exacerbations	Adverse Events
<ul style="list-style-type: none"> Number analyzed 	<ul style="list-style-type: none"> N=59 	sinus infections, 24 weeks <ul style="list-style-type: none"> N=59 	
BOD + medical management	<ul style="list-style-type: none"> Total score: 37.3 (SD 24.4) Symptom subscore: 48.7 (SD 28.7) Medication subscore: 26.0 (SD 26.6) 	<ul style="list-style-type: none"> 0.2 (0.4) 	<ul style="list-style-type: none"> 1 serious procedure-related adverse event (headache leading to hospital admission) No device-related adverse events Nonserious AEs: 58.6%
Sham + medical management	<ul style="list-style-type: none"> Total score: 21.8 (29.0) Symptom subscore: 27.2 (40.1) Medication subscore: 16.4 (24.0) 	<ul style="list-style-type: none"> 0.9 (0.9) 	<ul style="list-style-type: none"> Nonserious AEs: 60.0%
Between-group p-value	<ul style="list-style-type: none"> Total score:.0424 Symptom subscore:.0484 Medication subscore:.2607 	<ul style="list-style-type: none"> .0015 	<ul style="list-style-type: none"> Nonserious AEs: P = NS

CI: confidence interval; HR: hazard ratio; NNT: number needed to treat; OR: odds ratio; RCT: randomized controlled trial; RR: relative risk.

Tables 10 and 11 summarize the limitations of the RCTs of BOD in patients with RARS. Major limitations include no blinding of outcome assessors, a very small number of patients studied, and variation in the comparators and outcome measures used across the studies.

Table 10. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
REMODEL	3. Some outcomes not reported separately by diagnosis of RARS	1. Randomization of added patients occurred outside of key study			1. Differential loss post-randomization between study arms
Sikand et al (2019) <ul style="list-style-type: none"> CABERNET 	<ul style="list-style-type: none"> 		Medical regimen not standardized (customized by the treating investigator)	5. Clinically significant difference on primary outcome (CSS) not specified	

CABERNET: Comparison of Balloon Sinuplasty In-Office Versus Medical Management for Recurrent Acute Sinusitis Patients; REMODEL: randomized evaluation of maxillary antrostomy versus ostial dilation efficacy through long-term follow-up.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 11. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
REMODEL		1, 2. Not blinded			Not powered to detect differences by RARS subgroup	
Sikand et al (2019) CABERNET		Patients, but not outcome assessors, blinded				4. Confidence intervals not reported

CABERNET: Comparison of Balloon Sinuplasty In-Office Versus Medical Management for Recurrent Acute Sinusitis Patients; REMODEL: randomized evaluation of maxillary antrostomy versus ostial dilation efficacy through long-term follow-up.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Section Summary: Balloon Ostial Dilation as a Standalone Procedure for Patients with Recurrent Acute Rhinosinusitis

Two RCTs of BOD reported results separately for patients with RARS; 1 (REMODEL) compared BOD to FESS in a subgroup of 29 patients, and the other (CABERNET) compared BOD to medical care in 59 patients. In the REMODEL study BOD was non-inferior to FESS on measures of quality of life at 6 months and 12 months post-procedure; 24-month results were not reported separately for patients with RARS. One RCT comparing balloon ostial dilation plus medical care to medical care alone in patients with RARS found significantly improved quality of life and lower mean number of sinus infections after 24 months in the balloon dilation group. A third RCT included a mix of patients with chronic and RARS and found improved quality of life compared to FESS, but results were not reported separately by diagnosis.

Summary of Evidence

For individuals with CRS who receive BOD as a stand-alone procedure, the evidence includes RCTs, observational studies, and systematic reviews. Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. In the REMODEL RCT, balloon ostial dilation was non-inferior to FESS for patients with chronic rhinosinusitis. Durability of effect was demonstrated in uncontrolled studies that followed patients who received balloon dilation for up to 24 months. Evidence from RCTs is supported by multiple observational studies and a systematic review showing improved quality of life following BOD. In a retrospective cohort study that used data from a large commercial insurance database to examine adverse events reported in patients who underwent balloon dilation (n=2851), FESS (n=11,955), or a hybrid procedure (n=1234), the overall complication rate was 7.35% with FESS and 5.26% with balloon dilation. The evidence is sufficient to determine the effects that the technology results in an improvement in the net health outcome.

For individuals with RARS who receive BOD as a stand-alone procedure, the evidence includes RCTs. Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. In the REMODEL study of BOD compared to FESS, 32% of patients were diagnosed with recurrent acute rhinosinusitis (N=29). Balloon ostial dilation was non-inferior to FESS on measures of quality of life at 6 months and 12 months post-procedure. One RCT comparing balloon ostial dilation plus medical care to medical care alone in patients with RARS found significantly improved quality of life and lower mean number of sinus infections after 24 months in the balloon dilation group. A third RCT included a mix of patients with chronic and RARS and found improved quality of life compared to FESS, but results were not reported separately by diagnosis. The body of evidence is limited by the small number of patients studied, unblinded outcome assessment, lack of appropriate comparators, and heterogeneity in outcome measures used. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

American Academy of Otolaryngology – Head and Neck Surgery et al

In 2018, the American Academy of Otolaryngology - Head and Neck Surgery (AAO-HNS) published a clinical consensus statement on balloon dilation of the sinuses.¹⁷ Participating subgroups included the Triologic Society, the American Rhinologic Society, the American Academy of Otolaryngic Allergy, and the American Academy of Allergy, Asthma & Immunology. The expert panel used Delphi method surveys to assess consensus on proposed statements. Statements achieving a mean score of 7.00 or higher and having no more than 1 outlier (2 or more Likert points from the mean in either direction) met criteria for consensus. Strong consensus was defined as a mean Likert score of 8.00 or higher with no outliers. The following statements met consensus; statements reaching strong consensus are **highlighted**.

Patient Criteria:

- **Balloon dilation is not appropriate for patients who are without both sinonasal symptoms and positive findings on CT. (Strong consensus)**
- **Balloon dilation is not appropriate for the management of headache in patients who do not otherwise meet the criteria for chronic sinusitis or recurrent acute sinusitis. (Strong consensus)**

- **Balloon dilation is not appropriate for the management of sleep apnea in patients who do not otherwise meet the criteria for chronic sinusitis or recurrent acute sinusitis. (Strong consensus)**
- **CT scanning of the sinuses is a requirement before balloon dilation can be performed. (Strong consensus)**
- Balloon dilation is not appropriate for patients with sinonasal symptoms and a CT that does not show evidence of sinonasal disease.
- Balloon dilation can be appropriate as an adjunct procedure to FESS in patients with chronic sinusitis without nasal polyps.
- There can be a role for balloon dilation in patients with persistent sinus disease who have had previous sinus surgery.
- There is a role for balloon sinus dilation in managing patients with recurrent acute sinusitis as defined in the AAO-HNSF guideline based on symptoms and CT evidence of ostial occlusion and mucosal thickening.

Perioperative Considerations:

- **Surgeons who consider reusing devices intended for dilation of the sinuses should understand the regulations set forth by the U.S. Food and Drug Administration for reprocessing such devices and ensure that they are followed. (Strong consensus)**
- Balloon dilation can be performed under any setting as long as proper precautions are taken and appropriate monitoring is performed.
- Balloon dilation can be performed under local anesthesia with or without sedation.

Outcome:

- Balloon dilation can improve short-term quality-of-life outcomes in patients with limited CRS without polyposis.
- Balloon dilation can be effective in frontal sinusitis.

The AAO-HNS updated its statement on balloon ostial dilation, reaffirming its 2010 position statement: "Sinus ostial dilation ... is a therapeutic option for selected patient with chronic rhinosinusitis.... This approach may be used alone... or in conjunction with other instruments...."¹⁸,

In 2015, the Academy's Foundation updated its 2007 clinical practice guidelines on adult sinusitis, which do not discuss surgical therapy or use of balloon sinuplasty.¹

National Institute for Health and Care Excellence

In 2008, a guidance on balloon catheter dilation of paranasal sinus ostia from the National Institute for Health and Care Excellence (NICE) stated:

- "Current evidence on the short-term efficacy of balloon catheter dilation of paranasal sinus ostia for chronic sinusitis is adequate and raises no major safety concerns.
- This procedure should only be carried out by surgeons with experience of complex sinus surgery, and specific training in both the procedure and the use of fluoroscopy.
- Publication of long-term outcomes will be helpful in guiding the future use of this technique. NICE may review the procedure upon publication of further evidence."¹⁹,

In 2016, NICE published a recommendation on the use of the XprESS Multi-Sinus Dilation System for the treatment of chronic rhinosinusitis²⁰:

1.1 "The case for adopting the XprESS multi-sinus dilation system for treating uncomplicated chronic sinusitis after medical treatment has failed is supported by the evidence. Treatment with XprESS leads to a rapid and sustained improvement in chronic symptoms, fewer acute episodes and improved quality of life which is comparable to functional endoscopic sinus surgery (FESS).

1.2 XprESS should be considered in patients with uncomplicated chronic sinusitis who do not have severe nasal polyposis. In these patients, XprESS works as well as FESS, is associated with faster recovery times, and can more often be done under local anesthesia."

The recommendation was based on the results of the REMODEL study: the committee "considered that the evidence from REMODEL demonstrated that balloon dilation (with either XprESS or FinESS) is clinically non-inferior to FESS in terms of alleviating symptoms in patients with uncomplicated chronic sinusitis." Single-arm observational studies were of lower quality but were consistent with the findings of the REMODEL study.

American Rhinologic Society

A position statement, revised in 2017, from the American Rhinologic Society, stated that sinus ostial dilation is "a therapeutic option for selected patients with chronic rhinosinusitis (CRS) ... who have failed appropriate medical therapy."²¹

U.S. Preventive Services Task Force Recommendations

Not applicable.

Ongoing and Unpublished Clinical Trials

A search of ClinicalTrials.gov in November 2020 did not identify any ongoing or unpublished trials that would likely influence this review.

CODING

The following codes for treatment and procedures applicable to this policy are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

CPT/HCPCS

31295	Nasal/sinus endoscopy, surgical, with dilation (e.g., balloon dilation); maxillary sinus ostium, transnasal or via canine fossa
31296	Nasal/sinus endoscopy, surgical, with dilation (e.g., balloon dilation); frontal sinus ostium
31297	Nasal/sinus endoscopy, surgical, with dilation (e.g., balloon dilation); sphenoid sinus ostium
31298	Nasal/sinus endoscopy, surgical, with dilation (e.g., balloon dilation); frontal and sphenoid sinus ostia
31299	Unlisted procedure, accessory sinuses
C1726	Catheter, balloon dilation, nonvascular

ICD-10 Diagnoses

J32.0	Chronic maxillary sinusitis
J32.1	Chronic frontal sinusitis
J32.2	Chronic ethmoidal sinusitis
J32.3	Chronic sphenoidal sinusitis
J32.4	Chronic pansinusitis
J32.8	Other chronic sinusitis

REVISIONS

02-16-2011	Description Section updated. Rationale Section updated. References Section updated. In Coding section: <ul style="list-style-type: none"> ▪ Added CPT codes: 31295, 31296, 31297.
08-01-2011	In the Policy section: Liberalized the medical policy language from "Use of a catheter-based inflatable device (balloon sinuplasty) in the treatment of sinusitis is considered experimental / investigational." to "Use of a catheter-based inflatable device (balloon sinuplasty) in the treatment of medically refractory chronic sinusitis may be considered medically necessary as a minimally invasive alternative to endoscopic sinus surgery." In the Rationale section: <ul style="list-style-type: none"> ▪ In the summary section, removed the second paragraph, "In addition, more information is needed to determine which patients and which sinuses might be treated with the balloon technique and which require standard approaches. Given the limitations of the available data, the uncertain impact on clinical outcomes and questions about which patients might be candidates for this procedure, this approach is considered investigational." Added Policy Guidelines section.
09-21-2011	In the Policy Guidelines section: <ul style="list-style-type: none"> ▪ Added the new Item #2, "Beginning in 2011, there are specific category I CPT codes for these procedures (31295-31297). These codes may be used to describe balloon sinuplasty when no other surgical intervention has been performed on the same sinus site." This statement was moved from the Coding section to the Policy Guidelines section. ▪ Removed from the new Item #3, "Plans should be aware of this possibility." In the Coding section: removed S2344 (deleted 04/01/11).
09-24-2012	Description section updated. Rationale section updated. Reference section updated.
11-11-2016	Updated Description section. In Policy section: <ul style="list-style-type: none"> ▪ Removed "Use of a catheter-based inflatable device (balloon sinuplasty) in the treatment of medically refractory chronic sinusitis may be considered medically necessary as a minimally invasive alternative to endoscopic sinus surgery." ▪ Replaced with the following: "A. Balloon sinus ostial dilation is considered medically necessary for treating rhinosinusitis lasting longer than 12 weeks or 4 or more documented episodes of acute rhinosinusitis (in one year) <u>when all of the following are met</u>: 1. Confirmation with computed tomography (CT) scan and/or nasal endoscopy of findings of rhinosinusitis that includes <u>one or more of the following</u>: a) Nasal endoscopy showing purulent discharge or persistent inflammatory sinus pathology, OR b) CT evidence of mucosal thickening, bony remodeling, bony thickening, or obstruction of the ostiomeatal complex; AND 2. Balloon sinus ostial dilation is limited to the frontal, maxillary or sphenoid sinuses; AND 3. Balloon sinus ostial dilation is performed either as a stand-alone procedure or as part of functional endoscopic sinus surgery (FESS); AND 4. Balloon sinus ostial dilation is performed on patients whose symptoms persist despite maximal medical therapy (see Policy Guidelines). B. Balloon ostial dilation is considered not medically necessary in the following situations: 1. Nasal polyposis (Grade 2 or greater). 2. Samter's triad (aspirin sensitivity). 3. Severe sinusitis secondary to autoimmune or connective tissue disorders (i.e., including, but not limited to, sarcoidosis, Wegener's granulomatosis). 4. Severe sinusitis secondary to ciliary dysfunction, including, but not limited to, cystic fibrosis. 5. Contraindication to, or inability to tolerate local and/or topical anesthetic. 6. History of failed balloon procedure

	<p>in the sinus to be treated. 7. Sinusitis with extensive fungal disease 8. Isolated ethmoid sinus disease. 9. Significant neo-osteogenesis. C. Balloon sinus ostial dilation for treatment of chronic sinusitis is considered experimental / investigational for all other indications."</p> <p>In Policy Guidelines:</p> <ul style="list-style-type: none"> ▪ In Item 1, removed, "Beginning in 2011, t" and added "T" to read "There are specific category I CPT codes for these procedures (31295-31297). These codes may be used to describe balloon sinuplasty when no other surgical intervention has been performed on the same sinus site." ▪ Added Item 4, "Maximal medical therapy: a) Oral antibiotics of 2-4 weeks' duration for patients with CRS (chronic rhinosinusitis) (culture-directed if possible. b) Oral antibiotics with multiple 1-3 week courses for patients with RARS (recurrent acute rhinosinusitis). c) Systemic or topical steroids (discretion of physician). d) Saline irrigation (optional). e) Topical and/or systemic decongestants (optional).
	Updated Rationale section.
	Updated References section.
07-01-2017	Updated Description section.
	Updated Rationale section.
	Updated References section.
01-01-2018	<p>In Coding section:</p> <ul style="list-style-type: none"> ▪ Added CPT code: 31298. ▪ Removed ICD-9 codes.
03-14-2018	Updated Description section.
	Updated Rationale section.
03-27-2019	Updated Description section.
	Updated Rationale section.
	Updated References section.
01-01-2020	<p>In Coding section:</p> <ul style="list-style-type: none"> ▪ Revised CPT codes: 31295, 31296, 31297, 31298
04-19-2021	Updated Description section.
	Updated Rationale section.
	Updated References section.

REFERENCES

1. Rosenfeld RM, Piccirillo JF, Chandrasekhar SS, et al. Clinical practice guideline (update): adult sinusitis. *Otolaryngol Head Neck Surg.* Apr 2015; 152(2 Suppl): S1-S39. PMID 25832968
2. Hathorn IF, Pace-Asciak P, Habib AR, et al. Randomized controlled trial: hybrid technique using balloon dilation of the frontal sinus drainage pathway. *Int Forum Allergy Rhinol.* Feb 2015; 5(2): 167-73. PMID 25360863
3. Plaza G, Eisenberg G, Montojo J, et al. Balloon dilation of the frontal recess: a randomized clinical trial. *Ann Otol Rhinol Laryngol.* Aug 2011; 120(8): 511-8. PMID 21922974
4. Bizaki AJ, Taulu R, Numminen J, et al. Quality of life after endoscopic sinus surgery or balloon sinuplasty: a randomized clinical study. *Rhinology.* Dec 2014; 52(4): 300-5. PMID 25479206
5. Hopkins C, Browne JP, Slack R, et al. The Lund-Mackay staging system for chronic rhinosinusitis: how is it used and what does it predict?. *Otolaryngol Head Neck Surg.* Oct 2007; 137(4): 555-61. PMID 17903570
6. Greguric T, Trkulja V, Baudoin T, et al. Association between computed tomography findings and clinical symptoms in chronic rhinosinusitis with and without nasal polyps. *Eur Arch Otorhinolaryngol.* May 2017; 274(5): 2165-2173. PMID 28154930
7. Levy JM, Marino MJ, McCoul ED. Paranasal Sinus Balloon Catheter Dilation for Treatment of Chronic Rhinosinusitis: A Systematic Review and Meta-analysis. *Otolaryngol Head Neck Surg.* Jan 2016; 154(1): 33-40. PMID 26519456

8. Chandra RK, Kern RC, Cutler JL, et al. REMODEL larger cohort with long-term outcomes and meta-analysis of standalone balloon dilation studies. *Laryngoscope*. Jan 2016; 126(1): 44-50. PMID 26228589
9. Minni A, Dragonetti A, Sciuto A, et al. Use of balloon catheter dilation vs. traditional endoscopic sinus surgery in management of light and severe chronic rhinosinusitis of the frontal sinus: a multicenter prospective randomized study. *Eur Rev Med Pharmacol Sci*. Jan 2018; 22(2): 285-293. PMID 29424885
10. Cutler J, Bikhazi N, Light J, et al. Standalone balloon dilation versus sinus surgery for chronic rhinosinusitis: a prospective, multicenter, randomized, controlled trial. *Am J Rhinol Allergy*. Sep-Oct 2013; 27(5): 416-22. PMID 23920419
11. Bikhazi N, Light J, Truitt T, et al. Standalone balloon dilation versus sinus surgery for chronic rhinosinusitis: a prospective, multicenter, randomized, controlled trial with 1-year follow-up. *Am J Rhinol Allergy*. Jul-Aug 2014; 28(4): 323-9. PMID 24823902
12. Achar P, Duvvi S, Kumar BN. Endoscopic dilatation sinus surgery (FEDS) versus functional endoscopic sinus surgery (FESS) for treatment of chronic rhinosinusitis: a pilot study. *Acta Otorhinolaryngol Ital*. Oct 2012; 32(5): 314-9. PMID 23326011
13. Chaaban MR, Rana N, Baillargeon J, et al. Outcomes and Complications of Balloon and Conventional Functional Endoscopic Sinus Surgery. *Am J Rhinol Allergy*. Sep 2018; 32(5): 388-396. PMID 29947260
14. Gliklich RE, Metson R. Techniques for outcomes research in chronic sinusitis. *Laryngoscope*. Oct 2015; 125(10): 2238-41. PMID 26390386
15. Bizaki AJ, Numminen J, Taulu R, et al. A Controlled, Randomized Clinical Study on the Impact of Treatment on Antral Mucociliary Clearance: Uncinectomy Versus Balloon Sinuplasty. *Ann Otol Rhinol Laryngol*. May 2016; 125(5): 408-14. PMID 26611244
16. Sikand A, Ehmer DR, Stolovitzky JP, et al. In-office balloon sinus dilation versus medical therapy for recurrent acute rhinosinusitis: a randomized, placebo-controlled study. *Int Forum Allergy Rhinol*. Feb 2019; 9(2): 140-148. PMID 30452127
17. Piccirillo JF, Payne SC, Rosenfeld RM, et al. Clinical Consensus Statement: Balloon Dilation of the Sinuses. *Otolaryngol Head Neck Surg*. Feb 2018; 158(2): 203-214. PMID 29389303
18. American Academy of Otolaryngology - Head and Neck Surgery. Position Statement: Dilation of sinuses, any method (e.g., balloon, etc.). 2017; <http://www.entnet.org/content/position-statement-dilation-sinuses-any-method-e.g.-balloon-etc>. Accessed January 21, 2020.
19. National Institute of Health and Care Excellence (NICE). Balloon catheter dilation of paranasal sinus ostia for chronic sinusitis [IPG273]. 2008; <http://www.nice.org.uk/guidance/IPG273/chapter/1-Guidance>. Accessed January 21, 2020.
20. National Institute of Health and Care Excellence (NICE). XprESS multi sinus dilation system for treating chronic sinusitis [MTG30]. 2016; <https://www.nice.org.uk/guidance/mtg30>. Accessed January 21, 2020.
21. American Rhinologic Society (ARS). Ostial Balloon Dilation Position Statement. 2017; https://www.american-rhinologic.org/index.php?option=com_content&view=article&id=33:ostial-balloon-dilation-position-statement&catid=26:position-statements&Itemid=197. Accessed January 21, 2020.

Other References

1. Blue Cross and Blue Shield of Kansas Otolaryngology Liaison Committee, August 2010; August 2011; August 2014; July 2015; January 2016; May 2017.