

Medical Policy



Title: Electrical Stimulation Devices for Home and Provider Use

Professional / Institutional
Original Effective Date: July 1, 2006 / February 11, 2009
Latest Revision Date: February 27, 2024
Current Effective Date: February 27, 2024

State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).

The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.

The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.

If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.

Populations	Interventions	Comparators	Outcomes
Transcutaneous Electrical Stimulation Devices (TENS) and Form Fitting Conductive Garment			
Individuals: • With chronic pain	Interventions of interest are: • Transcutaneous electrical nerve stimulation	Comparators of interest are: • Physical therapy • Pharmacotherapy	Relevant outcomes include: • Symptoms • Functional outcomes • Quality of life • Medication use
Individuals: • With acute pain	Interventions of interest are: • Transcutaneous electrical nerve stimulation	Comparators of interest are: • Pharmacotherapy	Relevant outcomes include: • Symptoms • Medication use
Individuals: • With chronic or episodic migraine	Interventions of interest are: • Transcutaneous electrical nerve	Comparators of interest are: • Pharmacotherapy	Relevant outcomes include: • Symptoms • Functional outcomes

Populations	Interventions	Comparators	Outcomes
	stimulation for acute migraine treatment		<ul style="list-style-type: none"> Quality of life Medication use
Individuals: <ul style="list-style-type: none"> With chronic or episodic migraine 	Interventions of interest are: <ul style="list-style-type: none"> Transcutaneous electrical nerve stimulation for migraine prevention 	Comparators of interest are: <ul style="list-style-type: none"> Pharmacotherapy 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Functional outcomes Quality of life Medication use
Electrostimulation and Electromagnetic Therapy			
Individuals: <ul style="list-style-type: none"> With any wound type (acute or nonhealing) 	Interventions of interest are: <ul style="list-style-type: none"> Electrostimulation Electromagnetic therapy 	Comparators of interest are: <ul style="list-style-type: none"> Standard wound care 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Change in disease status Morbid events Quality of Life Treatment-related morbidity
Interferential Current Stimulation			
Individuals: <ul style="list-style-type: none"> With musculoskeletal conditions 	Interventions of interest are: <ul style="list-style-type: none"> Interferential current stimulation 	Comparators of interest are: <ul style="list-style-type: none"> Physical therapy Medication Different type of electrical stimulation 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Functional outcomes Quality of life Medication use Treatment-related morbidity
Individuals: <ul style="list-style-type: none"> With gastrointestinal disorders 	Interventions of interest are: <ul style="list-style-type: none"> Interferential current stimulation 	Comparators of interest are: <ul style="list-style-type: none"> Dietary changes Medication Different type of electrical stimulation 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Functional outcomes Quality of life Medication use Treatment-related morbidity
Individuals: <ul style="list-style-type: none"> With poststroke spasticity 	Interventions of interest are: <ul style="list-style-type: none"> Interferential current stimulation 	Comparators of interest are: <ul style="list-style-type: none"> Standard stroke rehabilitation 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Functional outcomes Quality of life Treatment-related morbidity
Electrical Stimulation for the Treatment of Arthritis			
Individuals: <ul style="list-style-type: none"> With osteoarthritis 	Interventions of interest are: <ul style="list-style-type: none"> Electrical stimulation 	Comparators of interest are: <ul style="list-style-type: none"> Pharmacologic therapy Physical therapy 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Functional outcomes Health status measures Treatment-related morbidity

DESCRIPTION

Transcutaneous electrical nerve stimulation (TENS) describes the application of electrical stimulation to the surface of the skin. In addition to more traditional settings such as a physician's office or an outpatient clinic, TENS can be self-administered in a patient's home.

OBJECTIVE

The objective of this evidence review is to determine whether the application of transcutaneous electrical nerve stimulation improves the net health outcome in individuals with a variety of health conditions including chronic and/or acute pain, essential tremor, and attention deficit hyperactivity disorder.

BACKGROUND

Transcutaneous electrical nerve stimulation (TENS) has been used to treat chronic intractable pain, migraine headache pain, postsurgical pain, and pain associated with active or post trauma injury unresponsive to other standard pain therapies. It has been proposed that TENS may provide pain relief through the release of endorphins in addition to potential blockade of local pain pathways. TENS has also been used to treat dementia by altering neurotransmitter activity and increasing brain activity that is thought to reduce neural degeneration and stimulate regenerative processes.

Percutaneous electrical nerve stimulation is similar to TENS but uses microneedles that penetrate the skin instead of surface electrodes. Interferential stimulation uses a modulated waveform for deeper tissue stimulation, and the stimulation is believed to improve blood flow to the affected area.

REGULATORY STATUS

TENS devices consist of an electrical pulse generator, usually battery-operated, connected by wire to 2 or more electrodes, which are applied to the surface of the skin at the site of the pain. Since 1977, a large number of devices have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. Marketing clearance via the 510(k) process does not require data on clinical efficacy; as a result, these cleared devices are considered substantially equivalent to predicate devices marketed in interstate commerce before May 1976, the enactment date of the Medical Device Amendments. The cleared devices are also equivalent to devices that have been reclassified and do not require a premarket approval application. FDA product code: GZJ.

In 2014, the Cefaly® (STX-Med), which is a TENS device, was granted a de novo 510(k) classification by the FDA for the prophylactic treatment of migraine in patients 18 years of age or older.¹ The Cefaly® Acute and Cefaly® Dual devices were cleared by the FDA through the 510(k) process for the acute treatment of migraine in patients in 18 years of age or older and for both the acute treatment and prophylaxis of migraines in adults, respectively, in 2017.^{2,3} Other TENS devices cleared by the FDA through the 510(k) process for the prophylactic treatment of migraine in patients include Allive (Nu Eyne Co), Relivion (Leurolief Ltd.) and HeadTerm (EEspress) among others.^{4,5,6} FDA product code: PCC.

In 2018, the FDA reviewed the Cala ONE™ TENS device (Cala Health) via the de novo pathway and granted approval for the device as an aid in the transient relief of hand tremors following stimulation in the affected hand of adults with essential tremor. This prescription device is contraindicated for use in patients with an implanted electrical medical device, those that have suspected or diagnosed epilepsy or other seizure disorder, those who are pregnant, and patients with swollen, infected, inflamed areas, or skin eruptions, open wounds, or cancerous lesions. In October 2020, the FDA granted breakthrough device designation to the Cala Trio™ device for the treatment of action tremors in the hands of adults with Parkinson's disease.⁷ In November 2022, the Cala kIQ™ device was approved via the 510(k) pathway (K222237). The device is indicated to aid in the temporary relief of hand tremors in the treated hand following stimulation in adults with essential tremor. It was also approved to aid in the temporary relief of postural and kinetic hand tremor symptoms that impact some activities of daily living in the treated hand of adults with Parkinson's disease.

In 2019, the FDA permitted marketing of the first medical device to treat attention deficit hyperactivity disorder (ADHD) - the Monarch® external Trigeminal Nerve Stimulation (eTNS) System by NeuroSigma.⁸ The FDA reviewed the system through the de novo premarket review pathway. This prescription only TENS device is indicated for patients 7 to 12 years of age who are not currently taking prescription ADHD medication. The Monarch eTNS System is intended to be used in the home under the supervision of a caregiver. The device generates a low-level electrical pulse and connects via a wire to a small patch that adheres to a patient's forehead, just above the eyebrow.

In 2021, the FDA approved the Axon Therapy device (Neuralace Medical, Inc.) for marketing through the 510(k) process for relief of chronic, intractable postsurgical or posttraumatic pain in adults.⁹ The Axon Therapy device is an electromagnetic transcutaneous peripheral nerve stimulator. FDA product codes: QPL, IPF.

POLICY**A. Transcutaneous Electrical Nerve Stimulation Devices (TENS)**

1. May be considered **medically necessary** for the treatment of refractory chronic pain (e.g., chronic musculoskeletal pain or neuropathic pain)
2. Are **not medically necessary** for:
 - a. non-musculoskeletal pain, including but not limited to, visceral abdominal pain, and pelvic pain
 - b. acute post op musculoskeletal pain
3. Are **not medically necessary** using 2 lead, localized stimulation (E0720).
4. TENS is considered **experimental / investigational** for the prevention or treatment of migraine headache.
5. The use of TENS for any other condition is considered **experimental / investigational**

B. Form Fitting Conductive Garment

Is considered **medically necessary** when it meets the indications outlined in Transcutaneous Electrical Nerve Stimulation Devices (TENS) and the patient:

1. Is unable to manage without the garment due to large area or large number of sites or
2. Has skin conditions that preclude the application of conventional electrodes, adhesive tapes and lead wires or
3. Is applying electrical stimulation beneath a cast for disuse atrophy

C. Neuromuscular Electrical Stimulation Devices (NMES) are denied **experimental / investigational.****D. Interferential Therapy is denied **experimental / investigational**. (e.g., Sanexas, RS-4i, Sequential, Stimulator, Medstar&trade 100, EMSI Flex-IT)****E. Galvanic Stimulation Devices are denied **experimental / investigational**.****F. Microcurrent Stimulation Devices (MENS) are denied **experimental / investigational**.****G. H-wave Stimulation Devices are denied **experimental / investigational**.****H. Sympathetic Therapy for the treatment of pain is denied **experimental / investigational**.****I. Electrostimulation and Electromagnetic Therapy for the treatment of chronic wounds-is denied **experimental / investigational**.****J. Pulsed Electrical Stimulation for the treatment of osteoarthritis is denied **experimental / investigational**.**

- K. Electrical Stimulation for the treatment of hyperemesis gravidarum is denied **experimental / investigational**.
- L. Neurostimulation for the treatment of migraine pain and prevention of migraine headaches is denied **experimental / investigational**.

POLICY GUIDELINES

- A. Form Fitting Conductive Garment –Is a garment with conductive fibers which are separated from the patients' skin by layers of fabric used for delivering TENS and NMES.
- B. Neuromuscular Electrical Stimulation Devices (NMES) - Attempts to stimulate motor nerves to cause contraction of muscles rather than alter the perception of pain. NMES are used to prevent disuse atrophy, relax muscle spasm, increase blood circulation, improve range of motion and re-educate muscles.
- C. Interferential Therapy (such as the Medstar™ 100 and the RS-4i) - Interferential current stimulation (IFS) is a type of electrical stimulation. IFS has primarily been investigated as a technique to reduce pain but has also been proposed to increase function of patients with osteoarthritis and to treat other conditions such as dyspepsia, irritable bowel syndrome, and constipation.
- D. Galvanic Stimulation Devices - High voltage, pulsed stimulation used primarily for local edema reduction through muscle pumping and polarity effect. Edema is comprised of negatively charged plasma proteins. Placing electrodes over the edematous site disperses the negatively charged proteins.
- E. Microcurrent Stimulation Devices (MENS) - Uses a reduced electrical stimulation compared to TENS and acts on naturally occurring electrical impulses to decrease pain.
- F. H-wave Stimulation Devices - H-wave stimulation is a distinct form of electrical stimulation for medical purposes that involve repeated muscle contractions. H-wave electrical stimulation has been evaluated primarily as a pain treatment, but it has also been studied for other indications such as wound healing and improving postsurgical range of motion.
- G. Sympathetic Therapy - Sympathetic therapy describes a type of electrical stimulation of the peripheral nerves that is designed to stimulate the sympathetic nervous system in an effort to "normalize" the autonomic nervous system and alleviate chronic pain. Unlike TENS (transcutaneous electrical nerve stimulation) or interferential electrical stimulation, sympathetic therapy is not designed to treat local pain, but is designed to induce a systemic effect on sympathetically induced pain. (Dynatron STS)
- H. Electrostimulation and Electromagnetic Therapy - Electrostimulation (electrical stimulation) refers to the application of electrical current through electrodes placed directly on the skin. Electromagnetic therapy involves the application of electromagnetic fields, rather than direct electrical current. Both are proposed as treatments for wounds, generally chronic wounds.

- I. Pulsed Electrical Stimulation - Pulsed electrical and electromagnetic stimulation are being investigated to improve functional status and relieve pain related to osteoarthritis (OA) unresponsive to other standard therapies. Electrical stimulation is provided by an electronic device that noninvasively delivers a subsensory low-voltage, monophasic electrical field to the target site of pain. Pulsed electromagnetic fields are delivered via treatment coils placed over the skin. (BioniCare BIO-1000).
- J. Electrical Stimulation for hyperemesis gravidarum (e.g., Prima Bella) – The PrimBella device is an FDA approved nerve stimulator device worn on the underside of the wrist to reduce nausea and vomiting symptoms during pregnancy.
- L. Neurostimulation for the treatment of migraine pain and prevention of migraine headaches – On March 11, 2014, FDA granted de novo 510(k) approval for marketing to Cefaly® (STX-med), Herstal, Belgium), which is a TENS device for the prophylactic treatment of migraine in patients 18 years of age or older.

Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

RATIONALE

This evidence review has been updated regularly with searches of the PubMed database. The most recent literature update was performed through September 27, 2023.

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life (QOL), and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Promotion of greater diversity and inclusion in clinical research of historically marginalized groups (e.g., People of Color [African-American, Asian, Black, Latino and Native American]; LGBTQIA

(Lesbian, Gay, Bisexual, Transgender, Queer, Intersex, Asexual); Women; and People with Disabilities [Physical and Invisible]) allows policy populations to be more reflective of and findings more applicable to our diverse members. While we also strive to use inclusive language related to these groups in our policies, use of gender-specific nouns (e.g., women, men, sisters, etc.) will continue when reflective of language used in publications describing study populations.

Transcutaneous Electrical Nerve Stimulation for Chronic Pain

A large number of systematic reviews, most conducted by Cochrane, have assessed the use of transcutaneous electrical nerve stimulation (TENS) in the treatment of a variety of pain conditions, including the topics of osteoarthritis, rheumatoid arthritis, pancreatitis, myofascial trigger points, temporomandibular joint pain, cancer pain, neck pain, acute pain, phantom limb pain, labor pain, and chronic back pain.^{10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26, [28,29,30,31,} In 2010, the American Academy of Neurology (AAN) published an evidence-based review of the efficacy of TENS for the treatment of pain in neurologic disorders, including low back pain and diabetic peripheral neuropathy.^{32,}

Clinical Context and Therapy Purpose

The purpose of TENS is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with chronic pain (eg, musculoskeletal, neuropathic, and mixed pain conditions).

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with chronic pain conditions (eg, musculoskeletal, neuropathic, and mixed pain conditions).

Interventions

The therapy being considered is TENS.

Comparators

The following therapies are currently being used to treat chronic pain: physical therapy and pharmacotherapy.

Outcomes

The general outcomes of interest are reductions in symptoms and medication use, and improvements in functional outcomes and QOL. Given the different types of pain conditions, follow-up will vary and some cases will be life-long (eg, fibromyalgia, arthritis).

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

LOW BACK PAIN

Systematic Reviews

Wu et al (2018) conducted a meta-analysis of RCTs comparing the efficacy of TENS with a control and other nerve stimulation therapies for the treatment of chronic back pain.³³ Reviewers searched 4 databases (PubMed, Cochrane, Google Scholar, ClinicalTrials.gov) and identified 12 RCTs involving 700 patients. Analysis indicated that TENS had efficacy for providing pain relief similar to control treatment (standard mean difference [SMD], -0.20; 95% confidence interval [CI], -0.5 to 0.18; $p=.293$) and that other types of nerve stimulation therapies were more effective than TENS (SMD , 0.86; 95% CI, 0.15 to 1.57; $p=.017$).

Dubinsky et al (2010), who conducted an evidence-based review for AAN, evaluated the efficacy of TENS for treating pain in neurologic disorders.³² The evidence on TENS for chronic low back pain of various etiologies (some neurologic) included 2 class I studies (prospective randomized trial with masked outcome assessment in a representative population) and 3 class II studies (randomized trial not meeting class I criteria or a prospective matched group cohort study in a representative population). The class I studies compared TENS with sham TENS for 4 or 6 weeks of treatment. Although both studies were adequately powered to find a 20% or greater difference in pain reduction by visual analog scale (VAS), after correction for multiple comparisons, no significant benefit was found for TENS compared with sham TENS. In 2 of the 3 class II studies, no significant differences were found between TENS and sham TENS. In the third class II study, the benefit was found in 1 of 11 patients treated with conventional TENS, 4 of 11 treated with burst-pattern TENS, and 8 of 11 treated with frequency-modulated TENS. Overall, evidence was conflicting. Because class I studies provide stronger evidence, AAN considered the evidence sufficient to conclude that TENS is ineffective for the treatment of chronic low back pain.

Cochrane reviews by Khadilkar et al (2005; 2008), concluded that there is limited and inconsistent evidence for the use of TENS as an isolated treatment for low back pain.^{18,19}

Randomized Controlled Trials

Jalavandi et al (2022) compared the effects of TENS ($n=22$) to back exercises (including strengthening and stretching; $n=22$) in operating room nurses with chronic low back pain.³⁴ After 6 weeks, average pain and disability scores significantly decreased in both treatment groups as compared to baseline. After adjusting for the baseline values, the TENS group had a significantly higher pain score reduction (mean difference [MD] , 4.23; $p=.030$) and a significantly greater decrease in the disability scores (MD , -3.99; $p=.021$) when compared to the back exercises group.

Leemans et al (2020) evaluated the effects of heat and TENS in 50 patients with chronic low back pain.³⁵ Patients were randomized to heat plus TENS or no treatment. At 24 hours after the procedure, there was no significant difference between the groups for average pain in the last 24 hours or maximum pain experienced in the last 24 hours. Measurements were repeated at 4 weeks and no significant differences in pain scores were found between groups at that time point either.

Keskin et al (2012) reported on an RCT of TENS for pregnancy-related low back pain.³⁶ Seventy-nine patients were randomized to 6 TENS sessions over 3 weeks, a home exercise program, acetaminophen, or no-treatment control. In the control group, pain intensity increased in 57% of participants. Pain decreased in 95% of participants in the exercise group and in all participants in the acetaminophen and TENS groups. The VAS score improved by a median of 4 points in the TENS group and by 1 point in the exercise and acetaminophen groups. In the control group, the VAS score worsened by 1 point. Roland-Morris Disability Questionnaire scores indicated significantly greater improvement in function in the TENS group (-8.5) compared with the control (+1), exercise (-3), and acetaminophen (-3) groups. This trial lacked a sham TENS control. In a subsequent RCT by Jamison et al (2019) that also lacked a sham control group and had fewer patients (n=33), compared to treatment-as-usual, use of high-frequency TENS along with a smartphone tracking app resulted in greater reductions in pain intensity.³⁷

DIABETIC PERIPHERAL NEUROPATHY

Systematic Reviews

The AAN's 2010 evidence-based review also identified 2 class II studies comparing TENS with sham TENS and 1 class III study comparing TENS with high-frequency muscle stimulation for patients with mild diabetic peripheral neuropathy.³² The studies found a modest reduction in VAS scores for TENS compared with sham, and a larger proportion of patients experiencing benefit with high-frequency muscle stimulation than with TENS. Reviewers concluded that, on the basis of these 2 class II studies, TENS was likely effective in reducing pain from diabetic peripheral neuropathy; however, no studies compared TENS with other treatment options.

Randomized Controlled Trials

A small RCT by Gossrau et al (2011) found no difference between microcurrent TENS (micro-TENS) compared with sham in 41 patients with diabetic peripheral neuropathy.³⁸ In this trial, the current was applied at an intensity of 30 to 40 microamps rather than the usual intensity of several milliamperes, and patients were treated for 30 minutes, 3 times per week. After 4 weeks of treatment, 29% of the micro-TENS group and 53% of the sham group showed a response to therapy, defined as a minimum 30% reduction in neuropathic pain score. Median Pain Disability Index was reduced to a similar extent in the TENS (23%) and sham (25%) groups.

CANCER PAIN

Systematic Reviews

For a Cochrane review by Robb et al (2008), which evaluated TENS for cancer pain, only 2 RCTs (N=64 participants) met the selection criteria.²⁸ There were no significant differences between TENS and placebo in the included studies. One RCT found no differences between TENS and placebo for pain secondary to breast cancer treatment. The other RCT examined acupuncture-type TENS in palliative care patients but was underpowered. The results of the review were considered inconclusive due to a lack of suitable RCTs. A 2012 update of the Cochrane review identified an additional RCT (a feasibility study of 24 patients with cancer bone pain) that met selection criteria.¹⁷ The small sample sizes and differences in patient study populations across the 3 RCTs precluded meta-analysis. Results on TENS for cancer pain remain inconclusive.

FIBROMYALGIA

Systematic Review

Amer-Cuenca et al (2023) conducted a systematic review and meta-analysis of TENS for analgesia in patients with fibromyalgia.³⁹ When the 11 included RCTs were analyzed with a random-effects model, there was no effect of TENS on pain ($p > .05$). In contrast, a mixed-effects model that considered the TENS dosage found significant effect sizes with the number of TENS sessions ($p = .005$), TENS frequency ($p = .014$), and TENS intensity ($p = .047$). The authors concluded that TENS can reduce fibromyalgia pain when used at high frequency, high intensity, or for more than 10 sessions. A limitation of the review is that about half of the included studies had a high risk of bias.

Randomized Controlled Trials

A placebo-controlled crossover randomized trial by Dailey et al (2013) investigated the effect of a single treatment of TENS in 41 patients with fibromyalgia.⁴⁰ Patients were blindly allocated to no treatment, active TENS treatment, or placebo treatment. Each treatment arm had therapy once weekly for a 3-week period. Patients rated the average pain intensity before and after treatment on a 0-to-10 scale and found less pain with movement during active TENS than with placebo or no TENS ($p < .05$). Patients also rated fatigue with movement and found that fatigue decreased with active TENS compared with placebo or no TENS ($p < .05$ and $p < .01$, respectively). Pressure pain threshold improvement was significantly greater with active TENS (30% ; $p < .05$) than with placebo (11%) or no TENS (14%).

Another RCT by Lauretti et al (2013) investigated TENS in fibromyalgia.⁴¹ However, there was no comparison between active treatment and placebo reported; only change from baseline within each group was reported. TENS was administered for 20 minutes at 12-hour intervals for 7 consecutive days. In the dual placebo group, VAS pain scores did not improve compared with baseline. Patients who had a single site of active TENS reported a reduction in pain of 2.5 cm ($p < .05$ vs. baseline), and patients in the dual TENS group experienced the greatest reduction in pain (4.2 cm; $p < .02$ vs. baseline). Consumption of medication for pain also decreased significantly from baseline in the single TENS ($p < .05$) and dual TENS groups ($p < .02$). Sleep improvements were reported by 10 patients in the dual TENS group, 8 in the single TENS group, and 4 in the placebo group. Fatigue increased for 3 patients in the placebo group but decreased in 7 patients in the dual TENS group; moreover, fatigue decreased for 5 patients in the single TENS group. No adverse events were reported.

Jamison et al (2021) evaluated the efficacy and safety of a wearable TENS device in adults with fibromyalgia.⁴² In this single-center, parallel-group study, 119 patients were randomly assigned to a wearable TENS device (Quell®; $n = 62$) or a sham device ($n = 57$) for 3 months. The primary outcome measure was the Patient Global Improvement of Change (PGIC), which represents the patient's overall belief about the efficacy of treatment on a 7 point categorical verbal rating scale. Selection of 1 means "no change or condition has gotten worse" to 7 meaning "a great deal better and a considerable improvement that has made all the difference." Overall, no differences were found between active and sham treatment on PGIC scores at 3 months (MD, 0.34; 95% CI, -0.37 to 1.04; $p = .351$) in the intention-to-treat population. In the higher pain sensitivity subgroup, the mean PGIC score at 3 months was 4.05 for active treatment versus 2.86 for sham treatment (MD, 1.19; 95% CI, 0.24 to 2.13; $p = .014$). After 3 months of active treatment, all secondary efficacy measures (eg, disease impact and health-related QOL) exhibited significant within-group improvement compared to pre-treatment baseline. A total of 12 (5 active, 7 sham)

adverse events were reported. Nine of the events were definitely or possibly related to TENS use, but were minor and self-limited. The authors concluded that the study demonstrated modest treatment effects of reduced disease impact, pain, and functional impairment from wearable TENS in patients with fibromyalgia.

REFRACTORY CHRONIC PELVIC PAIN

Observational Data

There is limited literature on the use of TENS for chronic pelvic pain. No RCTs were identified. An observational study by Schneider et al (2013) assessed 60 men consecutively treated with TENS for refractory chronic pelvic pain syndrome.⁴³ TENS was performed at home for 12 weeks with participants keeping a pain diary to calculate VAS scores. A successful treatment response was defined as a 50% or greater reduction in VAS and absolute VAS of less than 3 at the end of treatment. TENS was successful in 29 (48%) patients, and treatment response was sustained at a mean follow-up of 44 months (95% CI, 33 to 56 months). After 12 weeks of treatment, VAS scores decreased significantly ($p < .001$) from 6.6 to 3.9. QOL, assessed by the National Institutes of Health Chronic Prostatitis Symptom Index, improved significantly after 12 weeks of TENS treatment ($p < .001$). No adverse events were reported.

OSTEOARTHRITIS OF THE KNEE

Systematic Reviews

A Cochrane review by Rutjes et al (2009) found that the evidence on TENS for pain relief in patients with osteoarthritis of the knee was inconclusive.²⁹ Included in the review were 18 trials assessing 813 patients; 11 trials used TENS, 4 used interferential current stimulation, 1 used both TENS and interferential current stimulation, and 2 used pulsed electrostimulation. Methodologic quality and quality of reporting were rated as poor. Additionally, there was a high degree of heterogeneity among the trials, and the funnel plot for pain was asymmetrical, suggesting both publication bias and bias from small studies.

Randomized Controlled Trials

Additional randomized trials were published after the Rutjes et al (2009) systematic review.

Reichenbach et al (2022) compared treatment with TENS ($n=108$) to sham TENS ($n=112$) in patients with knee osteoarthritis in 6 outpatient clinics in Switzerland.⁴⁴ The primary outcome of mean Western Ontario and McMaster Universities Arthritis Index (WOMAC) pain subscale score at 3 weeks did not significantly differ between the TENS (2.20) and sham TENS group (2.34; MD, -0.06; 95% CI, -0.41 to 0.29; $p = .74$); there was also no significant between-group difference at 15 weeks (2.53 vs. 2.60, respectively; MD, 0.01; 95% CI, -0.37 to 0.39; $p = .98$).

Cherian et al (2016) compared TENS with the standard of care in the treatment of 70 patients who had knee osteoarthritis; all patients had previously taken part in a prospective 3-month trial of TENS, allowing researchers to collect data on the long-term efficacy of TENS (mean follow-up time, 19 months).⁴⁵ The follow-up study evaluated pain (using a VAS) and function (measured by new Knee Society Scale and Lower-Extremity Functional Scale scores) and a number of secondary outcomes, including medication usage, QOL, device use, and conversion to total knee arthroplasty. For all outcomes, reviewers reported a general trend of improvement for the TENS group compared with the standard of care group; however, no statistical analyses were provided

for secondary outcomes, and several differences were not significant among primary outcomes. When measured from pretreatment to final follow-up, Knee Society Scale ($p=.002$) and Lower-Extremity Functional Scale ($p<.001$) scores were significantly increased for the TENS group. The trial's limitations included its small sample size and possible variance in the amount of medication taken by each patient. Also, the interviews were not conducted in person, meaning that some conclusions about functional improvement were not confirmed by a physical examination.

An RCT by Palmer et al (2014) evaluated 224 participants with osteoarthritis of the knee who were assigned to 1 of 3 interventions: TENS combined with education and exercise ($n=73$), sham TENS combined with education and exercise ($n=74$), or education and exercise alone ($n=77$).⁴⁶ Investigators and participants were blinded to treatment. Participants were treated for 6 weeks and directed to use the TENS device as needed for pain relief. WOMAC pain, function, and total scores improved significantly over time from baseline to 24 weeks but did not vary between groups ($p>.05$). TENS as an adjunct to exercise did not elicit additional benefits.

In another RCT, Vance et al (2012) assessed 75 patients given a single session of high-frequency TENS, low-frequency TENS, or placebo TENS.⁴⁷ All 3 groups reported a reduction in pain at rest and during the Timed Up & Go test, and there were no differences in pain scores between groups.

An RCT by Chen et al (2013) compared intra-articular hyaluronic acid injections with TENS for the management of knee osteoarthritis in 50 participants.⁴⁸ Twenty-seven patients were randomized to hyaluronic acid and received 1 intra-articular injection weekly for 5 weeks. Twenty-three patients in the TENS group received 20-minute sessions of TENS 3 times weekly for 4 weeks. The TENS group exhibited a modest but significantly greater improvement ($p=.03$) than the hyaluronic acid group on VAS pain score (mean final score, 4.17 vs. 5.31, respectively) at 2 weeks, but there was no difference between groups at 2 or 3 months post-treatment. The TENS group also had greater improvement on the Lequesne Index at a 2-week follow-up compared with the hyaluronic acid group (mean final score, 7.78 vs. 9.85, respectively; $p=.01$) and at 3-month follow-up (mean final score, 7.07 vs. 9.2, respectively; $p=.03$). Both treatment groups reported significant improvements from baseline to 3 months on scores in walking time, patient global assessment, and disability in activities of daily life.

RHEUMATOID ARTHRITIS

Systematic Reviews

Two Cochrane reviews (2002, 2003) concluded that outcomes for patients with rheumatoid arthritis treated with TENS were conflicting.^{12,13}

MULTIPLE SCLEROSIS

Systematic Reviews

Sawant et al (2015) reported a systematic review of 4 RCTs of TENS for the management of central pain in multiple sclerosis.⁴⁹ Sample sizes ranged from 10 to 60 patients. One study examined the effect of TENS on upper-extremity pain, and the other 3 studied the effect of TENS on low back pain. The exact electrode placement could not be identified. Effect sizes, extracted from the 4 studies, showed a medium-sized effect of TENS (Hedges' $g=0.35$, $p=.009$). The

overall level of evidence was considered to be GRADE 2. Similar findings were reported in a subsequent review by Amatya et al (2018).⁵⁰

PHANTOM LIMB PAIN

Systematic Reviews

A Cochrane review by Johnson et al (2015) found no RCTs on TENS for phantom limb or stump pain after amputation.⁵¹ Reviewers concluded that the published literature on TENS for phantom limb pain in adults lacked the methodologic rigor and robust reporting needed to assess its effectiveness confidently and that RCT evidence is required.

NECK PAIN

Systematic Reviews

A Cochrane review reported by Martimbianco et al (2019) assessed the evidence of TENS for the treatment of chronic neck pain.²¹ Seven RCTs (N=651) comparing TENS alone or in combination with other treatments versus active or inactive treatments were included. Due to heterogeneity in interventions and outcomes, the results were not pooled for a meta-analysis. There was very low-certainty evidence from 2 trials about the effects of conventional TENS versus sham TENS at short-term (up to 3 months after treatment) follow-up. There was no statistically significant difference in outcomes between groups for pain, as assessed by the VAS, (MD, -0.10; 95% CI, -0.97 to 0.77) and the percentage of participants presenting improvement of pain (relative risk [RR], 1.57; 95% CI, 0.84 to 2.92). The authors concluded that there is insufficient evidence regarding the use of TENS in patients with chronic neck pain.

Randomized Controlled Trials

Martins-de-Sousa et al (2023) conducted an RCT of TENS combined with a therapeutic exercise program in patients with chronic neck pain.⁵² Patients were randomized to 8 sessions of placebo TENS (n=20), high frequency TENS (n=20), or low frequency TENS (n=20). The primary outcome, disability after 8 treatment sessions, was similar between groups (p>.05). Other outcomes including pain intensity at the end of treatment and 4 weeks after the end of treatment were also similar between groups. The small sample size may have limited the power to detect a difference between groups.

Diaz-Pulido et al (2021) compared the effects of manual therapy versus TENS on cervical active mobility and muscle endurance in 90 adults diagnosed with subacute and chronic mechanical neck disorders.⁵³ TENS (n=43) and manual therapy (n=47) interventions each consisted of 10 sessions, provided by primary care physical therapists for 30 minutes on alternate days. Outcome measures included active range of motion and endurance of the neck muscles; evaluated pre- and post-intervention and at 6 month follow-up. Of the 90 participants, 72 completed all interventions. Results revealed that manual therapy yielded a significant improvement in active mobility and endurance at post-intervention. At 6 month follow-up, the differences were only significant in endurance and in sagittal plane active mobility. No significant improvement was noted in the TENS group.

PAIN AFTER STROKE

Systematic Reviews

Evidence on the efficacy of TENS for shoulder pain after stroke was considered inconclusive in a Cochrane review by Price et al (2000).²⁶

PAIN AFTER SPINAL CORD INJURY**Systematic Reviews**

A Cochrane review by Boldt et al (2014) evaluating nonpharmacologic interventions for chronic pain in individuals with spinal cord injury identified an RCT on TENS.⁵⁴ This trial had a high-risk of bias, and no conclusion could be drawn on the effectiveness of TENS compared with sham for reducing chronic pain in this population.

FACIAL MYALGIA**Randomized Controlled Trials**

An RCT by De Giorgi et al (2017) evaluated the efficacy of TENS in treating subjective and objective pain in 49 women diagnosed with chronic facial myalgia; 34 patients received TENS treatment daily for 10 weeks and were evaluated for pain up to 25 weeks, and 15 patients received no treatment and were evaluated for pain up to 10 weeks.⁵⁵ TENS treatment consisted of daily 60-minute sessions at 50 Hz, and VAS scores were taken for average and maximum pain intensity in the previous 30 days, as well as the level of pain at an examination. The other primary outcome was the assessment of pain at muscular palpation sites, measured by the Pericranial Muscle Tenderness Score and Cervical Muscle Tenderness Score. For this outcome and that of VAS (mean and maximum measurements), patients in the TENS group had significantly lower pain levels than those for the control group at 10 weeks ($p < .05$). Within the TENS group, the trialists found that VAS scores tended to decrease during the trial, as did Pericranial Muscle Tenderness and Cervical Muscle Tenderness scores ($p < .05$). These differences were significant except for the period between 15 and 25 weeks. Secondary outcomes included mandibular movement and range of motion, and the TENS group showed no significant improvement over the control group for either outcome. Although a limitation of the trial was that observation of control patients ended at 10 weeks, these results confirmed the results of several similar studies of TENS in treating musculoskeletal pain. The trialists concluded that TENS is an effective treatment for chronic facial myalgia, although studies with more participants are needed.

TEMPOROMANDIBULAR DISORDER**Systematic Reviews**

de Castro-Carletti et al (2023) conducted a systematic review and meta-analysis of controlled trials with electrotherapy for orofacial pain.⁵⁶ The systematic review yielded 43 studies (N=1939) for temporomandibular disorder and none for other types of orofacial pain. The quality of evidence was low, but meta-analysis was performed with 20 studies. Regardless of the type of temporomandibular disorder, TENS did not demonstrate a significant benefit compared to placebo or other forms of electrotherapy for pain intensity, maximal mouth opening, or tenderness. A limitation of the analysis is that almost all studies (n=41) had a high risk of bias.

Serrano-Munoz et al (2023) conducted a systematic review and meta-analysis of electrical stimulation modalities for temporomandibular disorders.⁵⁷ Seven RCTs were included, 4 of which

evaluated TENS. Overall, TENS reduced pain intensity (MD, -1.09; 95% CI, -0.71 to -1.47; $I^2=72\%$). TENS did not have a significant effect on range of movement or muscle activity.

Randomized Controlled Trials

A randomized placebo-controlled trial by Ferreira et al (2017) evaluated TENS in the treatment of individuals with temporomandibular disorder; 40 patients (30 female, 10 male) were randomized into 2 groups (placebo or active TENS).⁵⁸ The trial used both high- and low-frequency TENS, allotting to the active TENS patients 25 minutes of 4 Hz followed by 25 minutes of 100 Hz; measuring pain intensity and pressure pain threshold immediately after treatment and again 48 hours later. When compared with baseline values, pain intensity was reduced for patients in the active TENS group, and pressure pain threshold was significantly increased ($p<.05$). For those in the placebo group, there were no significant improvements for either primary outcome. Limitations of the trial included the short duration of the assessment, and the absence of control groups either receiving no treatment or evaluating the same treatment in patients without the temporomandibular disorder.

MYOFASCIAL TRIGGER POINTS

Systematic Reviews

A systematic review by Ahmed et al (2019) evaluated the effects of various electric stimulation techniques in individuals with myofascial trigger points, including 13 RCTs of TENS compared with sham TENS. High-frequency TENS (>50 Hz) was used in the majority of RCTs. Unclear allocation concealment and blinding were the most common study limitations. Meta-analysis of post-treatment pain intensity scores found that TENS did not significantly reduce pain (SMD, -0.16; 95% CI, 0.39 to 0.07).⁵⁹

Randomized Controlled Trials

Effects of TENS combined with ultrasound were more positive in an RCT by Takla et al (2019) of 70 participants with acute mechanical neck pain and at least 2 active myofascial trigger points. Participants were randomized to 3 sessions per week for 4 consecutive weeks of low-frequency, high-intensity burst TENS combined with ultrasound, medium-frequency, low-intensity amplitude modulated frequency TENS combined with ultrasound or sham combined therapy. Pressure pain threshold and active cervical lateral flexion range of motion were improved in both combined therapy groups - more so in the high-intensity burst TENS combined with ultrasound - but not in the sham group.⁶⁰

MIXED CHRONIC PAIN CONDITIONS

Systematic Reviews

A systematic review and meta-analysis by Johnson et al (2022) investigated TENS for relief of various acute and chronic pain conditions in adults.⁶¹ In total, the review included 381 RCTs (N=24,543), with 164 RCTs having sufficient data for meta-analyses. In the subgroup of patients with chronic pain (31 RCTs; n=1417), TENS reduced pain intensity when compared to placebo (SMD, -0.87; 95% CI, -1.19 to -0.55). The authors concluded that for the overall population of patients with acute and chronic pain, there was moderate-certainty evidence that pain intensity is lower during or immediately after TENS compared with placebo. However, levels of evidence were downgraded because of small-sized trials contributing to imprecision in magnitude estimates.

An overview of a Cochrane review by Gibson et al (2019) evaluated the evidence from 8 Cochrane reviews consisting of 51 RCTs that compared TENS versus sham or usual care/no treatment/waiting list control in 2895 participants with various chronic pain conditions. As with previous reviews, due to the serious methodological limitations described below, authors were unable to draw conclusions about the effects of TENS on pain control, disability, health-related QOL, use of pain-relieving medications, global impression of change, or harms.⁶²

Section Summary: Transcutaneous Electrical Nerve Stimulation for Chronic Pain

For individuals who have chronic pain (eg, musculoskeletal, neuropathic, and mixed pain conditions) who receive TENS, the evidence includes numerous RCTs and systematic reviews. The overall strength of the evidence is weak. The best evidence exists for the treatment of chronic, intractable pain. Systematic reviews have found potential pain relief benefits with TENS for diabetic peripheral neuropathy and fibromyalgia. For low back pain and myofascial trigger points, available evidence suggests that TENS is ineffective. Available evidence from systematic reviews are inconclusive for cancer pain, osteoarthritis of the knee, rheumatoid arthritis, phantom knee pain, chronic neck pain, temporomandibular disorder, pain after stroke, and pain after spinal cord injury.

TRANSCUTANEOUS ELECTRICAL NERVE STIMULATION FOR ACUTE PAIN

Clinical Context and Therapy Purpose

The purpose of TENS is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with acute pain (eg, surgical, musculoskeletal, labor, and mixed pain conditions).

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with acute pain conditions (eg, surgical, musculoskeletal, labor, and mixed pain conditions).

Interventions

The therapy being considered is TENS.

Comparators

The following therapy is currently being used to treat acute pain: pharmacotherapy.

Outcomes

The general outcomes of interest are symptoms, functional outcomes, QOL, and medication use. Given the different types of pain conditions, follow-up at 2, 4, and 6 weeks is of interest to monitor outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

INJURY

Systematic Review

Davison et al (2022) conducted a systematic review of 4 studies that evaluated the effect of electrical stimulation after hip fracture.⁶³ Based on the results of one study, TENS decreased pain as assessed by VAS scores (MD, 3.3 points; $p < .001$), increased range of motion at 10 days (MD, 25.7 degrees; $p < .001$), and improved functional recovery ($p < .001$). Results were conflicting regarding the effects of TENS on muscle strength and mobility. The authors concluded that additional high quality trials were needed.

Randomized Controlled Trials

One double-blind, randomized, sham-controlled trial reported by Lang et al (2007) found that during emergency transport of 101 patients, TENS reduced posttraumatic hip pain (change in VAS score, 89 to 59), whereas the sham-stimulated group remained relatively unchanged (change in VAS score, 86 to 79).⁶⁴

SURGICAL PAIN

Systematic Reviews

Zimpel et al (2020) conducted a systematic review with meta-analysis to investigate the efficacy of various complementary alternative therapies, including TENS, for post-caesarean pain.³¹ Ten studies were included that evaluated TENS, with or without analgesia, for pain relief. One study (N=40) evaluated TENS with no treatment and found that it may reduce pain at 1 hour (MD, -2.26; 95% CI, -3.35 to -1.17). TENS plus analgesia, as compared to placebo plus analgesia, may reduce pain at 1 hour (SMD, -1.10; 95% CI -1.37 to -0.82 based on 3 studies with 238 women). Both findings were rated as low-certainty evidence by the Cochrane review.

Zhu et al (2017) conducted a systematic review with meta-analysis to investigate the efficacy of TENS on patients experiencing pain after total knee arthroplasty.⁶⁵ Two independent investigators searched PubMed, Embase, Web of Sciences, EBSCO, and Cochrane Library databases and identified 6 RCTs that assessed the effect TENS had on VAS scores of 529 patients who had a total knee arthroplasty. A meta-analysis indicated that, compared with a control intervention, TENS significantly reduced VAS scores over a 24-hour period (SMD, -0.47; 95% CI, -0.87 to -0.08; $p = .02$). The study was limited by the number of RCTs and sample sizes (4 of 6 selected RCTs had <100 patients), as well as differences in TENS intensities, differences in follow-up times, the ethnic diversity of patients, and possible unpublished or missing data.

Randomized Controlled Trials

Hatefi et al (2023) conducted a double-blind RCT of TENS for pain associated with chest tube removal in 120 patients who underwent coronary artery bypass grafting.⁶⁶ The 4 treatment groups were TENS, cold compress, TENS plus cold compress, and placebo (room temperature compress plus sham TENS), all administered for 15 minutes before chest tube removal. Mean pain intensity scores were lowest in the combined TENS plus cold compress group compared to the other groups at all time points (during chest tube removal, immediately after removal, and 15 minutes after removal [$p < .001$]). Safety of the intervention was not addressed.

Ramanathan et al (2017) published a prospective RCT of 66 patients having undergone total knee arthroplasty who were assigned to active or placebo TENS. Patients used the device as needed for 2 hours and had follow-up visits 2, 4, and 6 weeks after surgery.⁶⁷ For the primary outcome (reduction of opioid intake), no significant difference was observed between active and placebo TENS groups ($p = .60$). This was also the case for secondary outcomes, which included assessment of pain, function, and clinical outcomes. The trial was limited by a high withdrawal rate (only 66 of 116 patients enrolled completed the trial) and a lack of uniformity in the device settings chosen by patients. The investigators found no significant benefit of TENS treatment following total knee arthroplasty.

Parseliunas et al (2020) evaluated TENS use as a component of multimodal pain control after open inguinal hernia surgery in a randomized, double-blind, placebo-controlled trial.⁶⁸ Eighty male patients with unilateral inguinal hernia treated by elective surgery were enrolled and randomly allocated to TENS ($n = 40$) or placebo-TENS ($n = 40$) on the first postoperative day. The primary outcome measure was the change in pain intensity after each TENS application, using VAS and an algometer. Results revealed a significant reduction in VAS pain scores in the TENS group following the procedure ($p < .001$). Absolute and relative pain relief were significantly improved in the TENS group for pain at rest ($p < .01$), when walking ($p < .01$), and when standing up from the bed ($p < .01$). Administration of additional nonopioid analgesics was reduced in the TENS group on the first and second postoperative days ($p < .001$). No postoperative surgical complications or TENS-related adverse effects were seen.

Smaller studies with a higher risk of bias - often due to lack of a sham TENS group - have tended to support the use of TENS. In an RCT of 48 patients who had undergone abdominal surgery, compared to a control group that did not receive any electrical stimulation, Oztas et al (2019) found significantly lower pain scores and analgesic consumption in patients who underwent TENS.⁶⁹ In an assessor-blinded study of TENS in 74 living kidney donors, Galli et al (2015) found a modest reduction in pain at rest and during the measurement of pulmonary function 1 day postoperatively.⁷⁰ A patient-blinded study post abdominal surgery ($N = 55$) by Tokuda et al (2014) found that the application of TENS for 1 hour per day resulted in a significant reduction in pain, particularly at rest, measured both during and immediately after treatment compared with sham TENS.⁷¹ Pulmonary function (vital capacity, cough peak flow) was also significantly better in the active TENS arm. In a single-blind, randomized trial with 42 patients, Silva et al (2012) assessed the analgesic effect of TENS after laparoscopic cholecystectomy.⁷² Pain improved by a median of 2.4 points after TENS compared with 0.4 points after placebo treatment. The relative risk of nausea and/or emesis was 2.2 times greater for patients in the placebo group. In a double-blind RCT of 40 patients undergoing inguinal herniorrhaphy, DeSantana et al (2008) reported that two 30-minute sessions of TENS at 2 and 4 hours after surgery (vs. sham) reduced both analgesic

use and pain scores when measured up to 24 hours post-surgery.⁷³ Pulmonary function (vital capacity, cough peak flow) was also significantly better in the active TENS arm. One exception comes from a single-blind RCT by Forogh et al (2017) of 70 male athletes, which found that adding 20 sessions of high-frequency TENS for 35 minutes a day to semi-supervised exercise did not significantly improve VAS scores.⁷⁴

BONE MARROW SAMPLING

Randomized Controlled Trials

Tucker et al (2015) reported on a double-blind RCT of TENS administered during bone marrow sampling in 70 patients.⁷⁵ There was no significant difference in a numeric pain score between patients who received strong TENS impulses and the control group that received TENS just above the sensory threshold as reported immediately after the procedure (5.6 vs. 5.7, respectively). Over 94% of patients in both groups felt they benefited from TENS.

LOW BACK PAIN

Systematic Reviews

A systematic review by Binny et al (2019) included 3 placebo-controlled studies with 192 women with acute low back pain. Although a low-quality RCT found that TENS in an emergency-care setting provided clinically worthwhile pain relief for moderate to severe acute low back pain, evidence was inconclusive in the other 2 RCTs. Review authors concluded that, overall, the evidence is insufficient to support or refute the use of TENS for acute low back pain.⁷⁶

Koukoulithras et al (2021) reported a systematic review that included 13 RCTs evaluating the effectiveness of non-pharmaceutical interventions upon pregnancy-related low back pain in 2213 patients.⁷⁷ TENS and muscle relaxation exercises accompanied by music were found to be the most effective interventions, having a statistically significant impact on lumbar pain. There was high heterogeneity among the studies including sample sizes.

DYSMENORRHEA

Systematic Reviews

Arik et al (2020) conducted a meta-analysis evaluating the effectiveness of TENS for primary dysmenorrhea.⁷⁸ Four RCTs (N=260) that compared TENS to a sham device were included in the analysis. Pain, as measured by VAS scores, was statistically reduced in the TENS group compared to the sham group (SMD, 1.384; 95% CI, 0.505 to 2.262).

Randomized Controlled Trials

Guy et al (2022) reported on a crossover RCT that took place in France and compared TENS (n=20) to sham TENS (n=20) for primary dysmenorrhea.⁷⁹ The change in pain intensity (measured using VAS) after the first 2 applications (the primary outcome) was significantly greater with TENS (-36.6) versus sham TENS (-2.6; between-group difference, -34.1; p<.0001).

HYSTEROSCOPY

Randomized Controlled Trials

Platon et al (2020) reported the pain relief effects in 74 patients who were randomized to TENS or morphine 5 mg in the post-anesthesia care unit (PACU) after hysteroscopy.⁸⁰ At PACU discharge, both groups reported a significant reduction in pain, with a decrease of VAS scores from 5.6 to 1.4 in the TENS group and 5.1 to 1.3 in the opioid group. There were no significant differences between groups. Sixteen patients in each group reported a VAS ≥ 3 after initial treatment and were crossed over to receive the other treatment during the study as defined by the protocol.

Lison et al (2017) published an RCT assessing the effect of TENS on pain in women undergoing hysterectomy without sedation; the trial included 138 women receiving active TENS, placebo TENS, or neither treatment during the procedure.⁸¹ Women in the active TENS group reported significantly lower VAS scores than women in the control or placebo TENS groups reported. This was the case at each stage measured (entry, contact, biopsy [when necessary], and residual). To validate these measurements, the investigators included a second pain scale (Likert scale) and found a significant correlation with the VAS results ($p < .001$). For secondary endpoints (eg, procedure duration, vital parameters, vasovagal symptoms), the trialists reported that differences between the groups were not statistically significant. However, patient satisfaction was significantly higher in the active TENS group than in either placebo TENS or control groups ($p < .001$ and $p = .001$, respectively). Trial limitations included the failure to account for the use of a flexible hysteroscope, instead of using a rigid hysteroscope; this might have limited the generalizability of the results.

LABOR AND DELIVERY

Systematic Reviews

A Cochrane review by Deussen et al (2020) included 28 studies involving 2749 women experiencing uterine cramping after vaginal delivery.⁸² There was a very low certainty that TENS is better than no TENS for adequate pain relief as reported by 32 women in 1 applicable RCT.

A systematic review and meta-analysis by Thuvarakan et al (2020) evaluating the efficacy of TENS for labor pain included 26 studies with 3348 patients.⁸³ TENS showed a statistically significant effect in the reduction of pain intensity (pooled RR, 1.52; 95% CI, 1.35 to 1.70). The authors noted that there was high study heterogeneity ($I^2 = 89\%$) and the majority of included studies were judged to be low quality.

A Cochrane review by Dowswell et al (2009) included 19 studies with 1671 women in labor.¹⁶ Overall, there was little difference in pain ratings between TENS and control groups, although women receiving TENS to acupuncture points were less likely to report severe pain (RR, 0.41). Reviewers found limited evidence that TENS reduced pain in labor or had any impact (either positive or negative) on other outcomes for mothers or babies.

Randomized Controlled Trials

Kurata et al (2022) published the results of an RCT comparing TENS (n=60), sham TENS (n=60), and no TENS (n=60) after cesarean birth.⁸⁴ The primary outcome of median opioid consumption within 60 hours of cesarean delivery was 7.5 morphine milligram equivalents (MME) with TENS versus 0 MME with sham TENS ($p = .31$). In the no TENS group, the median opioid consumption within 60 hours of cesarean delivery was 7.5 MME ($p = .57$ vs. sham TENS).

A placebo-controlled, randomized trial by Kayman-Kose et al (2014) assessed 200 women who gave birth between January and July 2010.⁸⁵ One hundred women who gave birth vaginally were allocated to active TENS or sham TENS in a 1:1 ratio; this same assignment was performed for 100 women who gave birth by cesarean delivery. TENS was performed once for 30 minutes after childbirth was completed. After vaginal or cesarean delivery, but before the administration of TENS, the placebo and active groups did not significantly differ in VAS or verbal numeric scale scores. However, after active TENS in the cesarean group, there was a significant reduction in VAS ($p < .001$) and verbal numeric scale ($p < .001$) scores compared with the placebo group. A similar benefit was observed in the vaginal delivery group with the active treatment showing a significant reduction in VAS ($p = .022$) and verbal numeric scale ($p = .005$) scores. The investigators also assessed whether TENS reduced the need for additional analgesia. There was no difference between the active TENS and the placebo groups for vaginal delivery ($p = .83$), but, in the cesarean arm, the active treatment group had a significant reduction in analgesic need ($p = .006$). Results were consistent in a much smaller RCT by Baez Suarez et al (2019) of 10 women in labor with a breech vaginal delivery. In this RCT, only women who received active TENS experienced a clinically significant improvement in VAS scores.⁸⁶

Njogu et al (2021) assessed the effects of TENS during the first stage of labor in a single-blind RCT involving 326 adult pregnant women anticipating spontaneous vaginal delivery.⁸⁷ Enrolled patients were randomly assigned to TENS ($n = 161$) or routine obstetric care ($n = 165$) at the beginning of active labor until the second labor stage. The primary outcome was labor pain intensity as assessed by VAS immediately after randomization, at 30, 60, and 120 minutes after TENS therapy, and 2 to 24 hours post-delivery. Prior to the TENS intervention, there was no statistically significant difference in mean VAS scores between the groups ($p > .05$). The TENS group had significantly lower mean VAS scores as compared to control at all time points post-intervention and at 2 to 24 hours post-delivery (all $p < .0001$). The TENS group had a significantly shorter duration of the active labor phase compared to controls ($p < .001$) and the time of the second and third stages of labor were similar between the groups ($p > .05$). The authors concluded that TENS can be used as a non-pharmacologic therapy to reduce labor pain and shorten the active labor phase duration. Limitations cited were lack of a double-blind, sample size, single-center analysis, and inclusion of only a low-risk pregnancy population.

MEDICAL ABORTION

Randomized Controlled Trials

Goldman et al (2021) evaluated whether the use of TENS reduced pain with medical abortion in a randomized, placebo-controlled trial involving 40 patients.⁸⁸ Enrolled women underwent a medical abortion with mifepristone and misoprostol and were randomly assigned to high-frequency TENS (80 Hz; $n = 20$) or a sham device ($n = 20$) to use at home. The primary outcome was a comparison of maximum pain scores within the first 8 hours after misoprostol administration using an 11-point numeric rating scale. Thirty-seven patients had data evaluable for the primary outcome. Median maximum pain scores within 8 hours after misoprostol were 7 and 10 for the high-frequency TENS and sham device, respectively. Patients administered high-frequency TENS experienced a significant reduction in post-treatment pain score compared to those who were administered the sham device (-2.0 vs. 0 ; $p = .008$). No significant differences between the devices were found with regard to additional analgesia use, distribution of maximum pain scores at 24 hours, adverse effects, or measures of acceptability.

MIXED ACUTE PAIN CONDITIONS

Systematic Reviews

A systematic review by Johnson et al (2022) was previously introduced.⁶¹ In the subgroup of patients with acute pain (57 RCTs; n=3348), TENS significantly reduced pain intensity compared to placebo (SMD, -1.02; 95% CI, -1.24 to -0.79). The authors concluded that for the overall population of patients with acute and chronic pain, there was moderate-certainty evidence that pain intensity is lower during or immediately after TENS compared with placebo. However, levels of evidence were downgraded because of small-sized trials contributing to imprecision in magnitude estimates.

Randomized Controlled Trials

Butera et al (2018) conducted a trial to determine the efficacy of using TENS to reduce musculoskeletal pain and improve function after exercise-induced muscle pain.⁸⁹ In this RCT, 36 patients were divided into 3 groups and received TENS, placebo TENS, or no treatment as a control. Treatment was administered for 90 minutes at 24, 48, and 72 hours after the onset of muscle soreness. Analysis indicated that active TENS and placebo TENS had no significant effect on pain. Limitations included a small sample size of young, relatively healthy individuals.

TENNIS ELBOW

Randomized Controlled Trials

A multicenter RCT of TENS as an adjunct to primary care management for tennis elbow was reported by Chesterton et al (2013).⁹⁰ Thirty-eight general practices in the United Kingdom recruited 241 adults who had a new or first diagnosis of tennis elbow. Participants were randomized to TENS once a day for 45 minutes over 6 weeks or until resolution of pain plus primary care management (consultation with a general practitioner followed by information and advice on exercise) versus primary care management alone. Both groups saw large (>25%) within-group improvements in pain intensity, with the greatest improvement during the first 6 weeks of treatment. Intention-to-treat analysis revealed no difference in improvement of pain (-0.33; 95% CI, -0.96 to 0.31; p=.31) between the 2 groups at 6 weeks, 6 months (-0.20; 95% CI, -0.81 to 0.42; p=.526), or 12 months (0.45; 95% CI, -0.15 to 1.06; p=.139). However, adherence to exercise and TENS was very poor, with only 42 (35%) meeting prior adherence criteria. Per-protocol analyses only showed a statistically significant difference in favor of TENS at 12 months (p=.03).

Section Summary: Transcutaneous Electrical Nerve Stimulation for Acute Pain

The evidence for the use of TENS from high-quality trials remains inconclusive for most indications of acute pain. A systematic review of TENS for acute and chronic pain found some evidence that TENS reduces pain intensity over and above that seen with placebo and other control groups in patients with acute pain, but small-sized trials contributed to imprecision in magnitude estimates. Systematic reviews have found that TENS may help reduce pain in patients with post-operative pain (post-caesarean and total knee arthroplasty), dysmenorrhea, and pain associated with labor and delivery. For low back pain, systematic reviews have found insufficient evidence to support or refute the use of TENS. Randomized controlled trials have reported mixed results in the efficacy of TENS across various acute pain conditions.

ESSENTIAL TREMOR

Clinical Context and Therapy Purpose

The purpose of TENS is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with essential tremor.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with essential tremor.

Interventions

The therapy being considered is TENS of the median nerve. Stimulation of the median nerve has been shown to spike activity in the thalamus.

Comparators

The following therapies are currently being used to treat essential tremor: pharmacotherapy.

Outcomes

The general outcomes of interest are reductions in symptoms and medication use, and improvements in functional outcomes and QOL.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

Nonrandomized Studies

Isaacson et al (2020) evaluated the repeated home use of an FDA-cleared wrist-worn neuromodulation device in the Prospective Study for Symptomatic Relief of Essential Tremor with Cala Therapy (PROSPECT) trial.⁹¹ Key characteristics of the trial are summarized in Table 1. For each active treatment session, the device electrically stimulated the median and radial nerves for 40 minutes with an alternating burst pattern tuned to the frequency of each patient's tremor. The pre-specified co-primary endpoints were improvements on the clinician-rated Tremor Research Group Essential Tremor Rating Assessment Scale (TETRAS) and patient-rated Bain & Findley Activities of Daily Living (BF-ADL) dominant hand scores. Of the 263 enrolled patients, 205 completed the visit 3 follow-up and were included in the primary analysis. Results revealed a significant improvement in TETRAS and BF-ADL from pre- to post-stimulation at each clinic visit ($p < .0001$ for all comparisons). Pre-stimulation tremor levels were improved from Visit 1 to 3 on both TETRAS and BF-ADL ($p < .0001$ for both). Patients rated as "severe" or "moderate" improved with both TETRAS (49.3% at baseline to 21% at study exit) and BF-ADL (64.8% at baseline to

23% at study exit) scoring. Tremor power is a calculation of amplitude and frequency. Tremor power decreases with lower amplitude motions and lower frequency motions. Tremor power was also noted to significantly improve with therapy from pre- to post-stimulation ($p < .0001$). No device-related serious adverse events were reported. Non-serious device-related adverse events occurred in 18% of patients (eg, persistent skin irritation, sore/lesion, discomfort, electrical burns, and minor skin irritation). Conclusions were that the repeated in-home use of this neuromodulation device over 3 months was effective and safe for patients with essential tremor. Limitations identified were the open-label, single-arm design, the lack of consensus for the definition of clinically meaningful improvement in TETRAS or BF-ADL, as well as the exclusion of 58 patients who exited the study early from the pre-specified primary and secondary endpoint analyses.

Table 1. Summary of Key Nonrandomized Trial

Study	Study Type	Country	Participants	Treatment	Follow-Up
Isaacson et al (2020) ⁹¹ ,	Prospective, multicenter, single-arm, open-label	U.S. - 26 sites	263 patients (≥ 22 years) diagnosed with essential tremor having at least 1 dominant hand task scoring ≥ 2 on the clinician-rated TETRAS and ≥ 3 on the self-rated BF-ADL, and having a total score across all dominant hand tasks ≥ 6 on TETRAS and ≥ 8 on BF-ADL	Cala wrist-worn neuromodulation device; patients were instructed to use the device twice daily for 3 months	Three in-clinic visits: Visit 1 (patient screening and enrollment); Visit 2 (1 month follow-up); Visit 3 (3 month follow-up and study completion)

BF-ADL: Bain & Findley Activities of Daily Living; TETRAS: Tremor Research Group Essential Tremor Rating Assessment Scale.

Section Summary: Transcutaneous Electrical Nerve Stimulation for Essential Tremor

The evidence for the use of TENS for essential tremor includes results from a prospective, open-label, post-clearance, single-arm study. Results of this trial suggest that repeated in-home non-invasive neuromodulation therapy is effective and safe for patients with essential tremor. Limitations identified were the open-label, single-arm design, the lack of consensus for the definition of clinically meaningful improvement in TETRAS or BF-ADL, as well as the exclusion of 58 patients who exited the study early from the pre-specified primary and secondary endpoint analyses. Further studies comparing TENS to pharmacologic therapy for essential tremor are needed.

ATTENTION DEFICIT HYPERACTIVITY DISORDER

Clinical Context and Therapy Purpose

The purpose of TENS is to provide a treatment option that is an alternative to or an improvement on existing therapies in patients with attention deficit hyperactivity disorder (ADHD).

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with ADHD (7 to 12 years of age) who are not currently taking prescription ADHD medication.

Interventions

The therapy being considered is TENS. Monarch® external Trigeminal Nerve Stimulation (eTNS) System is based on a purported mechanism of action that the trigeminal nerve stimulates brain areas thought to be involved in ADHD. While the exact mechanism of action is not yet known, neuroimaging studies have shown that eTNS increases activity in the brain regions that are known to be important in regulating attention, emotion, and behavior.

Comparators

The following therapies are currently being used to treat ADHD: pharmacotherapy.

Outcomes

The general outcomes of interest are reductions in symptoms and medication use, and improvements in functional outcomes and QOL.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE**Randomized Controlled Trials**

McCough et al (2019) assessed the efficacy and safety of TENS in a double-blind, sham-controlled pilot study of pediatric patients with ADHD.⁹² Key characteristics of the trial are summarized in Table 2. The study was a 4-week trial followed by 1 blinded week without intervention. Clinical assessments included weekly clinician-administered ADHD-Rating and Clinical Global Impression (CGI) scales, and quantitative electroencephalography (EEG) at baseline and week 4. The primary outcome measure was the clinician completed ADHD-Rating Scale total score. Results revealed that ADHD-Rating Scale totals showed significant group-by-time interactions, demonstrating a differential treatment effect ($F=8.12$; $df=1/228$; $p=.005$). The CGI-Improvement scale also favored active treatment over sham ($p=.003$). Quantitative EEG readings were obtained in both groups but there was no participant specific correlations to other outcomes. No serious adverse events were observed in either group and no patient withdrew from the study due to adverse events. Significant increases in weight and pulse were seen with active TENS over the trial period; however, no differences between active and sham TENS with regard to blood pressure were seen. Conclusions were that TENS therapy is efficacious and well-tolerated in pediatric patients with ADHD. Limitations cited were sample size and short duration of treatment and follow-up.

Table 2. Summary of Key RCT Characteristics

Study	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
McGough et al (2019) ⁹² ,	US	1	NR	62 patients (8 to 12 years) with ADHD based on the KSADS and clinical interview with a minimum total of 24 on the clinician-administered parent ADHD-IV Rating Scale, baseline CGI-S ≥ 4 , and full-scale IQ ≥ 85 . Children were medication-free for at least 1 month prior to enrollment.	TENS device (Monarch eTNS System) administered nightly for 4 weeks (n=32)	Sham TENS device administered nightly for 4 weeks (n=30)

ADHD: attention deficit hyperactivity disorder; CGI-S: Clinical Global Impression-Severity; IQ: intelligence quotient; KSADS: Kiddie Schedule for Affective Disorders and Schizophrenia; NR: not reported; RCT: randomized controlled trial; TENS: transcutaneous electrical nerve stimulation.

Section Summary: Transcutaneous Electrical Nerve Stimulation for Attention Deficit Hyperactivity Disorder

The evidence for the use of TENS for ADHD includes an RCT. Results concluded that TENS is an effective and safe treatment option for pediatric patients with ADHD; however, the study included a small patient sample and was of relatively short duration.

MIGRAINE

Clinical Context and Therapy Purpose

The purpose of TENS in individuals with migraine is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with migraine.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with episodic or chronic migraine.

Migraine is categorized as episodic or chronic depending on the frequency of attacks. Generally, episodic migraine is characterized by 14 or fewer headache days per month and chronic migraine is characterized by 15 or more headache days per month.⁹³

Specific International Classification of Headache Disorders⁹⁴, diagnostic criteria are as follows:

- Episodic migraine:
 1. Untreated or unsuccessfully treated headache lasting 4 to 72 hours
 2. Headache has at least 2 of the following characteristics:
 1. Unilateral location
 2. Pulsating quality
 3. Moderate or severe pain intensity
 4. Aggravation by or causing avoidance of routine physical activity
 3. At least 1 of the following during headache:
 1. Nausea and/or vomiting
 2. Photophobia or phonophobia.
- Chronic migraine:
 1. Migraine-like or tension-type headache on 15 or more days per month for more than 3 months
 2. At least 5 headache attacks without aura meet episodic migraine criteria 1 to 3, and/or at least 5 headache attacks with aura meet episodic migraine criteria 2 to 3
 3. On more than 8 days per month for more than 3 months, fulfilling any of the following criteria:
 1. For migraine without aura, episodic migraine criteria 2 and 3
 2. For migraine with aura, episodic migraine criteria 1 and 2
 3. Believed by the patient to be migraine at onset and relieved by a triptan or ergot derivative.

Interventions

The therapy being considered is TENS. Several TENS devices are approved for both prevention and treatment of migraine.

Comparators

The following therapies are currently being used to treat acute migraine due to episodic or chronic migraine: Medical management or no treatment. A number of medications are used to treat acute migraine. First-line therapy for mild or moderate migraine includes oral non-steroidal anti-inflammatory drugs (NSAIDs) or acetaminophen. More severe migraine can be treated through the use of triptans or an NSAID-triptan combination through a variety of routes (eg, oral, nasal [spray or powder], subcutaneous). Antiemetics can be added for migraine accompanied by nausea or vomiting. Other pharmacologic interventions used to treat acute migraine include calcitonin-gene related peptide antagonists, which can be used in patients with an insufficient response or contraindications to triptans, lasmiditan, and dihydroergotamine.

The following therapies are currently being used to prevent acute migraine in individuals with episodic or chronic migraine: medical management or no treatment. A number of medications are used as prevention for migraine. For most adults with episodic migraines who may benefit from preventive therapy, initial therapy with an antiepileptic drug (divalproex sodium, sodium valproate, topiramate) or beta-blockers (metoprolol, propranolol, timolol) is recommended. Frovatriptan may be beneficial as initial therapy for prevention of menstrual-associated migraine.

Antidepressants (eg, amitriptyline, venlafaxine), alternative beta-blockers (atenolol, nadolol), and additional triptans (naratriptan, zolmitriptan for menstrual-associated migraine prevention) may be considered if initial therapy is unsuccessful. For preventive treatment of pediatric migraine, many children and adolescents who received placebo in clinical trials improved and most preventive medications were not superior to placebo. Possibly effective preventive treatment options for children and adolescents may include amitriptyline, topiramate, or propranolol.

Outcomes

For treatment of acute migraine, specific important health outcomes include freedom from migraine pain and bothersome symptoms, restored function (eg, return to normal activities), and patient-assessed global impression of treatment. Examples of relevant outcome measures appear in Table 3. Follow-up over several hours is needed to monitor for treatment effects.

For prevention of acute migraine in individuals with chronic or episodic migraine, specific important health outcomes include reduction of future attack frequency, severity, and duration, improved responsiveness to acute treatments, improved function and reduced disability, and prevention of progression of episodic migraine to chronic migraine. Follow-up over several days to months is needed to monitor for preventive treatment effects.

Table 3. Health Outcome Measures Relevant to Acute Migraine Attack^{93,95,96,}

Outcome	Description
Pain free	No pain at defined assessment time (eg, 2 hours)
Pain relief	Improvement of pain from moderate to severe at baseline to mild or none or pain scale improved at least 50% from baseline at defined assessment time (eg, 2 hours)
Sustained pain free	No pain at initial assessment (eg, 2 hours) and remains at follow-up assessment (eg, 1 day) with no use of rescue medication or relapse (recurrence) within that time frame
Sustained pain relief	Improvement of pain from moderate to severe at baseline to mild or none or pain scale improved at least 50% from baseline at defined assessment time (eg, 2 hours) and remains improved at follow-up assessment (eg, 1 day) with no use of rescue medication or relapse (recurrence) within that time frame
Symptom relief	Improvement of most bothersome symptom(s) from moderate to severe at baseline to mild or none at defined assessment time (eg, 2 hours)
Function relief	Improvement of function from moderate to severe at baseline to mild or none at defined assessment time (eg, 2 hours)
Restored function	No restriction to perform work or usual activities at a defined assessment time (eg, 2 hours)
Global impact of treatment	Patient assessment of functional disability and health-related quality of life using a Likert or other validated scale at a defined assessment time (eg, 2 hours)
Global evaluation of treatment	Patient assessment of overall treatment effect (pain, symptom relief, adverse events) using a Likert or other validated scale at a defined assessment time (eg, 2 hours)

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

TREATMENT OF ACUTE MIGRAINE

Randomized Controlled Trials

Three double-blind, sham-controlled RCTs evaluated TENS for acute migraine treatment (Tables 4 and 5). Two of the studies evaluated healthcare-provider administration of the device during a single episode in emergency departments, and 1 evaluated self-administration of the device at home during acute episodes over a 3-month period.

Chou et al (2019) conducted an RCT of TENS to the trigeminal nerve with the Cefaly device in 106 individuals experiencing migraine headaches with or without aura.⁹⁷ Eligibility criteria specified that participants may have used any acute medications to treat the attack, but not within the 3 hours before enrollment; 29% had treated the current migraine with an acute medication prior to enrollment. Patients received 1 hour of TENS or sham treatment. The primary outcome, mean pain intensity at 1 hour compared to baseline (using a VAS score of 0 to 10), improved by 3.46 ± 2.32 points in the TENS group versus 1.78 ± 1.89 points in the sham group ($p < .0001$). Patients without aura had significant improvement in pain intensity at 1 hour compared to sham ($p = .0006$) but there was no difference between treatments among patients with aura ($p = .06$). Seven minor adverse effects were reported, and there were no serious adverse events.

Hokonek et al (2021) conducted a single center RCT (N=78) to evaluate the use of TENS in individuals presenting to an emergency department with a migraine.⁹⁸ Participants had not received any medication prior to being admitted to the emergency department. Participants were randomized to TENS or a sham device, and their pain was assessed after 20 and 120 minutes. The change in VAS (0 to 100 mm) score from 0 to 20 min was -51.13 ± 2.94 for the TENS group, while the mean VAS score in the sham group was similar between baseline and 20 minutes (73 ± 3 vs. 72 ± 2). The change in VAS (0 to 100 mm) score from 0 to 120 min was -65 ± 25 for the TENS group and -9 ± 2 for the sham group ($p < .001$). Following randomization, 3 participants in the intervention group withdrew due to paresthesia caused by TENS administration and 2 in the control group withdrew due to severe pain; these individuals were not included in the analysis. Domingues et al (2021) evaluated the analgesic efficacy of a portable, disposable, and home self-applied TENS device during migraine attacks.⁹⁹ Participants (74 adults) who had been diagnosed with migraine by a specialist were randomized in this double-blind clinical trial to the active intervention (n=42) or a sham (n=32) with monthly follow-up for 3 months. The primary outcome measure was an evaluation of pain intensity following treatment. Subjects in both groups reported reduced pain scores; with significantly lower pain scores in the intervention

group compared to the sham group ($p=.004$). Patients in the active intervention group also showed a significant improvement in functional disability scores. No adverse effects were reported.

Study limitations are summarized in Tables 6 and 7. Strengths of the RCTs included the use of a sham device and blinded outcome assessment using validated outcome measures. Although short-term pain relief was demonstrated at some time points, the quality of the overall body of evidence was downgraded due to inconsistency of results and heterogeneity in study settings. Supporting evidence from additional RCTs is needed.

Table 4. Summary of Key RCT Characteristics

Study; Trial	Setting	Sites	Dates	Participants	Interventions	
					Active	Comparator
Chou et al (2019) ⁹⁷ ,	Emergency Departments, US	3	2016-2017	Adults (18 to 65 years of age) with IHS-defined acute migraine attack with or without aura for at least 3 hours before enrollment. Participants may have used any acute medications to treat the attack, but not within the 3 hours before enrollment. 29% had treated the current migraine with an acute medication prior to enrollment.	TENS (1 hour) with the Cefaly device (n=52)	Sham TENS (1 hour) using low-frequency pulses (n=54)
Hokonek et al (2021) ⁹⁸ ,	Emergency Department, Turkey	1	June-Oct 2019	Adults (ages 18 to 50 years) with IHS-defined migraine with or without aura, no preventive migraine treatment in the prior 30 days, presenting to the ED with an	TENS (20 minutes) (n=39)	Sham TENS using a device with an empty battery (20 minutes) (n=39)

Study; Trial	Setting	Sites	Dates	Participants	Interventions	
					Active	Comparator
				untreated acute migraine episode. Participants had not received any medication prior to being admitted to the ED.		
Domingues et al (2021) ⁹⁹ ,	Home, Brazil	NR	Nov 2017-Jul 2018	Adults (18 to 65 years of age) with IHS-defined migraine with or without aura. Most participants were under pharmacological treatment for migraine but specifics of treatment for acute episodes during the study period were not reported.	TENS (20 minutes, self-applied at home) (n=42)	Sham TENS using a device with settings that did not meet those required for effective analgesic treatment by TENS devices (20 minutes, self applied at home) (n=32)

ED: emergency department; IHS: International Headache Society; RCT: randomized controlled trial; TENS: transcutaneous electrical nerve stimulation.

Table 5. Summary of Key RCT Results

Study	Pain score	Rescue medication use	Adverse Events
Chou et al (2019) ⁹⁷ ,	N=106	N=106	
TENS	Mean change After 1 hour: -3.46±2.32 After 2 hours: -2.87±2.24 After 24 hours: -3.46±2.65	After 2 hours: 6% After 24 hours: 40%	No serious adverse events. Inability to tolerate paresthesia sensation: 2 discontinued before first 5 mins elapsed 3 discontinued before the full hour

Study	Pain score	Rescue medication use	Adverse Events
Sham TENS	Mean change After 1 hour: -1.78±1.89 After 2 hours: -1.85±1.96 After 24 hours: -2.38±2.27	After 2 hours: 4% After 24 hours: 41%	1 discontinued before first 5 mins elapsed 1 discontinued before the full hour
p for difference	1 hour: <.0001 .2 hours: .028 24 hours: .062	After 2 hours: 66 After 24 hours: 1.0	
Hokonek et al (2021) ⁹⁸ ,	N=78	78	
	Likert-type verbal scale (1=severe pain, 5=more than fine)	Additional analgesic medication required at 120 minutes	
TENS	change in pain intensity 1 hour 4.5	1/39 (2.6%)	3/83 withdrew due to paresthesia caused by TENS administration
Sham TENS	mean at 1 hour 1.2	30/39 (76.9%)	2/83 withdrew due to severe pain
p	<.001	74.3%; 95% CI 59.9% to 87.6%)	
Domingues et al (2021) ⁹⁹ ,	N=74	N=74	
TENS	Median (IQR) Month 1: -3 (-10 to 0) Month 2: -2 (-10 to 0) Month 3: -2 (-10 to 0)	0 (0 to 3)	
Sham TENS	Median (IQR) Month 1: 0 (-7 to 0) Month 2: -2 (-10 to 0) Month 3: -2 (-10 to 0)	1 (9 to 5)	No adverse events or intolerance to the electrical stimuli
p for difference	Month 1: .001 Month 2: <.001 Month 3: .129	.427	No adverse events or intolerance to the electrical stimuli

IQR: interquartile range; NR: not reported; RCT: randomized controlled trial; TENS: transcutaneous electrical nerve stimulation; VAS: visual analogue scale.

Table 6. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Chou et al (2019) ⁹⁷ ,	1. Intended use population is unclear (e.g., treatment naive,				

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
	those with contraindications to medication, or those who have failed pharmacologic treatment)				
Hokonek et al (2021) ⁹⁸ ,	1. Intended use population is unclear (e.g., treatment naive, those with contraindications to medication, or those who have failed pharmacologic treatment)			Pain measure described as "likert-type verbal scale," unclear if validated	
Domingues et al (2021) ⁹⁹ ,	1, 2. Intended use population is unclear (e.g., treatment naive, those with contraindications to medication, or those who have failed pharmacologic treatment); no details on timing or type of treatment of acute attacks during the study period.				Followup was for 3 months. There was no difference between groups in pain score at the 3-month timepoint. Longer follow-up could provide more information about the effectiveness of the device over time.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5: Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 7. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Chou et al (2019) ^{97,}						
Hokonek et al (2020) ^{98,}		1. Authors stated that TENS patients probably felt that the unit was active	1. No mention of registration	5/83 randomized not included in analysis (3 TENS, 2 sham); no ITT analysis		Confidence intervals NR for pain scale difference; post hoc analysis for scores at different timepoints. Table 2 does not provide a footnote to explain data points and no statistical comparison. Text provides means and p-value for pain scores but does not specify timepoint.
Domingues et al (2021) ^{99,}						

TENS: transcutaneous electrical nerve stimulation.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

MIGRAINE PREVENTION

Randomized Controlled Trial

One RCT evaluated TENS for acute migraine prevention in individuals with chronic or episodic migraine (Tables 8 and 9). The Cefaly device for prevention of migraine was evaluated in the Prevention of Migraine using the STS Cefaly (PREMICE) trial (2013).¹⁰⁰ PREMICE was a double-blind, sham-controlled, randomized trial conducted at 5 tertiary care headache clinics in Belgium. Sixty-seven individuals with at least 2 migraine attacks per month were randomized to active (n=34) or sham (n=33) neurostimulation for 3 months, and 59 (88%) completed the trial on protocol. No serious adverse events occurred, although 1 patient discontinued the trial because of a reported device-caused headache. After a 1-month run-in period, patients were instructed to use the device daily for 3 months. Adherence was recorded by the TENS device. Ninety stimulation sessions were expected, but on average, 56 sessions were completed by the active group, and 49 were completed by the sham group. Primary outcome measures were changes in the number of migraine days and the percent of responders.

In the intention-to-treat analysis, the change in the number of migraine days (run-in vs. 3-month) was -2.06 (95% CI, -0.54 to -3.58) for the TENS group and 0.32 (95% CI, -0.63 to +1.27) for the sham group; this difference was not statistically significant (p=.054). The proportion of responders ($\geq 50\%$ reduction in the number of migraine days/month) was 38% (95% CI, 22% to 55%) in the TENS group and 12% (95% CI, 1% to 23%) in the sham group (p=.014). The number of migraine attacks from the run-in period to the 3-month evaluation was significantly lower for the active TENS group (decrease of 0.82 in the TENS group vs. 0.15 in the sham group; p=.044). The number of headache days was lower in the TENS group than in the sham group (decrease of 2.5 vs. 0.2; p=.041). Patients in the active TENS group reported a 36.6% reduction in the number of acute antimigraine drugs taken compared with a 0.5% reduction in the sham group (p=.008). The severity of migraine days did not differ significantly between groups. No adverse effects were reported among the study participants.

Table 8. Summary of Key RCT Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Schoenen et al (2013); PREMICE ¹⁰⁰ ,	Belgium	5	2009-2011	Adults (18 to 65 years of age) with IHS-defined migraine with or without aura and at least 2 migraine attacks per month	TENS (20 minutes daily) for 3 months (n=34)	Sham TENS (20 minutes daily) for 3 months (n=33)

RCT: randomized controlled trial; TENS: transcutaneous electrical nerve stimulation.

Table 9. Summary of Key RCT Results

Study	Change in number of monthly migraine days at month 3	Responders at month 3	Change in antimigraine medication use at month 3
Schoenen et al (2013); PREMICE ¹⁰⁰ ,	N=67	N=67	N=67
TENS	-2.06 (-0.54 to -3.58)	38.24%	-36.6%
Sham TENS	0.32 (-0.63 to 1.27)	12.12%	0.5%
p	.054	.023	.0072

RCT: randomized controlled trial; TENS: transcutaneous electrical nerve stimulation.

Table 10. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-up ^e
Schoenen et al (2013); PREMICE ¹⁰⁰ ,					1. Follow-up limited to 3 months

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

Table 11. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Schoenen et al (2013); PREMICE ¹⁰⁰ ,			1. No mention of registration		4. Power calculated for a different outcome than the outcome described as primary	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias; 5. Other.

^b Blinding key: 1. Participants or study staff not blinded; 2. Outcome assessors not blinded; 3. Outcome assessed by treating physician; 4. Other.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication; 4. Other.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials); 7. Other.

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on

clinically important difference; 4. Other.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated; 5. Other.

Section Summary: Transcutaneous Electrical Nerve Stimulation for Migraine

The evidence for the use of TENS for treatment of acute migraine includes 3 double-blind, sham-controlled RCTs. Two of the RCTs evaluated healthcare-provider administration of a TENS device during a single episode in emergency departments, and 1 evaluated self-administration of the device at home during acute episodes over a 3-month period. The studies conducted in emergency departments showed clinically and statistically significant reductions in pain intensity and medication use within 2 hours of use. The self-administration study had mixed results: The difference in median pain scores before and after treatment was significantly higher in the TENS group at months 1 and 2, but at month 3 the difference was not statistically significant. Function and analgesic medication use did not differ between groups at any time point. Strengths of the RCTs included the use of a sham device and blinded outcome assessment using validated outcome measures. Although short-term pain relief was demonstrated at some time points, the quality of the overall body of evidence was downgraded due to inconsistency of results and heterogeneity in study settings. It is not clear whether the pain intensity reductions demonstrated in emergency department settings would generalize to other settings over longer time periods. Supporting evidence from RCTs is needed. Additionally, based on the existing evidence, it is unclear how TENS would fit into the current migraine treatment pathway, although it could provide benefit for those who do not receive adequate benefit from pharmacologic first- or second-line therapies, or who may have a contraindication to pharmacologic therapies. The specific intended use must be specified in order to adequately evaluate net health benefit.

The evidence for the use of TENS for prevention of acute migraine in individuals with chronic or episodic migraine includes 1 RCT (N = 67) that reported a greater proportion of patients achieving at least a 50% reduction in migraines with TENS than with sham placebo. The RCT also reported modest reductions in the number of total headache and migraine days. This manufacturer-sponsored trial needs corroboration before conclusions can be made about the efficacy of TENS for preventing migraine headaches. Additionally, based on the existing evidence, it is unclear how TENS would fit into the current migraine prevention pathway, although it could provide benefit for those who do not receive adequate benefit from pharmacologic first- or second-line therapies, or who may have a contraindication to pharmacologic therapies.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Academy of Neurology

In 2010, the American Academy of Neurology published an evidence-based review of the efficacy of TENS for the treatment of pain in neurologic disorders.³² The Academy did not recommend TENS for the treatment of chronic low back pain due to lack of proven efficacy (level A, established evidence from 2 class I studies), but stated that TENS should be considered for the treatment of painful diabetic neuropathy (level B, probably effective, based on 2 class II studies).

American College of Physicians

In 2017, the American College of Physicians published guidelines on noninvasive therapies for acute and low back pain.¹⁰¹ No recommendations for TENS were made; the College concluded that "evidence was insufficient to determine the effectiveness" of TENS and that there was no long-range data.

American Congress of Obstetricians and Gynecologists

In 2019 (reaffirmed in 2021), the ACOG guidelines on labor and delivery found that TENS may "help women cope with labor more than directly affect pain scores."¹⁰²

American Society of Anesthesiologists, et al

In 2010, the practice guidelines from the American Society of Anesthesiologists and American Society of Regional Anesthesia and Pain Medicine recommended that TENS be used as part of a multimodal approach to management for patients with chronic back pain and may be used for other pain conditions (eg, neck and phantom limb pain).¹⁰³

National Cancer Institute

The National Cancer Institute's Physician Data Query identifies TENS as a potential nonpharmacological modality for pain control for post thoracotomy pain syndrome.¹⁰⁴

National Comprehensive Cancer Network

National Comprehensive Cancer Network guidelines on adult cancer pain (v 2.2023) indicate that nonpharmacologic interventions, including TENS, may be considered in conjunction with pharmacologic interventions as needed (category 2A).¹⁰⁵

National Institute for Health and Care Excellence

In 2016, the NICE guidance on low back pain indicated that, despite the long history of use of TENS for back pain, the quality of research studies is poor. This guidance recommended against TENS as a treatment.¹⁰⁶

In 2014, the NICE guidance on osteoarthritis care and management in adults indicated that TENS be considered "as an adjunct to core treatments for pain relief."¹⁰⁷

In 2017, the NICE guidance on intrapartum care recommended against the use of TENS for "established labour."¹⁰⁸

North American Spine Society

In 2020, the North American Spine Society clinical guidelines on the diagnosis and treatment of low back pain provided guidance on the effectiveness of different physical medicine and rehabilitation therapies.¹⁰⁹ The guideline noted that there is conflicting evidence that TENS results in improvement in pain or function at short- to medium-term follow-up. The work group

further recommended that randomized clinical trials with long-term follow-up are needed to evaluate the benefits of TENS compared to exercise/physical therapy or as adjunctive use to usual care for low back pain.

In 2011, the North American Spine Society clinical guidelines on the diagnosis and treatment of cervical radiculopathy from degenerative disorders discussed the role of ancillary treatments such as bracing, traction, electrical stimulation, acupuncture, and TENS.¹¹⁰ A consensus statement from the Society recommended that ozone injections, cervical halter traction, and combinations of medications, physical therapy, injections, and traction have been associated with improvements in patient-reported pain in uncontrolled case series. Such modalities may be considered, recognizing that no improvement relative to the natural history of cervical radiculopathy has been demonstrated. There were no specific statements about the role of TENS in this population.

Osteoarthritis Research Society International

In 2014, the guidelines from the Osteoarthritis Research Society International recommended that TENS was inappropriate for use in patients with multi-joint osteoarthritis; moreover, the guidelines suggested that TENS has an uncertain value for the treatment of knee-only osteoarthritis pain.¹¹¹ Updated guidance (2019) on the non-surgical management of knee, hip, and polyarticular osteoarthritis does not address TENS nor include it in their patient-focused treatment recommendations.¹¹²

U.S. Preventive Services Task Force Recommendations

Not applicable.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review are listed in Table 3.

Table 3. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT05939804	The Effect of Transcutaneous Electrical Nerve Stimulation (TENS) Application on Patients' Pain Level and Analgesic Use in Patients Undergoing Hip Replacement	60	Jul 2024
NCT05812885	Transcutaneous Electrical Nerve Stimulation (TENS) and Chronic Low-Back Pain: A Randomized Crossover Trial	34	Dec 2024
NCT05991921	The Effect of TENS Applied in the Early Postpartum Period on Incision Healing, Pain and Comfort	138	Aug 2023
NCT04114149	Effective Postoperative Pain Relief After Laparoscopic Cholecystectomy With TENS Treatment for First Line of Treatment Compared to Conventional Treatment With Opioids	166	Mar 2024
NCT05320432	Transcutaneous Electrical Nerve Stimulation for Pain Control During First Trimester Abortion: a Blinded Randomized Controlled Trial	70	Jan 2024

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Unpublished</i>			
NCT04092088	Effectiveness of Cerebral and Peripheral Electrical Stimulation on Pain and Functional Limitations Associated With Carpal Tunnel Syndrome: A Randomized, Double-blind, Multi-center, Factorial Clinical Trial	180	Oct 2020 (unknown status)
NCT04851938	Evaluation of the Effect of Transcutaneous Electrical Nerve Stimulation Applied in Different Frequencies on Hormone Levels, Birth Pain Perception and Anxiety During Delivery	112	Jun 2021 (unknown status)
NCT02642796	Comparison of the Efficacy of 2 Different Transcutaneous Electrical Nerve Stimulation Application Sites in Reducing Postoperative Pain After Hip Fracture Surgery	120	Sep 2021

NCT: national clinical trial.

CODING

The following codes for treatment and procedures applicable to this policy are included below for informational purposes. This may not be a comprehensive list of procedure codes applicable to this policy.

Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

The code(s) listed below are medically necessary ONLY if the procedure is performed according to the "Policy" section of this document.

CPT/HCPCS	
A4541	Monthly supplies for use of device coded at E0733 (eff 01/01/2024)
A4542	Supplies and accessories for external upper limb tremor stimulator of the peripheral nerves of the wrist (eff 01/01/2024)
A4556	Electrodes (e.g., apnea monitor), per pair
A4557	Lead wires (e.g., apnea monitor), per pair
A4558	Conductive gel or paste, for use with electrical device (e.g., TENS, NMES), per oz.
A4595	Electrical stimulator supplies, 2 leads, per month (e.g., TENS, NMES)
A4630	Replacement batteries, medically necessary, transcutaneous electrical stimulator, owned by patient
A9900	Miscellaneous DME supply, accessory, and/or service component of another HCPCS code
E0720	Transcutaneous electrical nerve stimulation (TENS) device, 2 lead, localized stimulation
E0730	Transcutaneous electrical nerve stimulation (TENS) device, four or more leads, for multiple nerve stimulation
E0731	Form-fitting conductive garment for delivery of TENS or NMES (with conductive fibers separated from the patient's skin by layers of fabric)
E0733	Transcutaneous electrical nerve stimulator for electrical stimulation of the trigeminal nerve (eff 01/01/2024)
E0734	External upper limb tremor stimulator of the peripheral nerves of the wrist (eff 01/01/2024)
E0744	Neuromuscular stimulator for scoliosis
E0745	Neuromuscular stimulator, electronic shock unit
E0761	Nonthermal pulsed high frequency radiowaves, high peak power electromagnetic energy treatment device
E0762	Transcutaneous electrical joint stimulation device system, includes all accessories
E0765	FDA approved nerve stimulator, with replaceable batteries, for treatment of nausea and vomiting
E0769	Electrical stimulation or electromagnetic wound treatment device, not otherwise classified
E1399	Durable medical equipment, miscellaneous

CPT/HCPCS	
G0281	Electrical stimulation, (unattended), to one or more areas, for chronic Stage III and Stage IV pressure ulcers, arterial ulcers, diabetic ulcers, and venous stasis ulcers not demonstrating measurable signs of healing after 30 days of conventional care, as part of a therapy plan of care
G0282	Electrical stimulation, (unattended), to one or more areas, for wound care other than described in G0281
G0295	Electromagnetic therapy, to one or more areas, for wound care other than described in G0329 or for other uses
G0329	Electromagnetic therapy, to one or more areas for chronic Stage III and Stage IV pressure ulcers, arterial ulcers, diabetic ulcers and venous stasis ulcers not demonstrating measurable signs of healing after 30 days of conventional care as part of a therapy plan of care
S8130	Interferential current stimulator, 2 channel
S8131	Interferential current stimulator, 4 channel

There is no specific coding for the Cefaly device. It would most likely be reported with the miscellaneous durable medical equipment code E1399.

REVISIONS	
04-04-2011	<p>In Description section:</p> <ul style="list-style-type: none"> ▪ Removed "Functional Neuromuscular Stimulation (NMES) - Is a method being developed to restore function to patients with damaged or destroyed nerve pathways through use of an orthotic device with microprocessor controlled electrical neuromuscular stimulation." ▪ A stand-alone medical policy for this topic was created entitled Functional Neuromuscular Electrical Stimulation for Home Use. <p>In Policy section:</p> <ul style="list-style-type: none"> ▪ Removed, "Functional Neuromuscular Stimulation: Is denied experimental / investigational as a technique to restore function following nerve damage or nerve injury. This includes its use in the following situations: <ol style="list-style-type: none"> 1. As a technique to provide ambulation in patients with spinal cord injury; or 2. To provide upper extremity function in patients with nerve damage (e.g., spinal cord injury or post-stroke); or 3. To improve ambulation in patients with foot drop caused by nerve damage (e.g., post-stroke or in those with multiple sclerosis)" ▪ A stand-alone medical policy for this topic was created entitled Functional Neuromuscular Electrical Stimulation for Home Use. <p>In Coding section:</p> <ul style="list-style-type: none"> • Updated wording for HCPCS codes: E0730 • Removed CPT code: E0764, E0770 <p>Updated References section.</p>
12-31-2013	<p>Updated Description section.</p> <p>In Policy section:</p> <ul style="list-style-type: none"> ▪ In Item A 2 revised wording from: "Are not medically necessary for non-musculoskeletal pain, including but not limited to headache, visceral abdominal pain, and pelvic pain." to: "Are not medically necessary for: <ol style="list-style-type: none"> a. non-musculoskeletal pain, including but not limited to, visceral abdominal pain, and pelvic pain.

REVISIONS	
	<p>b. headache".</p> <p>In Coding section:</p> <ul style="list-style-type: none"> ▪ Added HCPCS codes: A9900, S8130, S8131 ▪ Added ICD-10 Diagnosis codes <p>Added Rationale section.</p> <p>In Revision section:</p> <ul style="list-style-type: none"> ▪ Removed Revision comments for the following dates: 01-01-2009, 02-11-2009, 03-22-2010, 06-29-2010 <p>References updated</p>
11-05-2014	<p>Policy revisions posted to bcbsks.com web site on 10-06-2014; effective 11-05-2014, 30 days after posting.</p> <p>Updated Description section.</p> <p>In Policy section:</p> <ul style="list-style-type: none"> ▪ Added Item L, "<u>Neurostimulation</u> for the treatment of migraine pain and prevention of migraine headaches (e.g., Cefaly® TENS device): Is denied experimental / investigational." <p>In Coding section:</p> <ul style="list-style-type: none"> ▪ Added HCPCS code E1399 <p>Updated Rationale section.</p> <p>Updated References section.</p>
09-05-2016	<p>Published 08-04-2016. Effective 09-05-2016.</p> <p>In Title section removed "See also: Functional Neuromuscular Electrical Stimulation for Home Use medical policy" as this policy is no longer on bcbsks.com.</p> <p>Description section updated</p> <p>In Policy section:</p> <ul style="list-style-type: none"> ▪ In Item A 1 added "refractory" and "pain (eg, chronic musculoskeletal pain or neuropathic pain)" and removed "intractable or acute post op musculoskeletal pain" to read "May be considered medically necessary for the treatment of refractory chronic pain (eg, chronic musculoskeletal pain or neuropathic pain)" ▪ In Item A 2 added "b. acute post op musculoskeletal pain" ▪ In Item A 2 removed "headache" as it is referenced in another section of the policy. ▪ Added Item A 4 "The use of TENS for any other condition is considered experimental / investigational." ▪ Removed from the TENS section "This policy reflects the long standing accepted standard of care despite lack of evidence of effectiveness." ▪ In Item C removed "when used in the home setting. Evidence is lacking regarding improved health outcomes." ▪ In Item D removed "(such as the RS-4i)" as device is referenced in the description section. ▪ In Item H removed "(Dynatron STS)" as device is referenced in the description section. ▪ In Item I removed "in the home setting:" ▪ In Item J removed "(BioniCare BIO-1000)" as device is referenced in the description section. ▪ In Item K removed "(e.g., Prima Bella)" as device is referenced in the description section. ▪ In Item L removed "(e.g., Cafaly TENS device)" as device is referenced in the description section. <p>Rationale section updated</p> <p>In Coding section:</p> <ul style="list-style-type: none"> ▪ Added HCPCS Codes: G0281, G0282, G0295, G0329 ▪ Removed ICD Codes: M79.643, M79.646

REVISIONS	
	References updated
10-01-2016	In Coding section: <ul style="list-style-type: none"> ▪ ICD-10 Codes Effective 10-01-2016: M25.541, M25.542, M50.121, M50.122, M50.123, M50.321, M50.322, M50.323 ▪ ICD-10 Codes Termed 09-30-2016: M50.12, M50.32
10-01-2017	In Coding section: <ul style="list-style-type: none"> ▪ Added ICD-10 Codes: M48.61, M48.62 ▪ Removed ICD-10 Code: M48.06
02-24-2021	Medical policy was reviewed with no changes made.
10-01-2021	Title Change to Electrical Stimulation Devices for Home and Provider Use In Coding section <ul style="list-style-type: none"> • Deleted ICD-10 code M54.5 (Effective 10-01-2021) • Added ICD-10 code M54.50, M54.41, M54.59 (Effective 10-01-2021)
02-22-2022	Updated Description Section Updated Policy Guidelines <ul style="list-style-type: none"> ▪ Moved product descriptions from under Regulatory Status section to Policy Guidelines Section Updated Rationale Section Updated Coding Section <ul style="list-style-type: none"> ▪ Removed ICD-9 Codes ▪ Added ICD-10 Codes G89.21-G89.8, G89.4, G90.50-G90.59, M54.81-M54.89, M54.9, M79.1, M79.2, R52 Updated References Section
02-27-2024	Updated Description Section Updated Policy Section <ul style="list-style-type: none"> ▪ Added to Section A: A5:TENS is considered experimental / investigational for the prevention or treatment of migraine headache. Updated Rationale Section Updated Coding Section <ul style="list-style-type: none"> ▪ Removed the ICD-10 codes ▪ Added A4541, A4542, E0733 and E0734 (eff. 01-01-2024) Updated References Section

REFERENCES

1. Food and Drug Administration. De Novo Classification Request for Cefaly Device. 2012; http://www.accessdata.fda.gov/cdrh_docs/reviews/K122566.pdf. Accessed September 24, 2023.
2. Food and Drug Administration. Cefaly Dual Device: K173006. 2017; https://www.accessdata.fda.gov/cdrh_docs/pdf17/K173006.pdf. Accessed September 25, 2023.
3. Food and Drug Administration. Cefaly Acute Device: K171446. 2017; https://www.accessdata.fda.gov/cdrh_docs/pdf17/K171446.pdf. Accessed September 26, 2023.
4. Food and Drug Administration. HeadTerm Device: K172450. 2018; https://www.accessdata.fda.gov/cdrh_docs/pdf17/K172450.pdf. Accessed September 23, 2023.

5. Food and Drug Administration. Allive Device: K192773. 2019; https://www.accessdata.fda.gov/cdrh_docs/pdf19/K192773.pdf. Accessed September 27, 2023.
6. Food and Drug Administration. Relivion Device: K203419. February 2021. https://www.accessdata.fda.gov/cdrh_docs/pdf20/K203419.pdf. Accessed October 6, 2023.
7. Cala Health news release. Cala Health receives FDA breakthrough device designation for Cala Trio therapy to treat action tremors in Parkinson's disease. <https://www.businesswire.com/news/home/20201022005276/en/Cala-Health-Receives-FDA-Breakthrough-Device-Designation-for-Cala-Trio%E2%84%A2-Therapy-to-Treat-Action-Tremors-in-Parkinsons-Disease>. Accessed September 27, 2023.
8. FDA news release. FDA permits marketing of first medical device for treatment of ADHD. April 19, 2019. <https://www.fda.gov/news-events/press-announcements/fda-permits-marketing-first-medical-device-treatment-adhd>. Accessed September 27, 2023.
9. Food and Drug Administration. Axon Therapy Device: K210021. 2021; https://www.accessdata.fda.gov/cdrh_docs/pdf21/K210021.pdf. Accessed October 4, 2023.
10. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). TENS or PENS in the treatment of chronic and postoperative pain. TEC Assessments. 1996; Volume 11, Tab 21. PMID
11. Bronfort G, Nilsson N, Haas M, et al. Non-invasive physical treatments for chronic/recurrent headache. *Cochrane Database Syst Rev.* 2004; (3): CD001878. PMID 15266458
12. Brosseau L, Judd MG, Marchand S, et al. Transcutaneous electrical nerve stimulation (TENS) for the treatment of rheumatoid arthritis in the hand. *Cochrane Database Syst Rev.* 2003; 2003(3): CD004377. PMID 12918009
13. Brosseau LU, Pelland LU, Casimiro LY, et al. Electrical stimulation for the treatment of rheumatoid arthritis. *Cochrane Database Syst Rev.* 2002; 2002(2): CD003687. PMID 12076504
14. Cameron M, Lonergan E, Lee H. Transcutaneous electrical nerve stimulation (TENS) for dementia. *Cochrane Database Syst Rev.* 2003; 2003(3): CD004032. PMID 12917999
15. Carroll D, Moore RA, McQuay HJ, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic pain. *Cochrane Database Syst Rev.* 2001; (3): CD003222. PMID 11687055
16. Dowswell T, Bedwell C, Lavender T, et al. Transcutaneous electrical nerve stimulation (TENS) for pain relief in labour. *Cochrane Database Syst Rev.* Apr 15 2009; (2): CD007214. PMID 19370680
17. Hurlow A, Bennett MI, Robb KA, et al. Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults. *Cochrane Database Syst Rev.* Mar 14 2012; 2012(3): CD006276. PMID 22419313
18. Khadilkar A, Milne S, Brosseau L, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic low-back pain. *Cochrane Database Syst Rev.* Jul 20 2005; (3): CD003008. PMID 16034883
19. Khadilkar A, Odebiyi DO, Brosseau L, et al. Transcutaneous electrical nerve stimulation (TENS) versus placebo for chronic low-back pain. *Cochrane Database Syst Rev.* Oct 08 2008; 2008(4): CD003008. PMID 18843638
20. Kroeling P, Gross A, Goldsmith CH, et al. Electrotherapy for neck pain. *Cochrane Database Syst Rev.* Oct 07 2009; (4): CD004251. PMID 19821322

21. Martimbianco ALC, Porfírio GJ, Pacheco RL, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic neck pain. *Cochrane Database Syst Rev*. Dec 12 2019; 12(12): CD011927. PMID 31830313
22. Milne S, Welch V, Brosseau L, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic low back pain. *Cochrane Database Syst Rev*. 2001; (2): CD003008. PMID 11406059
23. Mulvey MR, Bagnall AM, Johnson MI, et al. Transcutaneous electrical nerve stimulation (TENS) for phantom pain and stump pain following amputation in adults. *Cochrane Database Syst Rev*. May 12 2010; (5): CD007264. PMID 20464749
24. Nnoaham KE, Kumbang J. Transcutaneous electrical nerve stimulation (TENS) for chronic pain. *Cochrane Database Syst Rev*. Jul 16 2008; (3): CD003222. PMID 18646088
25. Osiri M, Welch V, Brosseau L, et al. Transcutaneous electrical nerve stimulation for knee osteoarthritis. *Cochrane Database Syst Rev*. 2000; (4): CD002823. PMID 11034768
26. Price CI, Pandyan AD. Electrical stimulation for preventing and treating post-stroke shoulder pain. *Cochrane Database Syst Rev*. 2000; 2000(4): CD001698. PMID 11034725
27. Proctor ML, Smith CA, Farquhar CM, et al. Transcutaneous electrical nerve stimulation and acupuncture for primary dysmenorrhoea. *Cochrane Database Syst Rev*. 2002; 2002(1): CD002123. PMID 11869624
28. Robb KA, Bennett MI, Johnson MI, et al. Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults. *Cochrane Database Syst Rev*. Jul 16 2008; (3): CD006276. PMID 18646140
29. Rutjes AW, Nuesch E, Sterchi R, et al. Transcutaneous electrostimulation for osteoarthritis of the knee. *Cochrane Database Syst Rev*. Oct 07 2009; 2009(4): CD002823. PMID 19821296
30. Walsh DM, Howe TE, Johnson MI, et al. Transcutaneous electrical nerve stimulation for acute pain. *Cochrane Database Syst Rev*. Apr 15 2009; (2): CD006142. PMID 19370629
31. Zimpel SA, Torloni MR, Porfírio GJ, et al. Complementary and alternative therapies for post-caesarean pain. *Cochrane Database Syst Rev*. Sep 01 2020; 9(9): CD011216. PMID 32871021
32. Dubinsky RM, Miyasaki J. Assessment: efficacy of transcutaneous electric nerve stimulation in the treatment of pain in neurologic disorders (an evidence-based review): report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. Jan 12 2010; 74(2): 173-6. PMID 20042705
33. Wu LC, Weng PW, Chen CH, et al. Literature Review and Meta-Analysis of Transcutaneous Electrical Nerve Stimulation in Treating Chronic Back Pain. *Reg Anesth Pain Med*. May 2018; 43(4): 425-433. PMID 29394211
34. Jalalvandi F, Ghasemi R, Mirzaei M, et al. Effects of back exercises versus transcutaneous electric nerve stimulation on relief of pain and disability in operating room nurses with chronic non-specific LBP: a randomized clinical trial. *BMC Musculoskelet Disord*. Mar 26 2022; 23(1): 291. PMID 35337314
35. Leemans L, Elma Ö, Nijs J, et al. Transcutaneous electrical nerve stimulation and heat to reduce pain in a chronic low back pain population: a randomized controlled clinical trial. *Braz J Phys Ther*. 2021; 25(1): 86-96. PMID 32434666
36. Keskin EA, Onur O, Keskin HL, et al. Transcutaneous electrical nerve stimulation improves low back pain during pregnancy. *Gynecol Obstet Invest*. 2012; 74(1): 76-83. PMID 22722614

37. Jamison RN, Wan L, Edwards RR, et al. Outcome of a High-Frequency Transcutaneous Electrical Nerve Stimulator (hfTENS) Device for Low Back Pain: A Randomized Controlled Trial. *Pain Pract.* Jun 2019; 19(5): 466-475. PMID 30636101
38. Gossrau G, Wähler M, Kuschke M, et al. Microcurrent transcutaneous electric nerve stimulation in painful diabetic neuropathy: a randomized placebo-controlled study. *Pain Med.* Jun 2011; 12(6): 953-60. PMID 21627767
39. Amer-Cuenca JJ, Badenes-Ribera L, Biviá-Roig G, et al. The dose-dependent effects of transcutaneous electrical nerve stimulation for pain relief in individuals with fibromyalgia: a systematic review and meta-analysis. *Pain.* Aug 01 2023; 164(8): 1645-1657. PMID 36893318
40. Dailey DL, Rakel BA, Vance CGT, et al. Transcutaneous electrical nerve stimulation reduces pain, fatigue and hyperalgesia while restoring central inhibition in primary fibromyalgia. *Pain.* Nov 2013; 154(11): 2554-2562. PMID 23900134
41. Lauretti GR, Chubaci EF, Mattos AL. Efficacy of the use of two simultaneously TENS devices for fibromyalgia pain. *Rheumatol Int.* Aug 2013; 33(8): 2117-22. PMID 23423539
42. Jamison RN, Edwards RR, Curran S, et al. Effects of Wearable Transcutaneous Electrical Nerve Stimulation on Fibromyalgia: A Randomized Controlled Trial. *J Pain Res.* 2021; 14: 2265-2282. PMID 34335055
43. Schneider MP, Tellenbach M, Mordasini L, et al. Refractory chronic pelvic pain syndrome in men: can transcutaneous electrical nerve stimulation help?. *BJU Int.* Jul 2013; 112(2): E159-63. PMID 23433012
44. Reichenbach S, Jüni P, Hincapié CA, et al. Effect of transcutaneous electrical nerve stimulation (TENS) on knee pain and physical function in patients with symptomatic knee osteoarthritis: the ETRELKA randomized clinical trial. *Osteoarthritis Cartilage.* Mar 2022; 30(3): 426-435. PMID 34826572
45. Cherian JJ, Harrison PE, Benjamin SA, et al. Do the Effects of Transcutaneous Electrical Nerve Stimulation on Knee Osteoarthritis Pain and Function Last?. *J Knee Surg.* Aug 2016; 29(6): 497-501. PMID 26540652
46. Palmer S, Domaille M, Cramp F, et al. Transcutaneous electrical nerve stimulation as an adjunct to education and exercise for knee osteoarthritis: a randomized controlled trial. *Arthritis Care Res (Hoboken).* Mar 2014; 66(3): 387-94. PMID 23983090
47. Vance CG, Rakel BA, Blodgett NP, et al. Effects of transcutaneous electrical nerve stimulation on pain, pain sensitivity, and function in people with knee osteoarthritis: a randomized controlled trial. *Phys Ther.* Jul 2012; 92(7): 898-910. PMID 22466027
48. Chen WL, Hsu WC, Lin YJ, et al. Comparison of intra-articular hyaluronic acid injections with transcutaneous electric nerve stimulation for the management of knee osteoarthritis: a randomized controlled trial. *Arch Phys Med Rehabil.* Aug 2013; 94(8): 1482-9. PMID 23628378
49. Sawant A, Dadurka K, Overend T, et al. Systematic review of efficacy of TENS for management of central pain in people with multiple sclerosis. *Mult Scler Relat Disord.* May 2015; 4(3): 219-27. PMID 26008938
50. Amatya B, Young J, Khan F. Non-pharmacological interventions for chronic pain in multiple sclerosis. *Cochrane Database Syst Rev.* Dec 19 2018; 12(12): CD012622. PMID 30567012
51. Johnson MI, Mulvey MR, Bagnall AM. Transcutaneous electrical nerve stimulation (TENS) for phantom pain and stump pain following amputation in adults. *Cochrane Database Syst Rev.* Aug 18 2015; 8(8): CD007264. PMID 26284511

52. Martins-de-Sousa PH, Fidelis-de-Paula-Gomes CA, Pontes-Silva A, et al. Additional effect of transcutaneous electrical nerve stimulation in a therapeutic exercise program for sedentary with chronic neck pain: A double-blind randomized controlled trial. *Physiother Res Int.* Jan 2023; 28(1): e1978. PMID 36252091
53. Díaz-Pulido B, Pérez-Martín Y, Pecos-Martín D, et al. Efficacy of Manual Therapy and Transcutaneous Electrical Nerve Stimulation in Cervical Mobility and Endurance in Subacute and Chronic Neck Pain: A Randomized Clinical Trial. *J Clin Med.* Jul 23 2021; 10(15). PMID 34362029
54. Boldt I, Eriks-Hoogland I, Brinkhof MW, et al. Non-pharmacological interventions for chronic pain in people with spinal cord injury. *Cochrane Database Syst Rev.* Nov 28 2014; (11): CD009177. PMID 25432061
55. De Giorgi I, Castroflorio T, Sartoris B, et al. The use of conventional transcutaneous electrical nerve stimulation in chronic facial myalgia patients. *Clin Oral Investig.* Jan 2017; 21(1): 275-280. PMID 27000071
56. de Castro-Carletti EM, Müggenborg F, Dennett L, et al. Effectiveness of electrotherapy for the treatment of orofacial pain: A systematic review and meta-analysis. *Clin Rehabil.* Jul 2023; 37(7): 891-926. PMID 36594219
57. Serrano-Muñoz D, Beltran-Alacreu H, Martín-Caro Álvarez D, et al. Effectiveness of Different Electrical Stimulation Modalities for Pain and Masticatory Function in Temporomandibular Disorders: A Systematic Review and Meta-Analysis. *J Pain.* Jun 2023; 24(6): 946-956. PMID 36801166
58. Ferreira AP, Costa DR, Oliveira AI, et al. Short-term transcutaneous electrical nerve stimulation reduces pain and improves the masticatory muscle activity in temporomandibular disorder patients: a randomized controlled trial. *J Appl Oral Sci.* 2017; 25(2): 112-120. PMID 28403351
59. Ahmed S, Plazier M, Ost J, et al. The effect of occipital nerve field stimulation on the descending pain pathway in patients with fibromyalgia: a water PET and EEG imaging study. *BMC Neurol.* Nov 12 2018; 18(1): 191. PMID 30419855
60. Takla MKN. Low-frequency high-intensity versus medium-frequency low-intensity combined therapy in the management of active myofascial trigger points: A randomized controlled trial. *Physiother Res Int.* Oct 2018; 23(4): e1737. PMID 30095858
61. Johnson MI, Paley CA, Jones G, et al. Efficacy and safety of transcutaneous electrical nerve stimulation (TENS) for acute and chronic pain in adults: a systematic review and meta-analysis of 381 studies (the meta-TENS study). *BMJ Open.* Feb 10 2022; 12(2): e051073. PMID 35144946
62. Gibson W, Wand BM, Meads C, et al. Transcutaneous electrical nerve stimulation (TENS) for chronic pain - an overview of Cochrane Reviews. *Cochrane Database Syst Rev.* Apr 03 2019; 4(4): CD011890. PMID 30941745
63. Davison P, Wilkinson R, Miller J, et al. A systematic review of using electrical stimulation to improve clinical outcomes after hip fractures. *Physiother Theory Pract.* Dec 2022; 38(12): 1857-1875. PMID 33890541
64. Lang T, Barker R, Steinlechner B, et al. TENS relieves acute posttraumatic hip pain during emergency transport. *J Trauma.* Jan 2007; 62(1): 184-8; discussion 188. PMID 17215752
65. Zhu Y, Feng Y, Peng L. Effect of transcutaneous electrical nerve stimulation for pain control after total knee arthroplasty: A systematic review and meta-analysis. *J Rehabil Med.* Nov 21 2017; 49(9): 700-704. PMID 28933513
66. Hatefi F, Kazemi M, Manglian P, et al. The effects of cold compress and transcutaneous electrical nerve stimulation on the pain associated with chest tube removal among

- patients with coronary bypass grafting. *J Cardiothorac Surg.* May 25 2023; 18(1): 186. PMID 37231409
67. Ramanathan D, Saleh A, Klika AK, et al. The Use of Transcutaneous Electrical Nerve Stimulation After Total Knee Arthroplasty: A Prospective Randomized Controlled Trial. *Surg Technol Int.* Jul 25 2017; 30: 425-434. PMID 28537354
 68. Parseliunas A, Paskauskas S, Kubiliute E, et al. Transcutaneous Electric Nerve Stimulation Reduces Acute Postoperative Pain and Analgesic Use After Open Inguinal Hernia Surgery: A Randomized, Double-Blind, Placebo-Controlled Trial. *J Pain.* May 2021; 22(5): 533-544. PMID 33309784
 69. Oztas B, Iyigun E. The effects of two different electrical stimulation methods on the pain intensity of the patients who had undergone abdominal surgery with a midline incision: Randomized controlled clinical trial. *Contemp Nurse.* 2019; 55(2-3): 122-138. PMID 31169066
 70. Galli TT, Chiavegato LD, Liebano RE. Effects of TENS in living kidney donors submitted to open nephrectomy: a randomized placebo-controlled trial. *Eur J Pain.* Jan 2015; 19(1): 67-76. PMID 24831862
 71. Tokuda M, Tabira K, Masuda T, et al. Effect of modulated-frequency and modulated-intensity transcutaneous electrical nerve stimulation after abdominal surgery: a randomized controlled trial. *Clin J Pain.* Jul 2014; 30(7): 565-70. PMID 24901753
 72. Silva MB, de Melo PR, de Oliveira NM, et al. Analgesic effect of transcutaneous electrical nerve stimulation after laparoscopic cholecystectomy. *Am J Phys Med Rehabil.* Aug 2012; 91(8): 652-7. PMID 22311059
 73. DeSantana JM, Walsh DM, Vance C, et al. Effectiveness of transcutaneous electrical nerve stimulation for treatment of hyperalgesia and pain. *Curr Rheumatol Rep.* Dec 2008; 10(6): 492-9. PMID 19007541
 74. Forogh B, Aslanpour H, Fallah E, et al. Adding high-frequency transcutaneous electrical nerve stimulation to the first phase of post anterior cruciate ligament reconstruction rehabilitation does not improve pain and function in young male athletes more than exercise alone: a randomized single-blind clinical trial. *Disabil Rehabil.* Mar 2019; 41(5): 514-522. PMID 29117738
 75. Tucker DL, Rockett M, Hasan M, et al. Does transcutaneous electrical nerve stimulation (TENS) alleviate the pain experienced during bone marrow sampling in addition to standard techniques? A randomised, double-blinded, controlled trial. *J Clin Pathol.* Jun 2015; 68(6): 479-83. PMID 25759407
 76. Binny J, Joshua Wong NL, Garga S, et al. Transcutaneous electric nerve stimulation (TENS) for acute low back pain: systematic review. *Scand J Pain.* Apr 24 2019; 19(2): 225-233. PMID 30849052
 77. Koukoulithras I, Stamouli A, Kolokotsios S, et al. The Effectiveness of Non-Pharmaceutical Interventions Upon Pregnancy-Related Low Back Pain: A Systematic Review and Meta-Analysis. *Cureus.* Jan 30 2021; 13(1): e13011. PMID 33728108
 78. Arik MI, Kiloatar H, Aslan B, et al. The effect of TENS for pain relief in women with primary dysmenorrhea: A systematic review and meta-analysis. *Explore (NY).* 2022; 18(1): 108-113. PMID 32917532
 79. Guy M, Foucher C, Juhel C, et al. Transcutaneous electrical neurostimulation relieves primary dysmenorrhea: A randomized, double-blind clinical study versus placebo. *Prog Urol.* Jul 2022; 32(7): 487-497. PMID 35249825
 80. Platon B, Thörn SE, Mannheimer C, et al. Effects of high-frequency, high-intensity transcutaneous electrical nerve stimulation versus intravenous opioids for pain relief after

- hysteroscopy: a randomized controlled study. *Obstet Gynecol Sci.* Sep 2020; 63(5): 660-669. PMID 32717773
81. Lisón JF, Amer-Cuenca JJ, Piquer-Martí S, et al. Transcutaneous Nerve Stimulation for Pain Relief During Office Hysteroscopy: A Randomized Controlled Trial. *Obstet Gynecol.* Feb 2017; 129(2): 363-370. PMID 28079781
 82. Deussen AR, Ashwood P, Martis R, et al. Relief of pain due to uterine cramping/involution after birth. *Cochrane Database Syst Rev.* Oct 20 2020; 10(10): CD004908. PMID 33078388
 83. Thuvarakan K, Zimmermann H, Mikkelsen MK, et al. Transcutaneous Electrical Nerve Stimulation As A Pain-Relieving Approach in Labor Pain: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Neuromodulation.* Aug 2020; 23(6): 732-746. PMID 32691942
 84. Kurata NB, Ghatnekar RJ, Mercer E, et al. Transcutaneous Electrical Nerve Stimulation for Post-Cesarean Birth Pain Control: A Randomized Controlled Trial. *Obstet Gynecol.* Aug 01 2022; 140(2): 174-180. PMID 35852266
 85. Kayman-Kose S, Arioiz DT, Toktas H, et al. Transcutaneous electrical nerve stimulation (TENS) for pain control after vaginal delivery and cesarean section. *J Matern Fetal Neonatal Med.* Oct 2014; 27(15): 1572-5. PMID 24283391
 86. Báez Suárez A, Martín Castillo E, García Andújar J, et al. Evaluation of the effectiveness of transcutaneous nerve stimulation during labor in breech presentation: a case series. *J Matern Fetal Neonatal Med.* Jan 2021; 34(1): 24-30. PMID 30654675
 87. Njogu A, Qin S, Chen Y, et al. The effects of transcutaneous electrical nerve stimulation during the first stage of labor: a randomized controlled trial. *BMC Pregnancy Childbirth.* Feb 24 2021; 21(1): 164. PMID 33627077
 88. Goldman AR, Porsch L, Hintermeister A, et al. Transcutaneous Electrical Nerve Stimulation to Reduce Pain With Medication Abortion: A Randomized Controlled Trial. *Obstet Gynecol.* Jan 01 2021; 137(1): 100-107. PMID 33278292
 89. Butera KA, George SZ, Borsa PA, et al. Prolonged Reduction in Shoulder Strength after Transcutaneous Electrical Nerve Stimulation Treatment of Exercise-Induced Acute Muscle Pain. *Pain Pract.* Nov 2018; 18(8): 954-968. PMID 29505689
 90. Chesterton LS, Lewis AM, Sim J, et al. Transcutaneous electrical nerve stimulation as adjunct to primary care management for tennis elbow: pragmatic randomised controlled trial (TATE trial). *BMJ.* Sep 02 2013; 347: f5160. PMID 23999980
 91. Isaacson SH, Peckham E, Tse W, et al. Prospective Home-use Study on Non-invasive Neuromodulation Therapy for Essential Tremor. *Tremor Other Hyperkinet Mov (N Y).* Aug 14 2020; 10: 29. PMID 32864188
 92. McGough JJ, Sturm A, Cowen J, et al. Double-Blind, Sham-Controlled, Pilot Study of Trigeminal Nerve Stimulation for Attention-Deficit/Hyperactivity Disorder. *J Am Acad Child Adolesc Psychiatry.* Apr 2019; 58(4): 403-411.e3. PMID 30768393
 93. Singh RBH, VanderPluym JH, Morrow AS, et al. Acute Treatments for Episodic Migraine [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2020 Dec. (Comparative Effectiveness Review, No. 239.) Available from: <https://www.ncbi.nlm.nih.gov/books/NBK566246/>. Accessed November 6, 2023.
 94. Ailani J, Burch RC, Robbins MS. The American Headache Society Consensus Statement: Update on integrating new migraine treatments into clinical practice. *Headache.* Jul 2021; 61(7): 1021-1039. PMID 34160823

95. Tassorelli C, Diener HC, Silberstein SD, et al. Guidelines of the International Headache Society for clinical trials with neuromodulation devices for the treatment of migraine. *Cephalalgia*. Oct 2021; 41(11-12): 1135-1151. PMID 33990161
96. Diener HC, Tassorelli C, Dodick DW, et al. Guidelines of the International Headache Society for controlled trials of acute treatment of migraine attacks in adults: Fourth edition. *Cephalalgia*. May 2019; 39(6): 687-710. PMID 30806518
97. Chou DE, Shnayderman Yugrakh M, Winegarner D, et al. Acute migraine therapy with external trigeminal neurostimulation (ACME): A randomized controlled trial. *Cephalalgia*. Jan 2019; 39(1): 3-14. PMID 30449151
98. Hokenek NM, Erdogan MO, Hokenek UD, et al. Treatment of migraine attacks by transcutaneous electrical nerve stimulation in emergency department: A randomized controlled trial. *Am J Emerg Med*. Jan 2021; 39: 80-85. PMID 31983598
99. Domingues FS, Gayoso MV, Sikandar S, et al. Analgesic efficacy of a portable, disposable, and self-applied transcutaneous electrical nerve stimulation device during migraine attacks: A real-life randomized controlled trial. *Pain Pract*. Nov 2021; 21(8): 850-858. PMID 34013542
100. Schoenen J, Vandersmissen B, Jeanette S, et al. Migraine prevention with a supraorbital transcutaneous stimulator: a randomized controlled trial. *Neurology*. Feb 19 2013; 80(8): 697-704. PMID 23390177
101. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. *Ann Intern Med*. Apr 04 2017; 166(7): 514-530. PMID 28192789
102. American Congress of Obstetricians and Gynecologists (ACOG) Committee Opinion Number 766 on Approaches to Limit Intervention During Labor and Birth. February 2019. <https://www.acog.org/Clinical-Guidance-and-Publications/Committee-Opinions/Committee-on-Obstetric-Practice/Approaches-to-Limit-Intervention-During-Labor-and-Birth>. Accessed September 27, 2023.
103. Benzon HT, Connis RT, De Leon-Casasola OA, et al. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology*. Apr 2010; 112(4): 810-33. PMID 20124882
104. National Cancer Institute. Pain (PDQ)-Health Professional Version. 2022. https://www.cancer.gov/about-cancer/treatment/side-effects/pain/pain-hp-pdq#_3. Accessed September 27, 2023.
105. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Adult Cancer Pain. Version 2.2023. https://www.nccn.org/professionals/physician_gls/pdf/pain.pdf. Accessed September 27, 2023.
106. National Institute for Health and Care Excellence (NICE). Low back pain and sciatica in over 16s: assessment and management [NG59]. 2016 (last updated 2020); <https://www.nice.org.uk/guidance/NG59>. Accessed September 25, 2023.
107. National Institute for Health and Care Excellence (NICE). Osteoarthritis in over 16s: diagnosis and management [NG226]. 2022. <https://www.nice.org.uk/guidance/ng226>. Accessed September 26, 2023.
108. National Institute for Health and Care Excellence (NICE). Intrapartum care for healthy women and babies [CG190]. 2014 (last updated 2022); <https://www.nice.org.uk/guidance/cg190>. Accessed September 27, 2023.

109. North American Spine Society. Diagnosis and Treatment of Low Back Pain. 2020. <https://www.spine.org/Portals/0/assets/downloads/ResearchClinicalCare/Guidelines/LowBackPain.pdf>. Accessed September 27, 2023.
110. Bono CM, Ghiselli G, Gilbert TJ, et al. An evidence-based clinical guideline for the diagnosis and treatment of cervical radiculopathy from degenerative disorders. *Spine J*. Jan 2011; 11(1): 64-72. PMID 21168100
111. McAlindon TE, Bannuru RR, Sullivan MC, et al. OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthritis Cartilage*. Mar 2014; 22(3): 363-88. PMID 24462672
112. Bannuru RR, Osani MC, Vaysbrot EE, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthritis Cartilage*. Nov 2019; 27(11): 1578-1589. PMID 31278997
113. Centers for Medicare & Medicaid. National Coverage Determination (NCD) for Transcutaneous Electrical Nerve Stimulators (TENS) (280.13). 2012; <https://www.cms.gov/medicare-coverage-database/view/ncd.aspx?NCDId=354&>. Accessed September 26, 2023.
114. Centers for Medicare & Medicaid. Decision Memo for Transcutaneous Electrical Nerve Stimulation for Chronic Low Back Pain (160.27). 2012; <https://www.cms.gov/medicare-coverage-database/details/nca-details.aspx?NCAId=256&DocID=CAG-00429N&NCDId=63&ncdver=2&bc=gAAAABAAAA&d>. Accessed September 27, 2023.
115. Centers for Medicare and Medicaid. National Coverage Determination (NCD) for Assessing Patient's Suitability for Electrical Nerve Stimulation Therapy (160.7.1). 2006; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=63&ncdver=2&DocID=160.7.1&bc=gAAAABAAAA&>. Accessed September 25, 2023.

OTHER REFERENCES:

1. MCMC board certified Obstetrics and Gynecology with subcertification in Gynecologic Oncology consultant (#2650), December 10, 2009.
2. AMR board certified Physical Medicine and Rehab, Pain Management consultant (#229785), December 21, 2010.