

Medical Policy



Title: Orthopedic Applications of Stem Cell Therapy (Including Allograft and Bone Substitute Products Used With Autologous Bone Marrow)

Related Policies:	<ul style="list-style-type: none"> ▪ <i>Recombinant and Autologous Platelet-Derived Growth Factors for Wound Healing and Other Non-Orthopedic Conditions</i> ▪ <i>Orthopedic Applications of Platelet-Rich Plasma</i> ▪ <i>Autologous Chondrocyte Implantation for Focal Articular Cartilage Lesions</i>
-------------------	---

Professional	Institutional
Original Effective Date: September 19, 2013	Original Effective Date: September 19, 2013
Revision Date(s): September 19, 2013; August 7, 2015; March 2, 2016; August 15, 2017; February 15, 2018; February 18, 2019; February 25, 2021; April 11, 2022; July 1, 2022; February 28, 2023	Revision Date(s): September 19, 2013; August 7, 2015; March 2, 2016; August 15, 2017; February 15, 2018; February 18, 2019; February 25, 2021; April 11, 2022; July 1, 2022; February 28, 2023
Current Effective Date: February 18, 2019	Current Effective Date: February 18, 2019

State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).

The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.

The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.

If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.

Populations	Interventions	Comparators	Outcomes
Individuals: • With cartilage defects	Interventions of interest are: • Stem cell therapy	Comparators of interest are: • Conservative management • Microfracture • Autologous chondrocyte implantation	Relevant outcomes include: • Symptoms • Morbid events • Functional outcomes • Quality of life • Treatment-related morbidity
Individuals: • With meniscal defects	Interventions of interest are: • Stem cell therapy	Comparators of interest are: • Conservative management	Relevant outcomes include: • Symptoms • Morbid events • Functional outcomes • Quality of life • Treatment-related morbidity
Individuals: • With joint fusion procedures	Interventions of interest are: • Stem cell therapy	Comparators of interest are: • Iliac crest bone graft	Relevant outcomes include: • Symptoms • Morbid events • Functional outcomes • Quality of life • Treatment-related morbidity
Individuals: • With osteonecrosis	Interventions of interest are: • Stem cell therapy	Comparators of interest are: • Core decompression	Relevant outcomes include: • Symptoms • Morbid events • Functional outcomes • Quality of life • Treatment-related morbidity

DESCRIPTION

Mesenchymal stem cells (MSCs) have the capability to differentiate into a variety of tissue types, including various musculoskeletal tissues. Potential uses of MSCs for orthopedic applications include treatment of damaged bone, cartilage, ligaments, tendons, and intervertebral discs.

OBJECTIVE

Mesenchymal stem cells (MSCs) have the capability to differentiate into a variety of tissue types, including various musculoskeletal tissues. Potential uses of MSCs for orthopedic applications include treatment of damaged bone, cartilage, ligaments, tendons, and intervertebral discs.

BACKGROUND

Mesenchymal Stem Cells

Mesenchymal stem cells (MSCs) are multipotent cells (also called multipotent stromal cells) that can differentiate into various tissues including organs, trabecular bone, tendon, articular cartilage, ligaments, muscle, and fat. MSCs are associated with the blood vessels within the bone marrow, synovium, fat, and muscle, where they can be mobilized for endogenous repair as occurs with the healing of bone fractures. Tissues such as cartilage, tendon, ligaments, and vertebral discs show limited capacity for endogenous repair because of the limited presence of the triad of functional tissue components: vasculature, nerves, and lymphatics. Orthobiologics is a term introduced to describe interventions using cells and biomaterials to support healing and repair. Cell therapy is the application of MSCs directly to a musculoskeletal site. Tissue engineering techniques use MSCs and/or bioactive molecules such as growth factors and scaffold combinations to improve the efficiency of repair or regeneration of damaged musculoskeletal tissues.¹

Bone marrow aspirate is considered the most accessible source and, thus, the most common place to isolate MSCs for the treatment of musculoskeletal disease. However, harvesting MSCs from bone marrow requires a procedure that may result in donor-site morbidity. Also, the number of MSCs in bone marrow is low, and the number and differentiation capacity of bone marrow-derived MSCs decreases with age, limiting their efficiency when isolated from older patients.

In vivo, the fate of stem cells is regulated by signals in the local 3-dimensional microenvironment from the extracellular matrix and neighboring cells. It is believed the success of tissue engineering with MSCs will also require an appropriate 3-dimensional scaffold or matrix, culture conditions for tissue-specific induction, and implantation techniques that provide appropriate biomechanical forces and mechanical stimulation. The ability to induce cell division and differentiation without adverse effects, such as the formation of neoplasms, remains a significant concern. Given that each tissue type requires different culture conditions, induction factors (signaling proteins, cytokines, growth factors), and implantation techniques, each preparation must be individually examined.

REGULATORY STATUS

The U.S. Food and Drug Administration (FDA) regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation, Title 21, parts 1270 and 1271. MSCs are included in these regulations.

The regulatory status of the stem cell or stem cell-containing products addressed in this review is summarized below.

Concentrated autologous MSCs do not require approval by the FDA. No products using engineered or expanded MSCs have been approved by the FDA for orthopedic applications.

The following products are examples of commercialized demineralized bone matrix (DBM) products. They are marketed as containing viable stem cells. In some instances, manufacturers have received communications and inquiries from the FDA related to the appropriateness of their

marketing products that are dependent on living cells for their function. The following descriptions are from the product literature.

- AlloStem® (AlloSource) is a partially demineralized allograft bone seeded with adipose-derived MSCs.
- Map3® (RTI Surgical) contains cortical cancellous bone chips, DBM, and cryopreserved multipotent adult progenitor cells (MAPC®).
- Osteocel Plus® (NuVasive) is a DBM combined with viable MSCs isolated from allogeneic bone marrow.
- Trinity Evolution Matrix™ (Orthofix) is a DBM combined with viable MSCs isolated from allogeneic bone marrow.
- Other products contain DBM alone and are designed to be mixed with bone marrow aspirate:
 - Fusion Flex™ (Wright Medical) is a dehydrated moldable DBM scaffold (strips and cubes) that will absorb autologous bone marrow aspirate;
 - Ignite® (Wright Medical) is an injectable graft with DBM that can be combined with autologous bone marrow aspirate.

A number of DBM combination products have been cleared for marketing by the FDA through the 510(k) process. FDA product code: MQV.

Table 1 provides a representative sample of these products; some of which are specifically labeled for mixing with bone marrow aspirate.

Table 1. Demineralized Bone Matrix Products Cleared by FDA

Product	Matrix Type	Mix With Autologous MSCs	Manufacturer or Sponsor	Date Cleared	510(k) No.
Vitoss® Bioactive Foam Bone Graft Substitute	Type I bovine collagen	X	Stryker	Nov 2008	K083033
NanOss BVF-E	Nanocrystalline hydroxyapatite		Pioneer Surgical	Aug 2008	K081558
OrthoBlast® II Demineralized bone matrix putty and paste	Human cancellous bone chips		SeaSpine	Sep 2007	K070751
CopiOs® Bone Void Filler (sponge and powder disc)	Type I bovine dermal collagen	X	Kensey Nash	May 2007	K071237
DBX® Demineralized bone matrix putty, paste and mix	Processed human bone and sodium hyaluronate	X	Musculoskeletal Transplant Foundation	Dec 2006	K053218

Product	Matrix Type	Mix With Autologous MSCs	Manufacturer or Sponsor	Date Cleared	510(k) No.
Integra MOZAIK™ Osteoconductive Scaffold-Putty	Human cancellous bone	X	IsoTis OrthoBiologics	Dec 2006	K062353
Formagraft™ Collagen Bone Graft Matrix	Bovine fibrillary collagen	X	R and L Medical	May 2005	K050789
DynaGraft® II Gel and Putty	Processed human bone particles		IsoTis Orthobiologics	Mar 2005	K040419

FDA: U.S. Food and Drug Administration; MSCs: mesenchymal stem cells.

In 2020, the FDA updated their guidance on "Regulatory Considerations for Human Cells, Tissues, and Cellular and Tissue-Based Products: Minimal Manipulation and Homologous Use."²

Human cells, tissues, and cellular and tissue-based products (HCT/P) are defined as human cells or tissues that are intended for implantation, transplantation, infusion, or transfer into a human recipient. If an HCT/P does not meet the criteria below and does not qualify for any of the stated exceptions, the HCT/P will be regulated as a drug, device, and/or biological product and applicable regulations and premarket review will be required.

An HCT/P is regulated solely under section 361 of the PHS Act and 21 CFR Part 1271 if it meets all of the following criteria:

- "1) The HCT/P is minimally manipulated;
- 2) The HCT/P is intended for homologous use only, as reflected by the labeling, advertising, or other indications of the manufacturer's objective intent;
- 3) The manufacture of the HCT/P does not involve the combination of the cells or tissues with another article, except for water, crystalloids, or a sterilizing, preserving, or storage agent, provided that the addition of water, crystalloids, or the sterilizing, preserving, or storage agent does not raise new clinical safety concerns with respect to the HCT/P; and
- 4) Either:
 - i) The HCT/P does not have a systemic effect and is not dependent upon the metabolic activity of living cells for its primary function; or
 - ii) The HCT/P has a systemic effect or is dependent upon the metabolic activity of living cells for its primary function, and: a) Is for autologous use; b) Is for allogeneic use in a first-degree or second-degree blood relative; or c) Is for reproductive use."

The FDA does not consider the use of stem cells for orthopedic procedures to be homologous use.

POLICY

- A. Mesenchymal stem-cell therapy is considered **experimental / investigational** for all orthopedic applications, including use in repair or regeneration of musculoskeletal tissue.
- B. Allograft bone products containing viable stem cells, including, but not limited to, demineralized bone matrix with stem cells, are considered **experimental / investigational** for all orthopedic applications.
- C. Allograft or synthetic bone graft substitutes that must be combined with autologous blood or bone marrow are considered **experimental / investigational** for all orthopedic applications.

POLICY GUIDELINES

This policy does not address unprocessed allograft bone.

Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

RATIONALE

This evidence review has been updated regularly with searches of the PubMed database. The most recent literature update was performed through December 5, 2022.

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life (QOL), and ability to function, including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The RCT is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Promotion of greater diversity and inclusion in clinical research of historically marginalized groups (e.g., People of Color [African-American, Asian, Black, Latino and Native American]; LGBTQIA

(Lesbian, Gay, Bisexual, Transgender, Queer, Intersex, Asexual); Women; and People with Disabilities [Physical and Invisible]) allows policy populations to be more reflective of and findings more applicable to our diverse members. While we also strive to use inclusive language related to these groups in our policies, use of gender-specific nouns (e.g., women, men, sisters, etc.) will continue when reflective of language used in publications describing study populations.

CARTILAGE DEFECTS

Clinical Context and Therapy Purpose

The purpose of stem cell therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies in patients with osteoarthritis (OA) or focal cartilage defects.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with OA or focal cartilage defects.

Interventions

The therapy being considered is treatment with mesenchymal stem cells (MSCs).

Comparators

Comparators of interest include conservative management with medication or hyaluronic acid (HA) injection, microfracture, and autologous chondrocyte implantation.

Outcomes

The general outcomes of interest are symptoms, morbid events, functional outcomes, QOL, and treatment-related morbidity (TRM). Specific scales may include the:

- Knee Injury and Osteoarthritis Outcome Score (KOOS; 5 subscales with 0-100 scale),
- Lysholm Knee Scale (LKS) score (0-100 scale),
- Tegner Activity Score (TAS); a visual analog scale (VAS) for pain (0-100 mm or 0-10 cm scale),
- Western Ontario and McMaster Universities Arthritis Index (WOMAC) which has 3 subscores: pain, which includes 5 items; stiffness, with 2 items; and physical function, with 17 items.
- WOMAC response criteria is an improvement of 20% in at least 2 items together with an improvement of 10 points in the overall scale.
- Cartilage is evaluated with the Magnetic Resonance Observation of Cartilage Repair Tissue (MOCART, 0-100 points, where higher scores indicate better cartilage repair).
- Follow-up over months to years is of interest for relevant outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

Systematic Reviews

A systematic review and meta-analysis by Borakati et al (2017) included 15 comparative studies (N=582) on the use of MSCs to treat OA or focal osteochondral lesions.³ The studies (13 published and 2 unpublished data) included 5 RCTs, 1 case-control, and 9 cohort studies. A majority of the studies were conducted in Asia, and the source of the MSCs varied (bone marrow, blood, amniotic fluid, adipose tissue). The largest trial had only 56 participants, giving low statistical power for the individual studies. The overall quality of the evidence was considered low, with 3 studies rated as "satisfactory" and the rest rated "poor" on the Jadad scale. Pain assessment results were noted for each of the controlled studies, resulting in a pooled standardized mean difference of -1.27 (95% confidence interval [CI], -1.95 to -0.58) in favor of the group treated with MSCs. Reviewers reported a Z-statistic effect size of 3.62, again in favor of the groups treated with MSCs ($p < .001$); although there was high heterogeneity across controlled studies ($I^2 = 92\%$). There was also suggestion of publication bias; the investigators found 79 trials on clinicaltrials.gov, of which only 3 were listed as 'complete with results,' many trials had been inactive for several years, and 9 had 'unknown' status.

A systematic review and meta-analysis by Maheshwer et al (2020) identified 25 studies with 439 participants that used MSCs for treatment of OA.⁴ Although 13 studies were considered level I RCTs by the authors (range of 7 to 40 participants), low quality RCTs would normally be downgraded to level II. Meta-analysis suggested improvement in self-reported function, but only in patients who underwent concomitant surgery, and there was no significant improvement in pain. Few studies reported on cartilage quality. Most of the studies were rated as poor or fair quality. Conclusions are limited due to substantial variability in MSC source, preparation, and concentration in the current literature.

Wiggers et al (2021) conducted a systematic review of RCTs evaluating autologous mesenchymal stem cell therapy on patient-reported outcome measures and disease severity.⁵ Fourteen RCTs were identified in searches conducted through December 2020. Meta-analysis was precluded because most of the original trial data were not available for pooling and due to heterogeneity across studies. A total of 408 patients with knee osteoarthritis received MSC therapy derived from bone marrow, adipose tissue, or activated peripheral blood. After 1 year, 19 of 26 (73%) clinical outcome measures improved with MSCs compared with control. In the MSC group, patients improved by 1.8 to 4.4 points on the Visual Analogue Scale (0 to 10) and 18 to 32 points on the Knee Osteoarthritis Outcome Score (0 to 100). Four studies showed better disease severity on imaging after MSC compared with control at 1 year. Although the reviewers found a positive effect of autologous MSC therapy compared with control treatments, the certainty of the evidence was rated low to very low due to high risk of bias in the included studies (e.g., 10 of 14 RCTs were at high risk of bias on all outcomes) and high heterogeneity in the source, method of preparation, and dosage of injected stem cells in included RCTs.

A more focused systematic review and meta-analysis of 6 RCTs (N=203) that evaluated cultured MSCs for OA was reported by Kim et al (2020).⁶ Four of the studies used bone marrow-derived MSCs, 1 used adipose-derived cells, and the other cultured placental cells. Only 2 of the 6 studies were rated as low risk of bias. Pain outcomes measured with VAS and WOMAC pain scales were improved at 6 to 12 months, but there was no significant improvement in measures of WOMAC function or cartilage measured by magnetic resonance imaging.

Jin et al (2022) also conducted a more focused systematic review and meta-analysis of 6 RCTs (N=452) that evaluated intra-articular MSC injection in patients undergoing high tibial osteotomy (HTO).⁷ Results demonstrated that there were no significant differences in the International Knee Documentation Committee (IKDC) score and KOOS Pain and Symptoms subscales in patients who underwent HTO with or without the MSC injection. However, patients who received MSC injection had significantly greater improvements in Lysholm scores (mean difference, 2.55; 95% CI, 0.70 to 4.40; $p=.007$), and greater proportions of International Cartilage Regeneration and Joint Preservation Society (ICRS) grade 1 ($p=.03$) and grade 2 ($p=.02$) cartilage repair in the medial femoral condyle and grade 2 cartilage repair in the tibial plateau ($p=.04$).

The source of MSCs may have an impact on outcomes, but this is not well-understood, and the available literature uses multiple sources of MSCs. Because of the uncertainty over whether these products are equivalent, the evidence is grouped by the source of MSC.

MESENCHYMAL STEM CELLS EXPANDED FROM BONE MARROW

Autologous Bone Marrow

Wakitani et al (2002) first reported on the use of expanded MSCs for repair of cartilage defects.⁸ Cells from bone marrow aspirate of 12 patients with OA knees were culture-expanded, embedded in collagen gel, transplanted into the articular cartilage defect, and covered with autologous periosteum at the time of HTO. Clinical improvement did not differ between the experimental group and a group of 12 control patients who underwent HTO alone. Wakitani et al (2007) have since published several cases of patients treated for isolated cartilage defects, with clinical improvement reported at up to 27 months.⁹ However, most of the defects appear to have been filled with fibrocartilage. A report from Wakitani et al (2011) was a follow-up safety study of 31 of the 41 patients (3 patients had died, 5 had undergone total knee arthroplasty) who had received MSCs for articular cartilage repair in their clinics between 1998 and 2008.¹⁰ At a mean of 75 months (range, 5-137 months) since the index procedure, no tumors or infections were identified. Functional outcomes were not reported.

A publication from Centeno et al (2010) of *Regenerative Sciences* in the United States described the use of percutaneously injected culture-expanded MSCs obtained from the iliac spine in 226 patients.¹¹ Following harvesting, cells were cultured with autologous platelet lysate and reinjected under fluoroscopic guidance into peripheral joints ($n=213$) or intervertebral discs ($n=13$). Culture-expanded MSCs requires approval by the U.S. Food and Drug Administration (FDA) and is no longer offered in the United States.

The largest study included in the systematic review by Borakati et al (2017) was by Wong et al (2013), who reported on an RCT of cultured MSCs in 56 patients with OA who underwent medial opening wedge HTO and microfracture of a cartilage lesion (See Tables 2 and 3)¹². Patients received an intra-articular injection of MSCs suspended in HA, or for controls, intra-articular injection of HA alone. The primary outcome was the IKDC score at 6 months, 1 year, and 2 years. Secondary outcomes were the TAS and LKS scores through 2 years and the MOCART scoring system by magnetic resonance imaging (MRI) at 1 year. All patients completed the 2-year follow-up. After adjusting for age, baseline scores, and time of evaluation, the group treated with MSCs showed significantly better scores on the IKDC (mean difference, 7.65 on 0-100 scale; $p=.001$), LKS (mean difference, 7.61 on 0-100 scale; $p=.02$), and TAS (mean difference, 0.64 on

a 0-10 scale; $p=.02$) scores. The clinical significance of these differences is uncertain. Blinded analysis of MRI results found higher MOCART scores in the MSC group. The group treated with MSCs had a higher proportion of patients who had complete cartilage coverage of their lesions (32% vs. 0%), greater than 50% cartilage cover (36% vs. 14%), and complete integration of the regenerated cartilage (61% vs. 14%).

Emadedin et al (2018) reported a triple-blind, placebo-controlled, phase 1/2 trial of expanded MSCs in 47 patients with OA of the knee.¹³ Compared to the placebo group, the MSC group showed statistically significant improvements in WOMAC pain and function subscales but not VAS. The WOMAC stiffness subscale improved to a similar extent in the 2 groups. Minimum Clinically Important Improvement and Patient Acceptable Symptom State were not significantly different between the 2 groups. Study limitations included the short duration of follow-up, statistical analysis, and lack of information regarding use of analgesic medications (see Tables 4 and 5).

Another phase 1/2 RCT of expanded MSCs was reported by Lamo-Espinosa et al (2016, 2018) in 30 patients with OA of the knee.^{14,15} Two doses of MSCs (10×10^6 , 100×10^6) were administered with HA and compared to injection of HA alone. VAS scores were significantly decreased in both MSC groups compared to baseline throughout the 12 months of follow-up, while the decrease in VAS in the control group was not statistically significant. Similarly, total WOMAC scores were statistically decreased only in the high dose group at 12 months. Four-year follow-up was available for 27 of the 30 participants. Two patients in the control group and 1 patient in the low dose group had undergone total knee arthroplasty. VAS scores were higher than at baseline in the HA control group but remained low in the 2 MSC groups. WOMAC scores at the long-term follow-up showed a similar course (see Table 3). Limitations of this study are described in Tables 4 and 5.

Table 2. Summary of Key RCT Characteristics

Study; Trial	Countries	Sites	Dates	Participants	Interventions	
					Active	Comparator
Wong et al (2013) ¹²	Asia	1	NR	Patients with OA who underwent HTO and microfracture (N=56)	Microfracture followed by expanded MSCs suspended in HA	Microfracture plus HA alone
Emadedin et al (2018) ¹³	Iran	1	2012-2016	Patients who met the ACR clinical and radiological criteria for knee OA (N=47)	40x10 ⁶ expanded MSCs with serum albumin (n=22)	Placebo (n=25)
Lamo-Espinosa et al (2016, 2018) ^{14,15}	Spain	2	2012-2014	Patients who met the ACR clinical and radiological criteria for	One of 2 doses of expanded MSCs with HA 10x10 ⁶ , 100x10 ⁶	HA alone

Study; Trial	Countries	Sites	Dates	Participants	Interventions
				knee OA (N=30)	

ACR: American College of Rheumatology; HA: hyaluronic acid; HTO: high tibial osteotomy; MSC: mesenchymal stem cell; NR: not reported; OA: osteoarthritis; RCT: randomized controlled trial.

Table 3. Summary of Key RCT Results

Study					
Wong et al (2013) ¹² ,	IKDC at 6 mo	IKDC at 2 yr	Tegner Activity Scale at 2 yr	Lysolm Knee Score at 2 yr	MOCART
N	56	56	56	56	56
Diff (95% CI)	7.65 (3.04 to 12.26)		0.64 (0.10 to 1.19)	7.61 (1.44 to 13.79)	19.6 (10.5 to 28.6)
p-Value	.001		.021	.016	<.001
Emadedin et al (2018) ¹³ ,	<i>WOMAC Total</i>	<i>WOMAC Pain</i>	<i>WOMAC Stiffness</i>	<i>WOMAC Function</i>	<i>VAS</i>
N	43	43	43	43	43
MSC (95% CI)	-25.7 (-35.4 to 16)	-35 (-44.9 to 25)	-16.9 (-30.4 to 3.5)	-22.9 (-32.9 to 12.9)	-20.8 (-34.5 to 7.1)
Placebo (95% CI)	5.5 (-2.8 to 13.8)	-12.2 (-18.5 to 5.9)	-13.1 (-20.7 to 5.4)	-9.5 (-21.8 to 2.7)	-15.7 (-33.9 to 2.4)
Diff (95% CI)	-13.5 (-24.3 to 2.7)	-21.8 (-33.8 to 9.9)	-7.4 (-25.4 to 10.5)	-11.3 (-22.1 to 0.4)	-5 (-28.1 to 18)
p-Value	.01	.001	.40	.04	.65
Effect size (95% CI)	0.7 (0.1 to 1.4)	1.1 (0.4 to 1.7)		0.6 (0.03 to 1.2)	
Lamo-Espinosa et al (2016, 2018) ^{14,15} ,	<i>WOMAC Total at 12 mo, median (IQR)</i>	<i>WOMAC Total at 4 yr, median (IQR)</i>	<i>VAS at 4 yr, median (IQR)</i>		
MSC low dose	21.5 (15, 26)	17 (13, 25.5)	2 (2, 5)		
MSC high dose	16.5 (12, 19)	16.5 (8, 23)	3 (3, 4)		
Control	13.5 (8, 33)	27 (17, 30)	7 (6, 7)		

CI: confidence interval; IKDC: International Knee Documentation Committee score; IQR: interquartile range; MOCART; Magnetic Resonance Observation of Cartilage Repair Tissue; MSC: mesenchymal stem cell; RCT: randomized controlled trial; VAS: visual analog scale; WOMAC: Western Ontario and McMaster Universities Arthritis Index.

Table 4. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
Wong et al (2013) ¹² ,	4. The population was restricted to patients younger than 55	4. The intervention included microfracture with/without stem cells			
Emadedin et al (2018) ¹³ ,			2. Did not use an active control and use of analgesics was not reported	1. Evaluation of cartilage was not performed.	1, 2. Follow-up was reported out to 6 mo.
Lamo-Espinosa et al (2016, 2018) ^{14,15} ,				1. Evaluation of cartilage was not performed.	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 5. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Wong et al (2013) ¹² ,	3. Patients selected from 1 of 2 identical envelopes	1, 2, 3. Not blinded except for evaluation of magnetic resonance imaging				
Emadedin et al (2018) ¹³ ,				.	3. Details of power analysis were not reported	1. The authors used non-inferiority compared to placebo and chi-square tests for continuous variables

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Data Completeness ^d	Power ^e	Statistical ^f
Lamo-Espinosa et al (2016, 2018) ^{14,15} ,		1, 2, 3. Not blinded			3. Details of power analysis were not reported	1. The authors used non-parametric tests for within-group comparisons rather than tests for repeated measures

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Mesenchymal Stem Cells from Allogeneic Bone Marrow

Vega et al (2015) reported on a small phase 1/2 RCT of 30 patients with OA unresponsive to conventional treatments.¹⁶ The MSC-treated group received an intra-articular injection of expanded allogeneic bone marrow MSCs from healthy donors, and the control group received an intra-articular injection of HA. Follow-up using standard outcome measures was performed at 3, 6, and 12 months post-injection. In the MSC-treated group, pain scores (VAS and WOMAC) decreased significantly between baseline and the 12-month follow-up, whereas pain scores in the control group did not improve significantly. A significant improvement in cartilage quality in the MSC group was supported by T2 MRI. Not reported was whether the patients or assessors were blinded to treatment.

Mesenchymal Stem Cells from Bone Marrow Aspirate Concentrate

Shapiro et al (2017) reported on the results of a prospective, single-blind, placebo-controlled trial assessing 25 patients with bilateral knee pain from bilateral OA.¹⁷ Patients were randomized to BMAC into 1 knee and to saline placebo into the other. Fifty-two milliliters of bone marrow was aspirated from the iliac crests and concentrated in an automated centrifuge. The resulting BMAC was combined with platelet-poor plasma for injection into the arthritic knee and was compared with a saline injection into the contralateral knee, thereby using each patient as his or her control. Safety outcomes, pain relief, and function as measured by Osteoarthritis Research Society International measures and a VAS score were tracked initially at 1 week, 3 months, and 6 months post-procedure. Study patients experienced a similar relief of pain in both BMAC- and saline-treated arthritic knees.

Adipose-Derived Mesenchymal Stem Cells

Adipose-derived stem cells are multipotent MSCs that can be harvested from multiple anatomic locations and with greater ease than bone marrow-derived MSCs. The literature on adipose-derived MSCs for articular cartilage repair comes primarily from research groups in Korea. One group appears to have been providing this treatment as an option for patients for a number of years. They compared outcomes of this new add-on treatment with those for patients who only received other cartilage repair procedures.

Koh et al (2014) reported on results of an RCT that evaluated cartilage healing after HTO in 52 patients with OA.¹⁸ Patients were randomized via sealed envelopes to HTO with the application of platelet-rich plasma (PRP) or to HTO with the application of PRP plus MSCs. A total of 44 patients completed second-look arthroscopy and 1- and 2-year clinical follow-ups. The primary outcomes were the KOOS (0-100 scale), the LKS score (0-100 scale), and a VAS for pain (0-100 scale). There were statistically significant differences between PRP only and PRP plus MSC on 2 of 5 KOOS subscales: pain (74 vs. 81.2, $p < .001$) and symptoms (75.4 vs. 82.8, $p = .006$), all respectively. There were also statistically significant differences on the final pain score between the PRP only (16.2) and PRP plus MSC groups (10.2; $p < .001$), but the final LKS score did not differ significantly between the PRP only (80.6) and PRP plus MSC groups (84.7; $p = .36$). Articular cartilage healing was rated as improved with MSCs following video review of second-look arthroscopy; blinding of this measure is unclear. There were limitations in study design (small sample size, short duration of follow-up). Also, significant improvements were found only on some outcomes, all significant differences in outcomes were modest in magnitude and, as a result, there is uncertainty about the clinical significance of the findings.

More recently, Zaffagnini et al (2022) reported on results of an RCT that evaluated a single intra-articular injection of microfragmented adipose tissue or PRP in patients (N=118) with knee OA.¹⁹ The primary outcomes were the IKDC subjective score and the KOOS pain subscore at 6 months. Overall, both treatments provided significant improvements from baseline in clinical outcomes, with no significant differences found between treatment groups. The IKDC scores significantly improved from baseline to 6 months, from 41.1 ± 16.3 to 57.3 ± 18.8 with microfragmented adipose tissue, and from 44.8 ± 17.3 to 58.4 ± 18.1 with PRP. The improvement in the KOOS pain subscore from baseline to 6 months was 58.4 ± 15.9 to 75.8 ± 17.4 with microfragmented adipose tissue and 63.5 ± 17.8 to 75.5 ± 16.1 with PRP. As a secondary outcome, more patients in the microfragmented adipose tissue group with moderate/severe knee OA reached the minimal clinically important difference for the IKDC score at 6 months compared with the PRP group (75.0% vs 34.6%, respectively; $p = .005$).

Mesenchymal Stem Cells from Peripheral Blood

A 2013 report from Asia has described a small RCT assessing the use of autologous peripheral blood MSCs for focal articular cartilage lesions. Fifty patients with grade 3 or 4 lesions of the knee joint underwent arthroscopic subchondral drilling followed by 5 weekly injections of HA. Half the patients were randomized to injections of peripheral blood stem cells or no further treatment. The peripheral blood stem cells were harvested after stimulation with recombinant human granulocyte colony-stimulating factor, divided in vials, and cryopreserved. At 6 months after surgery, HA and MSCs were re-administered over 3 weekly injections. At 18 months post-surgery, second-look arthroscopy on 16 patients in each group showed significantly higher histologic scores ($\gg 10\%$) for the MSC group (1066 vs. 957 by independent observers) while blinded

evaluation of MRI scans showed a higher morphologic score (9.9 vs. 8.5). There was no difference in IKDC scores between the 2 groups at 24 months after surgery.

Mesenchymal Stem Cells from Umbilical Cord Blood

Lim et al (2021) reported on a RCT of 114 patients with large, full-thickness cartilage defects (International Cartilage Repair Society grade 4) treated with either a composite of umbilical cord-derived MSCs plus 4% hyaluronate (MSC-HA) or microfracture.²⁰ The study consisted of a 48-week phase 3 clinical trial and a 5-year follow-up study. Of 114 patients randomized, 89 completed the phase 3 trial (78.1%), and 73 were enrolled in the follow-up study (64.0%). The primary outcome, proportion of participants with cartilage restoration equivalent to at least 1 grade improvement on the ICRS Macroscopic Cartilage Repair Assessment at 48-week arthroscopic evaluation, was 97.7% (42/43) in the MSC-HA group and 71.7% (33/46) in the microfracture group (odds ratio, 16.55; 95% CI, 2.06 to 133.03; $p=.001$). Both groups had significantly improved patient-reported pain scores (VAS pain, WOMAC, and IKDC scores) at 48 weeks versus baseline, but there was no significant difference between the 2 groups at this timepoint. From 36 to 60 months after intervention, the significant improvements from baseline were maintained in the MSC-HA group, whereas the improvements in VAS pain and WOMAC deteriorated in the microfracture group. This study had several limitations. There was no intervention group that received MSC alone, the comparator (microfracture) is not considered the standard of care for large, full-thickness cartilage defects, surgeons and participants were not blinded to treatment outcome, and there was high loss to follow-up. These limitations, along with a lack of improvement in patient-reported outcomes in the intervention group at 48 weeks, preclude drawing conclusions about the effectiveness of umbilical cord blood-derived MSCs in this population; higher quality evidence from RCTs is needed.

Section Summary: Cartilage Defects

The evidence on MSCs for cartilage repair is increasing, although nearly all studies to date have been performed outside of the United States with a variety of methods of MSC preparation. Overall, the quality of evidence is low and there is a possibility of publication bias. The strongest evidence base is on MSCs expanded from bone marrow, which includes several phase 1/2 RCTs. Compared to either placebo or an active HA control, treatment with expanded MSCs has been shown to provide some improvement in pain and function. Limitations in these initial trials preclude reaching conclusions, but the results to date do support future study in phase 3 trials. Alternative methods of obtaining MSCs that have been reported in RCTs include peripheral blood, adipose tissue, and bone marrow concentrate. These have been reported in a smaller number of trials and with mixed results. Additional study in a larger sample of patients with longer follow-up would be needed to evaluate the long-term efficacy and safety of these procedures. FDA approval for these methods has also not been obtained.

MENISCAL DEFECTS

Clinical Context and Therapy Purpose

The purpose of stem cell therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies in patients with meniscal defects.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with meniscal defects.

Interventions

The therapy being considered is stem cell therapy.

Comparators

Comparators of interest include conservative management.

Outcomes

The general outcomes of interest are symptoms, morbid events, functional outcomes, QOL, and TRM.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Damage to the meniscal cartilage in the knee is a very common orthopedic injury and predisposes to the development of OA. The tissue is relatively avascular and does not spontaneously heal well.

Whitehouse et al (2017) published a report on techniques of in vitro expansion of autologous-derived MSCs and a case series of the first-in-human implantation to treat meniscal defects in 5 patients.²¹ The regulatory framework in the United Kingdom allows cell manipulation and requires immunohistochemical documentation of the presence and volume of mesenchymal cells. Over the first 12 months postprocedure, 3 of the 5 patients were reported to have clinical symptom relief, which persisted through 24 months. MRI scans showing lack of meniscal displacement were the only other postoperative assessment. The 2 patients who failed to obtain symptom relief at 6 and 12 months had to repeat arthroscopic procedures with meniscectomy.

Vangsnest et al (2014) reported on an industry-sponsored phase 1/2 randomized, double-blind, multicenter Study of Chondrogen - Adult Universal Cell Delivered by Intra-Articular Injection Following Meniscectomy in Patients 18-60 Years (NCT00225095, NCT00702741) of cultured allogeneic MSCs (Chondrogen; Osiris Therapeutics) injected into the knee after partial meniscectomy.²² The 55 patients in this United States study were randomized to intra-articular injection of either 50×10^6 allogeneic MSCs, 150×10^6 allogeneic MSCs in HA, or an HA vehicle control at 7 to 10 days after meniscectomy. The cultured MSCs were derived from BMAC of unrelated donors. At 2-year follow-up, 3 patients in the low-dose MSC group had significantly increased meniscal volume measured by MRI (with an a priori determined threshold of at least 15%) compared with none in the control group or the high-dose MSC group. There was no significant difference between the groups in LKS scores. On subgroup analysis, patients with OA who received MSCs had a significantly greater reduction in pain at 2 years than patients who received HA alone. This trial appears to have been a post hoc analysis and, hence, should be

considered preliminary. No serious adverse events were reported as related to the investigational treatment.

Section Summary: Meniscal Defects

The evidence on the use of MSCs to repair or regenerate damaged meniscal tissue consists of preclinical animal studies, first-in-human uncontrolled implantation of expanded autologous MSCs into meniscal tears, and an early-phase randomized trial of cultured allogeneic MSCs injected into the site of partial meniscectomy. Results are preliminary.

JOINT FUSION PROCEDURES

Clinical Context and Therapy Purpose

The purpose of stem cell therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies in patients with joint fusion procedures.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with joint fusion procedures.

Interventions

The therapy being considered is stem cell therapy.

Comparators

Comparators of interest include iliac crest bone graft.

Outcomes

The general outcomes of interest are symptoms, morbid events, functional outcomes, QOL, and TRM.

Follow-up over months to years is of interest for relevant outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

There is limited evidence on the use of allografts with stem cells for bone fusion of the extremities or spine or the treatment of nonunion. The results of several industry-sponsored, early-phase trials are available.

A prospective, clinical, and radiographic 12-month outcomes study (2016) of patients undergoing single-level anterior cervical discectomy and fusion (ACDF) for symptomatic cervical degenerative disc disease using a novel viable allogeneic stem cell and cancellous bone matrix (Trinity

Evolution) was reported using historical controls as the comparator.²³ The ACDF procedure was performed using the polyetheretherketone interbody spacer and bone graft substitute (Trinity Evolution) in 31 patients at multiple clinical sites. At 6 and 12 months, the primary endpoint of radiographic fusion was evaluated as determined by an independent radiographic review and the fusion rate was 78.6% at 6 months and 93.5% at 12 months. Secondary endpoints included a function as assessed by Neck Disability Index scores, and neck and arm pain as assessed by individual VAS scores. Neck function and neck and arm pain were reported as significantly improved at both 6 and 12 months post-procedure. Reported adverse events included carpal tunnel syndrome, minor pain, numbness, permanent and/or unresolved pain, and swelling. Independent medical adjudication of the 26 adverse events occurring in 31 patients found that no adverse events were definitely or probably related to Trinity Evolution. However, 5 adverse events were found to be possibly related to Trinity Evolution with 3 events of mild severity and 2 of moderate severity.

A similar study (2017) involving several of the same investigators and clinical sites reported on the clinical and radiographic evaluation of an allogeneic bone matrix containing stem cells (Trinity Evolution Viable Cellular Bone Matrix) in patients undergoing 2-level ACDF.²⁴ This study involved 40 patients exposed to the ACDF and bone graft substitute procedure at 2 adjacent disc levels. A panel blinded to clinical outcomes reviewed 12-month dynamic motion plain radiographs and thin-cut computed tomography with multiplanar reconstruction. At 12 months, the per-subject and per-level fusion rates were 89.4% and 93.4%, respectively. The clinical function assessments using the Neck Disability Index and VAS scores were reported to have improved from baseline.

A 2015 prospective, multicenter, open-label clinical trial using a cryopreserved, donor mesenchymal cell scaffold (Trinity Evolution) was performed in subjects undergoing foot and/or ankle arthrodesis with surgeons' preferred technique.²⁵ A total of 103 subjects were prospectively enrolled at 10 participating sites. No restrictions were placed on the diagnosis, which included arthritis (primary OAs, posttraumatic OA, and rheumatoid), deformity, neuropathy (Charcot and diabetic), revision surgery, and degenerative joint disease, and arthrodesis was performed in 171 joints. The per-protocol population consisted of 92 patients at 6 months and 76 patients at 12 months, with 153 and 129 total arthrodeses, respectively. The primary endpoint was fusion at 6 months, as assessed from computed tomography scans and standard radiographs by an independent radiology consultant. At 6 months, the fusion rate for all patients was 68.5% and 81.1% for all joints. American Orthopaedic Foot and Ankle Society Hindfoot Scale scores for disability improved over time.

Eastlack et al (2014) reported on outcomes from a series of 182 patients treated with ACDF using Osteocel Plus in a polyetheretherketone cage and anterior plating.²⁶ At 24 months, 74% of patients (180/249 levels treated) were available for follow-up. These patients had significant improvements in clinical outcomes, with 87% of levels achieved solid bridging, and 92% of levels had a range of motion less than 3°. With 26% loss to follow-up at 24 months and lack of a standard of care control group, interpretation of these results is limited.

Section Summary: Joint Fusion Procedures

The evidence on the use of MSCs as a component of joint fusion procedures primarily comes from industry-sponsored, prospective, open-label procedures. Outcomes included radiologic assessments of fusion, sometimes made independently, and patient-reported measures (e.g., VAS scores). The MSCs used were cryopreserved allogeneic in origin. Presumptive benefits of

allogeneic MSCs are that patients undergoing an orthopedic intervention procedure do not need another graft harvesting procedure and that dose of stem cells can be managed.

OSTEONECROSIS

Clinical Context and Therapy Purpose

The purpose of stem cell therapy is to provide a treatment option that is an alternative to or an improvement on existing therapies in patients with osteonecrosis.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with osteonecrosis.

Interventions

The therapy being considered is therapy with MSCs.

Comparators

Comparators of interest include core decompression.

Outcomes

The general outcomes of interest are symptoms, morbid events, functional outcomes, QOL, and TRM.

Follow-up over months to years is of interest for relevant outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

At least 2 RCTs from Asia have evaluated the use of MSCs for osteonecrosis of the femoral head.

Mesenchymal Stem Cells Concentrated from Bone Marrow Aspirate Concentrate

Sen et al (2012) randomized 40 patients (51 hips) with early-stage femoral head osteonecrosis to core decompression plus concentrated bone marrow MSCs or core decompression alone.²⁷ Blinding of assessments in this small trial was not described. Harris Hip Score was significantly improved in the core decompression plus MSC group compared with the core decompression alone group at 12 months (scores, 83.65 vs. 76.68, $p < .016$) but not at 24 months (scores, 82.42 vs. 77.39; $p = .09$), all respectively. Kaplan-Meier analysis showed improved hip survival in the MSC group (mean, 51.9 weeks) compared with the core decompression group (mean, 46.7 weeks). There were no significant differences between groups in radiographic assessment or MRI results.

Mesenchymal Stem Cells Expanded From Bone Marrow

Zhao et al (2012) reported on a randomized trial that included 100 patients (104 hips) with early-stage femoral head osteonecrosis treated with core decompression and expanded bone marrow MSCs or with core decompression alone.²⁸ At 60 months post-surgery, 2 (3.7%) of the 53 hips treated with MSCs progressed and underwent vascularized bone grafting compared with 10 (23%) of 44 hips in the decompression group who progressed and underwent either vascularized bone grafting (n=5) or total hip replacement (n=5). The MSC group also had improved Harris Hip Scores compared with the control group on independent evaluation (data presented graphically). Lesion volume was also reduced by treatment with MSCs.

Section Summary: Osteonecrosis

Two small RCTs have compared core decompression alone with core decompression plus MSCs in patients with osteonecrosis of the femoral head. Both reported improvement in the Harris Hip Score in patients treated with MSCs, although it was not reported whether the patients or investigators were blinded to the treatment group. Hip survival was significantly improved following treatment with either expanded or concentrated MSCs. The effect appears to be larger with expanded MSCs than with concentrated MSCs. Additional, well-designed RCTs with a large number of patients are needed to permit greater certainty on the efficacy of this treatment for osteonecrosis.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Academy of Orthopaedic Surgeons

A 2020 guideline from American Association of Orthopaedic Surgeons on the management of glenohumeral joint OA, endorsed by several other societies, states that injectable biologics such as stem cells cannot be recommended in the treatment glenohumeral joint OA.²⁹ There was consensus from the panel that better standardization and high-quality evidence from clinical trials is needed to provide definitive evidence on the efficacy of biologics in glenohumeral OA. The strength of evidence was rated as no reliable scientific evidence to determine benefits and harms.

The 2021 guideline on treatment of osteoarthritis of the knee does not address stem cell injections.³⁰

American Association of Neurological Surgeons

In 2014, the American Association of Neurological Surgeons guidelines on fusion procedures for degenerative disease of the lumbar spine relevant to this evidence review have indicated that "The use of demineralized bone matrix (DBM) as a bone graft extender is an option for 1- and 2-

level instrumented posterolateral fusions. Demineralized Bone Matrix: Grade C (poor level of evidence).^{31,}

American College of Rheumatology and Arthritis Foundation

In 2019, guidelines from the American College of Rheumatology and Arthritis Foundation on osteoarthritis (OA) of the hand, hip, and knee gave a strong recommendation against stem cell injections in patients with knee and/or hip OA, noting the heterogeneity in preparations and lack of standardization of techniques.³² No recommendation was made for hand OA, since efficacy of stem cells has not been evaluated.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 6.

Table 6. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT02582489	Prospective, Randomized, Double-blind Clinical Trial to Investigate the Efficacy of Autologous Bone Marrow Aspirate Concentrate Post-Meniscectomy	100	Dec 2022
NCT03067870	Transplantation of Autologous Purified Bone Marrow Derived Specific Populations of Stem Cells and Mesenchymal Stem Cells in Patients With Rheumatoid Arthritis	100	Feb 2022
NCT04368806 ^a	A 48-Weeks, Phase 2b/3a, Double-Blind, Randomized, Placebo Controlled, Multi-center, Superiority Study to Evaluate the Efficacy and Safety of JointStem, Autologous Adipose Tissue Derived Mesenchymal Stem Cells in Patients Diagnosed as Knee Osteoarthritis	140	Dec 2023
NCT02838069	A Phase IIb, Prospective, Multicentre, Double-blind, Triple-arm, Randomized Versus Placebo Trial, to Assess the Efficacy of a Single Injection of Either 2 or 10 x 10 ⁶ Autologous Adipose Derived Mesenchymal Stromal Cells (ASC) in the Treatment of Mild to Moderate Osteoarthritis (OA) of the Knee, Active and Unresponsive to Conservative Therapy for at Least 12 Months	153	Mar 2024
NCT04448106 ^a	Clinical Study for Subjects With Osteoarthritis of Knees, Hips, and Shoulders Using a Combination of Intravenous Infusions With Intra-articular Injection of Autologous Adipose Tissue-Derived Mesenchymal Stem Cells (AdMSCs)	300	Aug 2025
NCT04427930	Long-Term Safety and Efficacy Extension Study Of Autologous Adipose-Derived MesenchymalStem Cells (JOINTSTEM) in Patients With Knee Osteoarthritis: A Phase III Extension Study	260	Dec 2027

NCT No.	Trial Name	Planned Enrollment	Completion Date
NCT05288725	A Study to Evaluate the Safety, and Efficacy of Minimally Manipulated Autologous Bone Marrow Aspirate to Treat Knee Osteoarthritis in Patients	120	Dec 2024
NCT05517434	Intra-Articular Autologous Bone Marrow Aspirate Concentrate vs Placebo Injection and Lipoaspirate Concentrate With Leukocyte-Poor Platelet Rich Plasma vs Placebo Injection Evaluations for Treatment of Knee OsteoArthritis: The ABLE OA Double-Blinded Randomized Clinical Trial	148	Dec 2024
<i>Unpublished</i>			
NCT04310215 ^a	A Multi-center, Single-blind, Randomized, Phase III Clinical Trial to Evaluate the Efficacy and Safety of Adding CARTISTEM® on Microfracture in Patients With Talar Chondral or Osteochondral Defect	102	Nov 2022
NCT03818737	Randomized Multicenter Phase 3 Single-blind Trial Comparing the Efficacy of Corticosteroid Control to Mesenchymal Stem Cell Preparations From Autologous Bone Marrow Concentrate (BMAC), Adipose-derived Stem Cells in the Form of Stromal Vascular Fraction (SVF), and Third-party Human Mesenchymal Stem Cells Manufactured From Umbilical Cord Tissue for the Treatment of Unilateral Knee Osteoarthritis (OA)	480	Jul 2022
NCT04043819 ^a	Evaluation of Safety and Exploratory Efficacy of PSC-01, an Autologous Adipose-derived Stromal Vascular Fraction Cell Therapy Product for the Treatment of Knee Osteoarthritis	125	Jan 2021
NCT03990805 ^a	Multi-center, Randomized, Double-Blind, Placebo Controlled Phase 3 Clinical Trial to Evaluate Efficacy and Safety of Mesenchymal Stem Cells JointStem in Patients With Knee Osteoarthritis	260	Dec 2020

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

CODING

The following codes for treatment and procedures applicable to this policy are included below for informational purposes. This may not be a comprehensive list of procedure codes applicable to this policy.

Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

The code(s) listed below are medically necessary ONLY if the procedure is performed according to the "Policy" section of this document.

CPT/HCPCS	
20930	Allograft, morselized, or placement of osteopromotive material, for spine surgery only (List separately in addition to code for primary procedure). NOTE: This is a generic graft add on injection code to spine surgery which could be used for stem cells injection.
20999	Unlisted procedure, musculoskeletal system, general (Use for aspiration of bone marrow for the purpose of bone grafting, other than spine surgery and other therapeutic musculoskeletal applications)
38205	Blood-derived hematopoietic progenitor cell harvesting for transplantation, per collection; allogeneic
38206	Blood-derived hematopoietic progenitor cell harvesting for transplantation, per collection; autologous
38230	Bone marrow harvesting for transplantation; allogeneic
38232	Bone marrow harvesting for transplantation; autologous
38240	Hematopoietic progenitor cell (HPC); allogeneic transplantation per donor
38241	Hematopoietic progenitor cell (HPC); autologous transplantation
0263T	Intramuscular autologous bone marrow cell therapy, with preparation of harvested cells, multiple injections, one leg, including ultrasound guidance, if performed; complete procedure including unilateral or bilateral bone marrow harvest
0264T	Intramuscular autologous bone marrow cell therapy, with preparation of harvested cells, multiple injections, one leg, including ultrasound guidance, if performed; complete procedure excluding bone marrow harvest
0265T	Intramuscular autologous bone marrow cell therapy, with preparation of harvested cells, multiple injections, one leg, including ultrasound guidance, if performed; unilateral or bilateral bone marrow harvest only for intramuscular autologous bone marrow cell therapy
0489T	Autologous adipose-derived regenerative cell therapy for scleroderma in the hands; adipose tissue harvesting, isolation and preparation of harvested cells including incubation with cell dissociation enzymes, removal of non-viable cells and debris, determination of concentration and dilution of regenerative cells
0490T	Autologous adipose-derived regenerative cell therapy for scleroderma in the hands; multiple injections in one or both hands
0565T	Autologous cellular implant derived from adipose tissue for the treatment of osteoarthritis of the knees; tissue harvesting and cellular implant creation

CPT/HCPCS	
0566T	Autologous cellular implant derived from adipose tissue for the treatment of osteoarthritis of the knees; injection of cellular implant into knee joint including ultrasound guidance, unilateral
0717T	Autologous adipose-derived regenerative cell (ADRC) therapy for partial thickness rotator cuff tear; adipose tissue harvesting, isolation and preparation of harvested cells, including incubation with cell dissociation enzymes, filtration, washing and concentration of ADRCs
0718T	Autologous adipose-derived regenerative cell (ADRC) therapy for partial thickness rotator cuff tear; injection into supraspinatus tendon including ultrasound guidance, unilateral
C9359	Porous purified collagen matrix bone void filler (Integra Mozaik Osteoconductive Scaffold Putty, Integra Os Osteoconductive Scaffold Putty), per 0.5 cc
C9362	Porous purified collagen matrix bone void filler (Integra Mozaik Osteoconductive Scaffold Strip), per 0.5 cc

ICD 10-DIAGNOSES
Experimental / Investigational for all diagnoses related to this policy.

REVISIONS	
09-19-2013	Policy added to the bcbsks.com web site on 08-20-2013 for an effective date of 09-19-2013 for professional and institutional.
08-07-2015	Updated Description section.
	In Policy section: <ul style="list-style-type: none"> ▪ In Item B, removed "is" and added "are" to read, "Allograft bone productions containing viable stem cells including, but not limited to, demineralized bone matrix (DBM) with stem cells are considered experimental/investigational for all orthopedic applications. ▪ Added Item C, "Allograft or synthetic bone graft substitutes that must be combined with autologous blood or bone marrow are considered experimental/investigational for all orthopedic applications." ▪ In Policy Guidelines, removed "Note:", to read "This policy does not address unprocessed allograft bone."
	Updated Rationale section.
	Updated References section.
03-02-2016	Updated Description section.
	Updated Rationale section.
	Updated References section.
08-15-2017	Updated Description section.
	Updated Rationale section.
	Updated References section.
02-15-2018	Updated Description section.
	Updated Rationale section.
	In Coding section: <ul style="list-style-type: none"> ▪ Added CPT codes: 0263T, 0264T, 0265T. ▪ Removed CPT code: 38241.
	Updated References section.

REVISIONS	
02-18-2019	Updated Description section.
	Updated Rationale section.
	In Coding section: ▪ Added CPT code: 38232.
	Updated References section.
02-25-2021	Changed title from "Orthopedic Applications Of Stem Cell Therapy" to "Orthopedic Applications of Stem Cell Therapy (Including Allograft and Bone Substitute Products Used With Autologous Bone Marrow)"
	Updated Description section
	Updated Rationale section
	In coding section: • Added CPT codes: 20932, 20933, 20934, 20999, 38205, 38240, 38241, 0565T, 0566T, C9359, C9362
Updated Reference section	
04-11-2022	Updated Description Section
	Updated Rationale Section
	Updated Coding Section ▪ Added CPT codes: 20930, 0489T, 0490T ▪ Removed CPT codes: 20932, 20933, 20934
	Updated References Section
07-01-2022	Updated Coding Section • Added 0717T and 0718T
02-28-2023	Updated Description Section
	Updated Rationale Section
	Updated References Section

REFERENCES

1. Goldberg A, Mitchell K, Soans J, et al. The use of mesenchymal stem cells for cartilage repair and regeneration: a systematic review. *J Orthop Surg Res.* Mar 09 2017; 12(1): 39. PMID 28279182
2. U.S. Food & Drug Administration. Regulatory Considerations for Human Cells, Tissues, and Cellular and Tissue-Based Products: Minimal Manipulation and Homologous Use. <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/regulatory-considerations-human-cells-tissues-and-cellular-and-tissue-based-products-minimal>. Accessed December 5, 2022.
3. Borakati A, Mafi R, Mafi P, et al. A Systematic Review And Meta-Analysis of Clinical Trials of Mesenchymal Stem Cell Therapy for Cartilage Repair. *Curr Stem Cell Res Ther.* Feb 23 2018; 13(3): 215-225. PMID 28914207
4. Maheshwer B, Polce EM, Paul K, et al. Regenerative Potential of Mesenchymal Stem Cells for the Treatment of Knee Osteoarthritis and Chondral Defects: A Systematic Review and Meta-analysis. *Arthroscopy.* Jan 2021; 37(1): 362-378. PMID 32497658
5. Wiggers TG, Winters M, Van den Boom NA, et al. Autologous stem cell therapy in knee osteoarthritis: a systematic review of randomised controlled trials. *Br J Sports Med.* Oct 2021; 55(20): 1161-1169. PMID 34039582
6. Kim SH, Djaja YP, Park YB, et al. Intra-articular Injection of Culture-Expanded Mesenchymal Stem Cells Without Adjuvant Surgery in Knee Osteoarthritis: A Systematic Review and Meta-analysis. *Am J Sports Med.* Sep 2020; 48(11): 2839-2849. PMID 31874044

7. Jin L, Yang G, Men X, et al. Intra-articular Injection of Mesenchymal Stem Cells After High Tibial Osteotomy: A Systematic Review and Meta-analysis. *Orthop J Sports Med.* Nov 2022; 10(11): 23259671221133784. PMID 36452339
8. Wakitani S, Imoto K, Yamamoto T, et al. Human autologous culture expanded bone marrow mesenchymal cell transplantation for repair of cartilage defects in osteoarthritic knees. *Osteoarthritis Cartilage.* Mar 2002; 10(3): 199-206. PMID 11869080
9. Wakitani S, Nawata M, Tensho K, et al. Repair of articular cartilage defects in the patello-femoral joint with autologous bone marrow mesenchymal cell transplantation: three case reports involving nine defects in five knees. *J Tissue Eng Regen Med.* 2007; 1(1): 74-9. PMID 18038395
10. Wakitani S, Okabe T, Horibe S, et al. Safety of autologous bone marrow-derived mesenchymal stem cell transplantation for cartilage repair in 41 patients with 45 joints followed for up to 11 years and 5 months. *J Tissue Eng Regen Med.* Feb 2011; 5(2): 146-50. PMID 20603892
11. Centeno CJ, Schultz JR, Cheever M, et al. Safety and complications reporting on the re-implantation of culture-expanded mesenchymal stem cells using autologous platelet lysate technique. *Curr Stem Cell Res Ther.* Mar 2010; 5(1): 81-93. PMID 19951252
12. Wong KL, Lee KB, Tai BC, et al. Injectable cultured bone marrow-derived mesenchymal stem cells in varus knees with cartilage defects undergoing high tibial osteotomy: a prospective, randomized controlled clinical trial with 2 years' follow-up. *Arthroscopy.* Dec 2013; 29(12): 2020-8. PMID 24286801
13. Emadedin M, Labibzadeh N, Liastani MG, et al. Intra-articular implantation of autologous bone marrow-derived mesenchymal stromal cells to treat knee osteoarthritis: a randomized, triple-blind, placebo-controlled phase 1/2 clinical trial. *Cytotherapy.* Oct 2018; 20(10): 1238-1246. PMID 30318332
14. Lamo-Espinosa JM, Mora G, Blanco JF, et al. Intra-articular injection of two different doses of autologous bone marrow mesenchymal stem cells versus hyaluronic acid in the treatment of knee osteoarthritis: long-term follow up of a multicenter randomized controlled clinical trial (phase I/II). *J Transl Med.* Jul 31 2018; 16(1): 213. PMID 30064455
15. Lamo-Espinosa JM, Mora G, Blanco JF, et al. Intra-articular injection of two different doses of autologous bone marrow mesenchymal stem cells versus hyaluronic acid in the treatment of knee osteoarthritis: multicenter randomized controlled clinical trial (phase I/II). *J Transl Med.* Aug 26 2016; 14(1): 246. PMID 27565858
16. Vega A, Martín-Ferrero MA, Del Canto F, et al. Treatment of Knee Osteoarthritis With Allogeneic Bone Marrow Mesenchymal Stem Cells: A Randomized Controlled Trial. *Transplantation.* Aug 2015; 99(8): 1681-90. PMID 25822648
17. Shapiro SA, Kazmerchak SE, Heckman MG, et al. A Prospective, Single-Blind, Placebo-Controlled Trial of Bone Marrow Aspirate Concentrate for Knee Osteoarthritis. *Am J Sports Med.* Jan 2017; 45(1): 82-90. PMID 27566242
18. Koh YG, Kwon OR, Kim YS, et al. Comparative outcomes of open-wedge high tibial osteotomy with platelet-rich plasma alone or in combination with mesenchymal stem cell treatment: a prospective study. *Arthroscopy.* Nov 2014; 30(11): 1453-60. PMID 25108907
19. Zaffagnini S, Andriolo L, Boffa A, et al. Microfragmented Adipose Tissue Versus Platelet-Rich Plasma for the Treatment of Knee Osteoarthritis: A Prospective Randomized Controlled Trial at 2-Year Follow-up. *Am J Sports Med.* Sep 2022; 50(11): 2881-2892. PMID 35984721

20. Lim HC, Park YB, Ha CW, et al. Allogeneic Umbilical Cord Blood-Derived Mesenchymal Stem Cell Implantation Versus Microfracture for Large, Full-Thickness Cartilage Defects in Older Patients: A Multicenter Randomized Clinical Trial and Extended 5-Year Clinical Follow-up. *Orthop J Sports Med.* Jan 2021; 9(1): 2325967120973052. PMID 33490296
21. Whitehouse MR, Howells NR, Parry MC, et al. Repair of Torn Avascular Meniscal Cartilage Using Undifferentiated Autologous Mesenchymal Stem Cells: From In Vitro Optimization to a First-in-Human Study. *Stem Cells Transl Med.* Apr 2017; 6(4): 1237-1248. PMID 28186682
22. Vangsness CT, Farr J, Boyd J, et al. Adult human mesenchymal stem cells delivered via intra-articular injection to the knee following partial medial meniscectomy: a randomized, double-blind, controlled study. *J Bone Joint Surg Am.* Jan 15 2014; 96(2): 90-8. PMID 24430407
23. Vanichkachorn J, Peppers T, Bullard D, et al. A prospective clinical and radiographic 12-month outcome study of patients undergoing single-level anterior cervical discectomy and fusion for symptomatic cervical degenerative disc disease utilizing a novel viable allogeneic, cancellous, bone matrix (trinity evolution™) with a comparison to historical controls. *Eur Spine J.* Jul 2016; 25(7): 2233-8. PMID 26849141
24. Peppers TA, Bullard DE, Vanichkachorn JS, et al. Prospective clinical and radiographic evaluation of an allogeneic bone matrix containing stem cells (Trinity Evolution® Viable Cellular Bone Matrix) in patients undergoing two-level anterior cervical discectomy and fusion. *J Orthop Surg Res.* Apr 26 2017; 12(1): 67. PMID 28446192
25. Jones CP, Loveland J, Atkinson BL, et al. Prospective, Multicenter Evaluation of Allogeneic Bone Matrix Containing Viable Osteogenic Cells in Foot and/or Ankle Arthrodesis. *Foot Ankle Int.* Oct 2015; 36(10): 1129-37. PMID 25976919
26. Eastlack RK, Garfin SR, Brown CR, et al. Osteocel Plus cellular allograft in anterior cervical discectomy and fusion: evaluation of clinical and radiographic outcomes from a prospective multicenter study. *Spine (Phila Pa 1976).* Oct 15 2014; 39(22): E1331-7. PMID 25188591
27. Sen RK, Tripathy SK, Aggarwal S, et al. Early results of core decompression and autologous bone marrow mononuclear cells instillation in femoral head osteonecrosis: a randomized control study. *J Arthroplasty.* May 2012; 27(5): 679-86. PMID 22000577
28. Zhao D, Cui D, Wang B, et al. Treatment of early stage osteonecrosis of the femoral head with autologous implantation of bone marrow-derived and cultured mesenchymal stem cells. *Bone.* Jan 2012; 50(1): 325-30. PMID 22094904
29. American Academy of Orthopaedic Surgeons. Management of Glenohumeral Joint Osteoarthritis Evidence-Based Clinical Practice Guideline. <https://www.aaos.org/globalassets/quality-and-practice-resources/glenohumeral/gjo-cpg.pdf>. Updated March 23, 2020. Accessed December 5, 2022.
30. American Academy of Orthopaedic Surgeons. Management of Osteoarthritis of the Knee (Non-Arthroplasty). <https://www.aaos.org/globalassets/quality-and-practice-resources/osteoarthritis-of-the-knee/oak3cpg.pdf>. Updated August 31, 2021. Accessed December 4, 2022.
31. Kaiser MG, Groff MW, Watters WC, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 16: bone graft extenders and substitutes as an adjunct for lumbar fusion. *J Neurosurg Spine.* Jul 2014; 21(1): 106-32. PMID 24980593
32. Kolasinski SL, Neogi T, Hochberg MC, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Management of Osteoarthritis of the

Hand, Hip, and Knee. Arthritis Care Res (Hoboken). Feb 2020; 72(2): 149-162. PMID 31908149

OTHER REFERENCES

1. Blue Cross and Blue Shield of Kansas Orthopedic Liaison Committee, June 2016; August 2017.