

Medical Policy



Title: Percutaneous Vertebroplasty and Sacroplasty

Professional

Original Effective Date: October 18, 2004
 Revision Date(s): April 21, 2005;
 September 7, 2005; December 14, 2005;
 February 21, 2006; May 9, 2006;
 July 27, 2006; September 14, 2006;
 October 31, 2006; January 1, 2007
 July 23, 2009; January 1, 2012;
 October 4, 2013; December 31, 2013;
 January 1, 2015; October 21, 2015;
 January 1, 2016; January 19, 2018;
 May 23, 2018; August 1, 2018; May 21, 2019;
 May 23, 2021
 Current Effective Date: May 23, 2021

Institutional

Original Effective Date: July 1, 2005
 Revision Date(s): September 7, 2005;
 December 14, 2005; February 21, 2006;
 May 9, 2006; July 27, 2006;
 September 14, 2006; October 31, 2006;
 January 1, 2007; July 23, 2009;
 January 1, 2012; October 4, 2013;
 December 31, 2013; January 1, 2015;
 October 21, 2015; January 1, 2016;
 January 19, 2018; May 23, 2018;
 August 1, 2018; May 21, 2019; May 23, 2021
 Current Effective Date: May 23, 2021

State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).

The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.

The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.

If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"> With symptomatic osteoporotic vertebral fractures between 6 weeks and 1 year old 	Interventions of interest are: <ul style="list-style-type: none"> Vertebroplasty 	Comparators of interest are: <ul style="list-style-type: none"> Conservative management 	Relevant outcomes include: <ul style="list-style-type: none"> Symptoms Functional outcomes Quality of life Hospitalizations Medication use Treatment-related morbidity

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"> • With symptomatic osteoporotic vertebral fractures less than 6 weeks old 	Interventions of interest are: <ul style="list-style-type: none"> • Vertebroplasty 	Comparators of interest are: <ul style="list-style-type: none"> • Conservative management 	Relevant outcomes include: <ul style="list-style-type: none"> • Symptoms • Functional outcomes • Quality of life • Hospitalizations • Medication use • Treatment-related morbidity
Individuals: <ul style="list-style-type: none"> • With sacral insufficiency fractures 	Interventions of interest are: <ul style="list-style-type: none"> • Sacroplasty 	Comparators of interest are: <ul style="list-style-type: none"> • Conservative management 	Relevant outcomes include: <ul style="list-style-type: none"> • Symptoms • Functional outcomes • Quality of life • Hospitalizations • Medication use • Treatment-related morbidity

DESCRIPTION

Percutaneous vertebroplasty is an interventional technique involving the fluoroscopically guided injection of polymethyl methacrylate (PMMA) into a weakened vertebral body. The technique has been investigated to provide mechanical support and symptomatic relief in patients with osteoporotic vertebral compression fractures or in those with osteolytic lesions of the spine (e.g., multiple myeloma, metastatic malignancies); as a treatment for sacral insufficiency fractures; and as a technique to limit blood loss related to surgery.

Objective

The objective of this evidence review is to evaluate whether vertebroplasty or sacroplasty improves the net health outcome in individuals with osteoporotic vertebral compression fractures or sacral insufficiency fractures.

Background

Treatment of Vertebral Compression Fracture

Chronic symptoms do not tend to respond to the management strategies for acute pain such as bed rest, immobilization or bracing device, and analgesic medication, sometimes including narcotic analgesics. The source of chronic pain after vertebral compression fracture may not be from the vertebra itself but may be predominantly related to strain on muscles and ligaments secondary to kyphosis. This type of pain frequently does not improve with analgesics and may be better addressed through exercise or physical therapy. Improvements in pain and ability to function are the principal outcomes of interest for treatment of osteoporotic fractures.

Treatment of Sacral Insufficiency Fractures

Similar interventions are used for sacral and vertebral fractures and include bed rest, bracing, and analgesics. Initial clinical improvements may occur quickly; however, resolution of all symptoms may not occur for 9 to 12 months.^{1,2}

Vertebral and Sacral Body Metastasis

Metastatic malignant disease of the spine generally involves the vertebrae/sacrum, with pain being the most frequent complaint.

Treatment of Vertebral and Sacral Body Metastasis

While radiotherapy and chemotherapy are frequently effective in reducing tumor burden and associated symptoms, pain relief may be delayed days to weeks, depending on tumor response.

Further, these approaches rely on bone remodeling to regain strength in the vertebrae/sacrum, which may necessitate supportive bracing to minimize the risk of vertebral/sacral collapse during healing. Improvements in pain and function are the primary outcomes of interest for treatment of bone malignancy with percutaneous vertebroplasty or sacroplasty.

SURGICAL TREATMENT OPTIONS

Percutaneous Vertebroplasty

Vertebroplasty is a surgical procedure that involves the injection of synthetic cement (e.g., polymethylmethacrylate, bis-glycidal dimethacrylate [Cortoss]³) into a fractured vertebra. It has been suggested that vertebroplasty may provide an analgesic effect through mechanical stabilization of a fractured or otherwise weakened vertebral body. However, other mechanisms of effect have been postulated, including thermal damage to intraosseous nerve fibers.

Percutaneous Sacroplasty

Sacroplasty evolved from the treatment of insufficiency fractures in the thoracic and lumbar vertebrae with vertebroplasty. The procedure, essentially identical to vertebroplasty, entails guided injection of polymethylmethacrylate through a needle inserted into the fracture zone. Although first described in 2000 as a treatment for symptomatic sacral metastatic lesions,^{4,5} it is most often described as a minimally invasive alternative to conservative management^{6,7,8} for sacral insufficiency fractures.

Pain and function are subjective outcomes and, thus, may be susceptible to placebo effects. Furthermore, the natural history of pain and disability associated with these conditions may vary. Therefore, controlled comparison studies would be valuable to demonstrate the clinical effectiveness of vertebroplasty and sacroplasty over and above any associated nonspecific or placebo effects and to demonstrate the effect of treatment compared with alternatives such as continued medical management.

In all clinical situations, adverse events related to complications from vertebroplasty and sacroplasty are the primary harms to be considered. Principal safety concerns relate to the incidence and consequences of leakage of the injected polymethyl methacrylate or another injectate.

REGULATORY STATUS

Vertebroplasty is a surgical procedure and, as such, is not subject to U.S. Food and Drug Administration (FDA) approval.

Polymethylmethacrylate bone cement was available as a drug product before enactment of the FDA's device regulation and was at first considered what the FDA terms a "transitional device." It was transitioned to a class III device requiring premarketing applications. Several orthopedic companies have received approval of their bone cement products since 1976. In 1999, polymethylmethacrylate was reclassified from class III to class II, which requires future 510(k) submissions to meet "special controls" instead of "general controls" to assure safety and effectiveness. Thus, use of polymethylmethacrylate in vertebroplasty represented an off-label use of an FDA-regulated product before 2005. In 2005, polymethylmethacrylate bone cements such as Spine-Fix® Biomimetic Bone Cement and Osteopal® V were cleared for marketing by the FDA through the 510(k) process for the fixation of pathologic fractures of the vertebral body using vertebroplasty procedures.

The use of polymethylmethacrylate in sacroplasty is an off-label use of an FDA-regulated product (bone cements such as Spine-Fix® Biomimetic Bone Cement [Teknimed] and Osteopal® V [Heraeus]) because the 510(k) approval was for the fixation of pathologic fractures of the vertebral body using vertebroplasty procedures. Sacroplasty was not included. FDA product code: NDN. In 2009, Cortoss® (Stryker) Bone Augmentation Material was cleared for marketing by the FDA through the 510(k) process. Cortoss® is a nonresorbable synthetic material that is a composite resin-based, bis-glycidyl dimethacrylate. The FDA classifies this product as a polymethylmethacrylate bone cement.

In 2010, the Parallax® Contour® Vertebral Augmentation Device (ArthroCare) was cleared for marketing by FDA through the 510(k) process. The device creates a void in cancellous bone that can then be filled with bone cement. FDA product code: HXG.

POLICY

- A. Percutaneous vertebroplasty may be considered **medically necessary** for:
1. The treatment of severe pain due to osteolytic lesions of the spine related to multiple myeloma or metastatic malignancies; **OR**
 2. The treatment of symptomatic osteoporotic vertebral fractures that have failed to respond to conservative treatment (e.g., rest, analgesics, physical therapy) for at least 6 weeks; **OR**
 3. The treatment of symptomatic osteoporotic vertebral fractures that are less than 6 weeks in duration that have led to hospitalization or persist at a level that prevents ambulation.
- B. Percutaneous vertebroplasty is considered **experimental / investigational** for all other indications, including use in acute vertebral fractures due to osteoporosis or trauma.
- C. Percutaneous sacroplasty is considered **experimental / investigational** for all indications, including use in sacral insufficiency fractures due to osteoporosis and spinal lesions due to metastatic malignancies or multiple myeloma.

RATIONALE

This evidence review has been updated regularly with searches of the PubMed database. The most recent literature update was performed through February 11, 2020.

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function, including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

PERCUTANEOUS VERTEBROPLASTY FOR VERTEBRAL COMPRESSION FRACTURES OF BETWEEN 6 WEEKS AND 1 YEAR OLD

Clinical Context and Therapy Purpose

Osteoporotic compression fractures are common. It is estimated that up to one-half of women and approximately one-quarter of men will have a vertebral fracture at some point in their lives. However, only about one-third of vertebral fractures reach clinical diagnosis, and most symptomatic fractures will heal within a few weeks or one month with medical management. Nonetheless, some individuals with acute fractures will have severe pain and decreased function that interferes with the ability to ambulate and is not responsive to usual medical management. Also, a minority of patients will exhibit chronic pain following osteoporotic compression fracture that presents challenges for medical management.

The purpose of vertebroplasty is to provide a treatment option that is an alternative to or an improvement on existing therapies in patients with symptomatic osteoporotic or osteolytic vertebral fractures between 6 weeks and 1 year old.

The question addressed in this evidence review is: Does vertebroplasty improve the net health outcome in individuals with symptomatic osteoporotic or osteolytic vertebral fractures between 6 weeks and 1 year old?

The following PICO was used to select literature to inform this review.

Patients

The relevant population of interest is individuals with symptomatic osteoporotic or osteolytic vertebral fractures between 6 weeks and 1 year old. With acute fractures, these individuals experience severe pain, decreased ambulatory function, and a lessened response to conservative medical management. Risk factors for osteoporotic or osteolytic vertebral fractures can include osteopenia, osteoporosis, advanced age, inactivity, corticosteroid use, female sex, and depression.

Interventions

The therapy being considered is vertebroplasty, a procedure for stabilizing compression fractures in the spine, during which bone cement is injected into the fractured vertebra through a small hole in the skin in order to relieve back pain. The vertebroplasty procedure is performed in an outpatient setting by interventional radiologists or orthopedic surgeons.

Comparators

Comparators of interest include conservative management. Conservative management includes measures to reduce pain and improve mobility. Physical therapy, analgesics, narcotics, and hormone treatments can be prescribed to achieve this. Bed rest and braces may also be utilized as conservative management; however, these modalities are associated with prolonged immobilization which can further exacerbate bone loss and fail to relieve systems. Patients who receive conservative management are typically managed by pediatricians, physical therapists, and primary care providers in an outpatient clinical setting.

Outcomes

The general outcomes of interest are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. Negative outcomes can include complications with sedation, further injury during transfer to the radiology table, and the possibility

of abuse after the prescription of narcotics. The outcomes of interest for vertebroplasty as a treatment for symptomatic vertebral fractures have varying follow-up times to fully examine the impact on the patient, ranging from shorter term outcomes like medication use to outcomes that require extended follow-up, such as functional outcomes. Given that the existing literature evaluating vertebroplasty as a treatment for symptomatic vertebral fractures between 6 weeks and 1 year old has varying lengths of follow-up, ranging from 6 months to 2 years, follow-up timing of 1 year is appropriate to demonstrate efficacy.

Disability, a major factor on quality of life, is measured using various tools throughout the literature. Three such tools include the Roland-Morris Disability Questionnaire,⁹ the visual analogue scale,¹⁰ and QUALEFFO (a quality of life questionnaire in patients with vertebral fractures). The Roland-Morris Disability Questionnaire is a self-administered disability measure in which greater levels of disability are reflected by higher numbers on a 24-point scale and on visual analogue scale. The Roland-Morris Disability Questionnaire has been shown to yield reliable measurements, which are valid for inferring the level of disability, and to be sensitive to change over time for groups of patients with low back pain. Visual analogue scale is commonly used as the outcome measure for such studies. It is usually presented as a 100-mm horizontal line on which the patient's pain intensity is represented by a point between the extremes of "no pain at all" and "worst pain imaginable." With QUALEFFO (a quality of life questionnaire in patients with vertebral fractures), quality of life is measured by the scale 0 to 100, higher scores indicating worse quality of life.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

1. To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
2. In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
3. To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
4. Studies with duplicative or overlapping populations were excluded.

This evidence review was informed by a TEC Assessment (2000), which was updated periodically through 2010.^{11,12,13,14,15,16} Subsequent evidence includes a number of RCTs, 2 of which included a sham control, and numerous RCTs that compared vertebroplasty with conservative management.

Review of Evidence Systematic Reviews

Buchbinder et al (2018) published a Cochrane review of the literature up to November 2014¹⁷. Studies compared vertebroplasty versus placebo (2 studies with 209 randomized participants), usual care (6 studies with 566 randomized participants), and kyphoplasty (4 studies with 545 randomized participants). The majority of participants were female, between 63.3 and 80 years of age, with symptom duration ranging from 1 week to more than 6 months. At 1 month, disease-specific quality of life measured by the QUALEFFO (a quality of life questionnaire in patients with vertebral fractures; scale 0 to 100, higher scores indicating worse quality of life) was 0.40 points worse in the vertebroplasty group. Based upon moderate quality evidence from 3 trials (1 placebo, 2 usual care, 281 participants) with up to 12 months follow-up, it is unclear if vertebroplasty increases the risk of new symptomatic vertebral fractures. Similarly, based upon moderate quality evidence from 2 placebo-controlled trials, it is unclear to what extent risk of other adverse events

exists. There were 3/106 adverse events observed in the vertebroplasty group compared with 3/103 in the placebo group; risk ratio, 1.01 (95% confidence interval [CI]: 0.21 to 4.85). Serious adverse events that have been reported with vertebroplasty included osteomyelitis, cord compression, thecal sac injury, and respiratory failure.

Staples et al (2011) conducted a patient-level meta-analysis of the 2 sham-controlled trials to determine whether vertebroplasty is more effective than sham in specific subsets of patients.¹⁸ This subset analysis focused on duration of pain (≤ 6 weeks vs. > 6 weeks) and severity of pain (score < 8 or ≥ 8 on an 11-point numeric rating scale). The analysis included 209 participants (78 from the Australian trial, 131 from the U.S. trial); 27% had pain of recent onset and 47% had severe pain at baseline. The primary outcome measures (pain scores and function on the Roland-Morris Disability Questionnaire at 1 month) did not differ significantly between groups. Responder analyses were also conducted based on a 3-unit improvement in pain scores, a 3-unit improvement in Roland-Morris Disability Questionnaire scores, and a 30% improvement in each of the pain and disability outcomes. The only difference observed between groups was a trend in the vertebroplasty group to achieve at least 30% improvement in pain scores (relative risk, 1.32; 95%CI, 0.98 to 1.76; $p=0.07$), a result that may have been confounded by the greater use of opioid medications in that group.

Xie et al (2017), in a meta-analysis of RCTs, evaluated the efficacy and safety in percutaneous vertebroplasty and conservative treatment for patients with osteoporotic vertebral compression fractures.¹⁹ Thirteen studies were selected ($n=1231$ patients; 623 to vertebroplasty, 608 to conservative treatment). Outcomes included pain relief (from 1 week to 6 months), quality of life assessments, and the rate of adjacent-level vertebral fracture. Vertebroplasty was superior for pain relief at 1 week and at 1 month. It was inferior to conservative treatment for pain relief at 6 months. Vertebroplasty showed improvement over conservative treatment for quality of life, as measured using the Quality of Life Questionnaire of the European Foundation for Osteoporosis. No statistically significant differences were found between treatments for the rate of adjacent-level vertebral fractures. Limitations included the inclusion of several studies with inadequate blinding and heterogenous reporting of patient characteristics outcomes.

Table 1. Characteristics of Systematic Reviews and Meta-Analyses on Percutaneous Vertebroplasty for Vertebral Compression Fractures of Between 6 Weeks and 1 Year Old

Study	Dates	Trials	Participants	Intervention	N (Range)	Design
Buchbinder et al (2018) ¹⁷ ,	2007-2016	21	Patients with osteoporotic vertebral fractures (mean age ranged from 63.3 to 80 years); symptom duration ranged from 1 week to > 6 months.	Vertebroplasty	2862 (46-404)	RCT
Staples et al (2011) ¹⁸ ,	NR	2	Participants with 1-2 painful osteoporotic vertebral fractures > 12 months duration and unhealed, as confirmed by MRI, were randomly assigned	Vertebroplasty vs. placebo (5 studies); kyphoplasty (7 studies); facet joint steroid injection (1)	209 (78-131)	RCT

Study	Dates	Trials	Participants	Intervention	N (Range)	Design
			to vertebroplasty or to a sham procedure.			
Xie et al (2017) ¹⁹ ,	NR-2017	13	Patients with OVCFs	PVP vs. conservative treatment	2561 (NR)	RCT

NR: not reported; OVCF: osteoporotic vertebral compression fracture; PVP: percutaneous vertebroplasty; RCT: randomized controlled trial

Table 2. Results of Systematic Reviews and Meta-Analyses on Percutaneous Vertebroplasty for Vertebral Compression Fractures of Between 6 Weeks and 1 Year Old

Study	Quality of Life	New Fractures
	QUALEFFO	
Buchbinder et al (2018) ¹⁷ ,		
Placebo group at 1-month, score (N)	4.58 (71)	NR
Vertebroplasty group at 1-month, score (N)	5.38 (71)	NR
Absolute change between groups	0.4% worse (5% worse-5% better [n=71])	NR
Relative change between groups	0.7% worse (9% worse-8% better [n=71])	NR
N. Intervention group (%)	NR	28 (19.58)
N. Placebo group (%)	NR	19 (50.00)
RR (CI)	NR	1.47 (0.39 to 5.50)
	Duration of Pain	
Staples et al (2011) ¹⁸ ,		
Mean change score (SD) of pain, at 2 weeks, PVP vs placebo	2.2 (2.8) vs. 2.5 (3.0)	NR
Adjusted between group difference (CI) at 2 weeks	~0.2 (~0.9-0.6)	
Mean change score (SD) of pain, at 1 month, PVP vs placebo	2.08 (3.0) vs. 2.2 (3.2)	NR
Adjusted between group difference (CI) at 2 weeks	0.6 (~0.2-1.4)	
	Pain relief	
Xie et al (2017) ¹⁹ ,	N=1231	NR
At 1-week (vertebroplasty superior), MD (CI)	1.36 (0.55 to 2.17)	NR
At 1-month (vertebroplasty superior), MD (CI)	1.56 (0.43 to 2.70)	NR
At 6-months (vertebroplasty inferior), MD (CI)	-1.59 (-2.9 to -0.27) p<0.05	NR
Total (vertebroplasty superior), MD (CI)	-5.03 (7.94 to -2.12)	NR

CI: 95% confidence interval; MD: mean difference; ; (score <8 or ≥8 on an 11-point numeric rating scale); NR: not reported; PVP: percutaneous vertebroplasty; QUALEFFO-41 Questionnaire: a quality of life questionnaire in patients with vertebral fractures; RR: relative risk; SD: standard deviation.

Randomized Controlled Trials

Vertebroplasty Versus Medical Management With Sham Controls

Three sham-controlled trials compared vertebroplasty with medical management using a sham control (that included local anesthetic), which mimicked the vertebroplasty procedure up to the point of cement injection.^{20,21} Buchbinder et al (2009) reported on results for a 4-center, randomized, double-blind, sham-controlled trial with 78 patients with 1 or 2 painful osteoporotic vertebral fractures with a duration of less than 1 year.²⁰ Patients were assigned to vertebroplasty or sham procedure (i.e., injection of local anesthetic into the facet capsule and/or periosteum). Ninety-one percent of participants completed 6 months of follow-up. The participants, investigators (other than the radiologists performing the procedure), and outcome assessors were blinded to the treatment assignment. Kroon et al (2014) reported results of the same trial at 12 and 24 months, maintaining blinding throughout the follow-up period.²² The primary outcome was overall pain measured on a visual analogue scale from 0 to 10, with 1.5 points representing the minimal clinically important difference. For the primary outcome, reviewers reported no significant differences in visual analogue scale pain score at 3, 12, or 24 months. With reductions in pain and improvements in quality of life observed in both groups, the authors concluded routine use of vertebroplasty provided no benefit.

Kallmes et al (2009) conducted a multicenter, randomized, double-blind, sham-controlled, Investigational Vertebroplasty Safety and Efficacy Trial in which 131 participants with 1 to 3 painful osteoporotic vertebral fractures were assigned to vertebroplasty or sham procedure (injection of local anesthetic into the facet capsule and/or periosteum).²¹ Participants had back pain for no more than 12 months and had a current pain rating of at least 3 on visual analogue scale at baseline. Participants were evaluated at various time points to 1 year postprocedure. Ninety-seven percent completed a 1-month follow-up; 95% completed 3 months. The primary outcomes were Roland-Morris Disability Questionnaire scores and average back pain intensity during the preceding 24 hours at 1 month, with a reduction of 30% in Roland-Morris Disability Questionnaire and visual analogue scale pain scores considered a clinically meaningful difference.²³

For the primary endpoints at one month, there were no significant between-group differences. There was a trend toward a higher clinically meaningful improvement in pain at 1 month (30% reduction from baseline) in the vertebroplasty group (64% vs. 48%, respectively; $p=0.06$). At 3 months, 51% from the control group and 13% in the vertebroplasty group crossed over ($p<0.001$). Comstock et al (2013) reported on patient outcomes at one year, at which point 16% of patients who underwent vertebroplasty and 60% of control subjects had crossed over to the alternative procedure ($p<0.001$).²⁴ The as-treated analysis found no significant difference in Roland-Morris Disability Questionnaire or pain scores between the 2 groups. Intention-to-treat analysis found a modest 1-point difference in pain rating and no significant difference in Roland-Morris Disability Questionnaire score. There was a significant difference in the percentage of patients showing a 30% or greater improvement in pain (70% of patients randomized to vertebroplasty vs. 45% of patients randomized to the control group). One limitation of this study is that at 14 days, 63% of patients in the control group correctly guessed they had the control intervention, and 51% of patients in the vertebroplasty group correctly guessed they had the vertebroplasty.

Firanesu et al (2018) published the results of a randomized, double-blind, sham-controlled clinical trial performed in 4 community hospitals in the Netherlands from 2011 to 2015.²⁵ The main outcome measured was mean reduction in visual analogue scale scores at 1 day, 1 week, and 1, 3, 6, and 12 months. The mean reduction in visual analogue scale score was statistically significant in the vertebroplasty and sham procedure groups at all follow-up points after the procedure compared with baseline. These changes in visual analogue scale scores did not, however, differ statistically significantly between the groups during 12 months' follow-up.

Table 3. Summary of Characteristics of Key RCT Comparing Vertebroplasty Versus Medical Management With Sham Controls

Study	Countries	Sites	Dates	Participants (N)	Interventions	
					Active (n)	Comparator (n)
Buchbinder et al (2009) ²⁰ ,	US	4	2003-2008	Patients with 1-2 painful OCVF, duration <1 year	Vertebroplasty (38)	sham procedure (40) ¹
Kallmes et al (2009) ²¹ ,	US, UK, Aus	10	2004-2008	Participants with 1-3 painful OCVF, pain < 12 mo, current pain VAS > 3	Vertebroplasty (68)	sham procedure ¹ (63)
Firanesu et al (2018) ²⁵ ,	Netherlands	4	2011-2015	Participants with acute OCVF	Vertebroplasty (91)	sham procedure ¹ (89)

OCVF: osteoporotic vertebral compression fracture; RCT: randomized controlled trial; VAS: visual analogue scale.

¹ Injection of local anesthetic into the facet capsule and/or periosteum.

Table 4. Summary of Results of Key RCT Comparing Vertebroplasty Versus Medical Management with Sham Controls

Study	VAS	RMDQ
Buchbinder et al (2009) ²⁰ ,	N=73, at 3-months	
Intervention (mean±SD)	Reduction: 2.6±2.9	
Control (mean±SD)	Reduction: 1.9±3.3	
Adjusted between-group difference (CI)	0.6 (-0.7-1.8)	
Kallmes et al (2009) ²¹ ,		
Day 14 Mean difference between groups (CI)	0.1 (-0.8-1.1)	-0.6 (-2.4-1.2)
P-value	0.77	0.35
Month 1 Mean difference between groups (CI)	0.7 (-0.3-1.70)	0.7 (-1.3-2.8)
P-value	0.19	0.49
Firanesu et al (2018) ²⁵ ,	N=180	
Day 1 Mean difference between groups (CI)	-0.43 (-1.17 - 0.31)	
Week 1 Mean difference between groups (CI)	-0.11 (-0.85 - 0.63)	

Study	VAS	RMDQ
Month 1 Mean difference between groups (CI)	0.41 (-0.33 - 1.15)	
Month 3 Mean difference between groups (CI)	0.21 (-0.54 - 0.96)	
Month 6 Mean difference between groups (CI)	0.39 (-0.33 - 1.15)	
Month 12 Mean difference between groups (CI)	0.45 (-0.37-1.24)	

CI: 95% confidence interval; NR: not reported; RCT: randomized controlled trial; RMDQ: Roland-Morris Disability Questionnaire; SD: standard deviation; VAS: visual analogue score.

Table 5. Relevance Study Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-up ^e
Buchbinder et al (2009) ^{20,}					
Kallmes et al (2009) ^{21,}				3. No reporting of harms. 5. investigator modified pain window from 6 to 9 weeks.	
Firanesu et al (2018) ^{25,}	2. Lack of screening for co-occurring pain conditions. 2. MRI was not conducted.			5. investigator modified pain window from 6 to 9 weeks.	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not established and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 6. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Follow-Up ^d	Power ^e	Statistical ^f
Buchbinder et al (2009) ^{20,}			2. 30% of eligible participants declined to participate, selection bias cannot be ruled out.>			

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Follow-Up ^d	Power ^e	Statistical ^f
Kallmes et al (2009) ²¹ ,		1. At 14 days, > 50% of participants in either arm correctly identified their intervention assignment.		4. Due to high crossover the group differences in outcomes were complicated.		
Firanesco et al (2018) ²⁵ ,	4. Screening logs not retained.					

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Follow-Up key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Intervention is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Intervention is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Vertebroplasty Versus Medical Management Without Sham Controls

Chen et al (2014) reported on a nonblinded RCT comparing vertebroplasty with conservative management.²⁶ The trial included 89 patients with chronic compression fractures confirmed by magnetic resonance imaging and persistent severe pain for 3 months or longer. The evaluation was performed at 1 week and 1, 3, 6, and 12 months. Over the course of 1 year, pain scores decreased from 6.5 to 2.5 in the vertebroplasty group and from 6.4 to 4.1 in the control group (p<0.001). Complete pain relief was reported by 84.8% of patients in the vertebroplasty group and 34.9% of controls. The final Oswestry Disability Index score was 15.0 in the vertebroplasty group and 32.1 in the conservative management group (p<0.001), and the final Roland-Morris Disability Questionnaire score was 8.1 for vertebroplasty and 10.7 for controls (p<0.001).

Farrokhi et al (2011) reported on a blinded RCT that compared vertebroplasty with optimal medical management in 82 patients.²⁷ Patients had painful osteoporotic vertebral compression fractures that were refractory to analgesic therapy for at least 4 weeks and less than 1 year. Control of pain and improvement in quality of life were measured by independent raters before treatment and at 1 week and 2, 6, 12, 24, and 36 months after treatment began. Radiologic evaluation to measure vertebral body height and correction of deformity was performed before and after treatment and after 36 months of follow-up. Adverse events include new symptomatic adjacent fractures in 1 patient in the treatment group and 6 in the control group. Additionally, 1 patient experienced epidural cement leakage, which caused severe lower extremity pain and weakness, and had to be treated with bilateral laminectomy and evacuation of the bone cement.

Table 7. Summary of Key RCT Characteristics - Vertebroplasty Versus Medical Management Without Sham Controls

Study	Countries	Sites	Dates	Participants (N)	Interventions	
					Active	Comparator
Chen et al (2014) ²⁶ ,	China	1	2007-2012	Patients with chronic compression fractures confirmed by MRI and persistent severe pain for <3 months (89)	Vertebroplasty	Conservative Management
Farrokhi et al (2011) ²⁷ ,	Iran	1	2004-2005	Patients with painful osteoporotic vertebral compression fractures refractory to analgesic therapy for >4, but <1 year (82)	Vertebroplasty	Optimal Medical Management

RCT: randomized controlled trial; MRI: magnetic resonance imaging.

Table 8. Summary of Key RCT Results

Study	Pain Score	ODI score	RMDQ
	Overall pain (scale 0-10)		
Chen et al (2014) (N=89) ²⁶ ,			
Intervention Group, Pooled at 1-year	2.5	15.0	8.1
Control Group, Pooled at 1-year	4.1	32.1	10.7
P-value	P<0.001	P<0.001	P<0.001
Farrokhi et al (2011) ²⁷ ,	VAS Score		
Week 1 Mean difference between groups (CI)p-value	-3.1 (-3.72 to -2.28)<0.001	-14.0 (-15.00 to -12.82)<0.028	
Month 2 Mean difference between groups (CI)p-value	-2.9 (-4.9 to -0.82)<0.011	-15.0 (-16.76 to -13.24)<0.019	
Month 6 Mean difference between groups (CI)p-value	-1.9 (-3.25 to -0.55)<0.021	-11.0 (-12.17 to -7.83)<0.011	
Month 12 Mean difference between groups (CI)p-value	-1.9 (-2.9 to 0.9)<0.11	-12.0 (-13.5 to -11.5)<0.021	

CI: confidence interval;; NNT: number needed to treat; NR: not reported; ODI: Oswestry Disability Index, RCT: randomized controlled trial; RMDQ: Roland-Morris Disability Questionnaire; VAS: visual analogue scale.

Table 9. Study Relevance Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
Chen et al (2014) ²⁶ ,			3. Investigator modified duration of the conservative therapy from 6 to 4 weeks		
Farrokhi et al (2011) ²⁷ ,				4. Language translation of Oswestry scale not validated.	

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 10. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Follow-Up ^d	Power ^e	Statistical ^f
Chen et al (2014) ²⁶ ,		1,2. This study was not blinded.	.			
Farrokhi et al (2011) ²⁷ ,						

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Follow-Up key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Intervention is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Intervention is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Nonrandomized Comparative Studies

Edidin et al (2011, 2015) reported on mortality risk rates in Medicare patients who had vertebral compression fractures and were treated with vertebroplasty, kyphoplasty, or nonoperatively.^{28,29} These studies were industry funded. In the 2015 report, they identified 1,038,956 patients who had vertebral compression fractures between 2005 and 2009. The dataset included 141,343 kyphoplasty patients and 75,364 vertebroplasty patients. The matched cohort included 100,649 nonoperated patients, 36,657 kyphoplasty patients, and 24,313 vertebroplasty patients. Survival was calculated from the index diagnosis date until death or the end of follow-up (up to 4 years). Analysis of the whole data set before matching indicated that patients in the nonoperated cohort had a 55% (95% CI, 53% to 56%; $p < 0.001$) higher risk of mortality than the kyphoplasty cohort and a 25% (95% CI, 23% to 26%; $p < 0.001$) higher mortality risk than the vertebroplasty cohort. After propensity matching, the risk of mortality at 4 years was 47.2% in the nonoperated group compared with 42.3% in the kyphoplasty group ($p < 0.001$) and 46.2% in the vertebroplasty group ($p < 0.001$).

Lin et al (2017) reported on mortality risk in elderly patients (>70 years old) who had vertebral compression fractures and were treated with early vertebroplasty (within 3 months) or conservative therapy.³⁰ The data set consisted of 10,785 Taiwanese patients who were selected through the National Health Insurance Research Database, of whom 1,773 patients received vertebroplasty, and 5,324 did not; a minority of these patients had osteoarthritis. The authors found that a "significant difference in survival curves of mortality and respiratory failure" existed between both groups of patients ($p < 0.05$). The incidence of death at 1 year in the vertebroplasty group was 0.46 per 100 person-months (95% CI, 0.38 to 0.56). The incidence of death at 1 year in the nonvertebroplasty group was 0.63 per 100 person-months (95% CI, 0.57 to 0.70). With regard to respiratory failure, hazard ratio between groups was 1.46 (95% CI, 1.04 to 2.05; $p = 0.028$). Limitations of this study included the broad selection of the population, which was not restricted only to patients with osteoporotic lesions. Also, authors were limited by the database, which did not report on pain or functional outcomes.

Section Summary: Percutaneous Vertebroplasty for Vertebral Compression Fractures of Between 6 Weeks and 1 Year Old

Despite evidence from numerous RCTs, including 2 with sham controls, the efficacy of vertebroplasty for painful osteoporotic compression fractures of less than 1 year remains uncertain. Four meta-analysis studies are present, including an update, 3 of which include the 2 randomized, sham-controlled trials from 2009 but have mixed results. There remains some uncertainty related to the interpretation of these conclusions. While the use of a sham procedure is a major methodologic strength to control for nonspecific (placebo) effects, the sham used is controversial, given that the effect of injecting local anesthetic in the facet capsule and/or periosteum is unknown. Also, the appropriateness of outcome measures used to detect clinically meaningful differences in pain might not have been optimal, because the studies were underpowered to detect differences in clinical response rates. Questions have also been raised about the low percentage of patients screened who participated in the trial, the volume of polymethylmethacrylate injected, and the inclusion of patients with chronic pain.

PERCUTANEOUS VERTEBROPLASTY FOR VERTEBRAL COMPRESSION FRACTURES OF LESS THAN 6 WEEKS OLD

Clinical Context and Therapy Purpose

The purpose of vertebroplasty is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as conservative management, in patients with symptomatic osteoporotic vertebral fractures less than 6 weeks old.

The question addressed in this evidence review is: Does vertebroplasty improve the net health outcome in individuals with symptomatic osteolytic vertebral fractures less than 6 weeks old?

The following PICO was used to select literature to inform this review.

Patients

The relevant population of interest is individuals with symptomatic osteoporotic vertebral fractures less than 6 weeks old. With acute fractures, these individuals experience severe pain, decreased ambulatory function, and a lessened response to conservative medical management.

Interventions

The therapy being considered is vertebroplasty, which is typically performed by an interventional radiologist in an outpatient clinical setting.

Comparators

Comparators of interest include conservative management. A detailed review of the comparators is listed in the above indication. Patients receiving conservative management are typically managed by physical therapists and primary care providers in an outpatient clinical setting.

Outcomes

The general outcomes of interest are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. Symptoms can include back pain and demonstrated fracture on radiography. The most current research available tracks follow-up to 12 months or more. A number of studies have longer term follow-up at more than 5 years, which is ideal for understanding all of the outcomes, particularly the occurrence of new vertebral compression fractures after vertebroplasty.

Study Selection Criteria

Methodologically credible studies were selected using the principles described in the first indication.

Review of Evidence

Randomized Controlled Trials

Vertebroplasty Versus Medical Management With Sham Controls

Clark et al (2016) reported on results from the Safety and Efficacy of Vertebroplasty of Acute Painful Osteoporotic Fractures (VAPOUR) trial (see Table 11).³¹ VAPOUR was a multicenter, double-blind trial of vertebroplasty in 120 patients with vertebral fractures of less than 6 weeks in duration and back pain of at least 7 out of 10 on a numeric rating scale. This trial followed a similar protocol as that used in the Kallmes et al (2009) trial (discussed above). The primary outcome (the percentage of patients with a numeric rating scale score <4 out of 10 at 14 days postprocedure) was met in a greater percentage of patients in the vertebroplasty group (44%) than in the sham control group (21%). This between-group difference was maintained through 6 months.

Other outcome measures were significantly improved in the vertebroplasty group at one or both of the time points (see Table 11). The benefit of vertebroplasty was found predominantly in the thoracolumbar subgroup, with 48% (95% CI, 27% to 68%) more patients meeting the primary endpoint (61% in the vertebroplasty group vs. 13% in the control group). The investigators commented that the thoracolumbar junction is subject to increased dynamic load, and fractures at this junction have the highest incidence of mobility. No benefit from vertebroplasty was found in the non-thoracolumbar subgroup. Postprocedural hospital stay was reduced from a mean of 14 days in the control group to 8.5 days after vertebroplasty, even though physicians who determined the discharge date remained blinded to treatment. In the vertebroplasty group, there were 2 serious adverse events due to sedation and transfer to the radiology table. In the control group, 2 patients developed spinal cord compression; one underwent decompressive surgery and the other, not a surgical candidate, became paraplegic.

Vertebroplasty Versus Medical Management Without Sham Controls

Klazen et al (2010) reported on the Vertebroplasty versus Conservative Treatment in Acute Osteoporotic Vertebral Compression Fractures, an open-label randomized trial of 202 patients at 6 hospitals in the Netherlands and Belgium.³² Of 431 patients eligible for randomization, 229 (53%) had spontaneous pain relief during assessment. Participants with at least one painful osteoporotic vertebral fracture of 6 weeks or less in duration were assigned to vertebroplasty or conservative management. The primary outcome was pain relief of 3 points measured on a 10-point visual analogue scale at 1 month and 1 year.

A total of 101 subjects were enrolled in the treatment group and the control arm; 81% completed 12-month follow-up. There were no significant differences in the primary outcome (pain relief of 3 points) measured at 1 month and 1 year. Vertebroplasty resulted in greater pain relief than did medical management through 12 months (<0.001); there were significant between-group differences in mean visual analogue scale scores at 1 month or at 1 year. Survival analysis showed significant pain relief was quicker (29.7 days vs. 115.6 days) and was achieved by more patients after vertebroplasty than after conservative management.

Yi et al (2014) assessed the occurrence of new vertebral compression fractures after treatment with cement augmenting procedures (vertebroplasty or kyphoplasty) vs conservative treatment in an RCT with 290 patients (363 affected vertebrae).³³ Patients treated conservatively had a mean length of stay of 13.7 days. Return to usual activity occurred at 1 week for 87.6% of operatively treated patients and 2 months for 59.2% of conservatively treated patients. All patients were evaluated with radiographs and magnetic resonance imaging at 6 months and then at yearly intervals until the last follow-up session. At a mean follow-up of 49.4 months (range, 36-80 months), 10.7% of patients had experienced 42 new symptomatic vertebral compression fractures. There was no significant difference in the incidence of new vertebral fractures between the operative (18 total; 9 adjacent, 9 nonadjacent) and conservative (24 total; 5 adjacent, 16 nonadjacent, 3 same level) groups but the mean time to a new fracture was significantly shorter in the operative group (9.7 months) than in the nonoperative group (22.4 months).

Leali et al (2016) published a brief report on a multicenter RCT enrolling 400 patients with osteoporotic thoracic or lumbar vertebral compression fractures who were treated with vertebroplasty or conservative therapy.³⁴ Fractures were treated within 2 weeks of pain onset. Details of randomization and rates of follow-up were not reported. At 1 day after treatment, the vertebroplasty group had a reduction in pain scores and improvement in physical function, with visual analogue scale pain scores decreasing from 4.8 (maximum, 5.0) to 2.3 ($p=0.023$) and

Oswestry Disability Index scores improving from 53.6% to 31.7% (p=0.012). Sixty-five percent of patients treated with vertebroplasty had stopped all analgesic use within 48 hours. The conservatively managed group showed no benefit in the first 48 hours, but by 6 weeks visual analogue scale and Oswestry Disability Index scores were described as similar in both groups (specific data not reported). Evaluation of this trial was limited by incomplete reporting.

Yang et al (2016) compared vertebroplasty with conservative therapy in 135 patients over 70 years of age with severe back pain due to an osteoporotic vertebral fracture after minor or mild trauma.³⁵ Vertebroplasty was performed at a mean of 8.4 days after pain onset. Patients in the conservative therapy group were placed on bed rest and analgesics for at least 2 weeks after diagnosis, followed by bracing and assistive devices. All patients receiving vertebroplasty could stand and walk with a brace at 1 day posttreatment, while only 12 (23.5%) patients in the control group could stand up and walk after 2 weeks of bed rest. The average duration of bed rest from pain onset was 7.8 days (range, 2-15 days) in the vertebroplasty group compared with 32.5 days (range, 14-60 days) in the conservative therapy group. At 1-year follow-up, there was a similar percentage of additional compression fractures but a significantly higher complication rate in the conservative therapy group (35.3%) than in the vertebroplasty group (16.1%; p<0.001). Complications included pneumonia, urinary tract infection, deep vein thrombosis, depression, and sleep disorders.

Table 11. Summary of Key RCT Characteristics Involving Vertebroplasty vs Medical Management without Sham Controls

Study; Trial	Countries	Sites	Dates	Participants (N)	Interventions	
					Active (n)	Comparator (n)
Klazen et al (2010) ³² ,	EU	6	2005-2008	Patients >50 years with radiographically confirmed VCF, back pain r <6 weeks, VAS >5	Vertebroplasty (101)	Medical Management without Sham Controls (101)
Yi et al (2014) ³³ ,	China	1	2005-2009	Patients with OCVF	PVP or PKP(169)	Conservative treatment(121)
Leali et al (2010) ³⁴ ,	International	4	NR	Post-menopausal women with 1 thoracic or lumbar symptomatic OVCF caused by primary or secondary osteoporosis.	PVP including analgesic and osteoporosis medication (200)	Conservative care including analgesic and osteoporosis medication (200)
Yang et al (2015) ³⁵ ,	China	1	2009-2011	Patients > 70 years with acute OVCF, severe pain from minor or mild trauma	PVP (56 at 1 y)	Conservative treatment (51 at 1 y)

NR: not reported; OVCF: osteoporotic vertebral compression fractures; PKP; percutaneous kyphoplasty; PVP: percutaneous vertebroplasty; RCT: randomized controlled trial; VCF: vertebral compression fracture; VAS: visual analog scale.

Table 12. Summary of Key RCT Results Involving Vertebroplasty Versus Medical Management without Sham Controls

Study	VAS	Quality of Life	Refracture Rate
Klazen et al (2010) ^{32,}			
Mean difference between groups in reduction of mean VAS score from baseline		RMDQ1	Median follow-up of 12.0 months (range: 1-24)
Month 1 (CI)	2.0 (1.13-2.80)	PVP: 12.5	PVP: 18 (16.48%)
p-value	< 0.0001	Control: 13.5	Control: 30 (24.71%)
Month 12 (CI)	2.0 (1.13-2.80)	PVP: 9	
p-value	<0.0001	Control: 12	
Yi et al (2014)			
Month 12 (%)	-	-	PVP/PKP: 18 (8.28%)
	-	-	Control: 24 (19.83%)
	-	-	Time interval of recompression
Intervention	-	-	9.7 ± 17.8 months
Control			22.4 ± 7.99 months
p-value			0.017
	-	ODI, %	
Leali et al (2016)			-
Intervention 24 hours after surgery, mean	2.3	31.7	-
p-value	≤0.023	≤0.012	
Yang et al (2015) ²			
Analysis of variance models, Month 1 (SD)	PVP: 2.4±1 Control: 4.8±1	PVP: 48±10 Control: 71±7	
Analysis of variance models, Month 12 (SD) p-value	PVP: 1.8±0.3 Control: 3±0.5-	PVP: 30±5 Control:-	PVP: 5 (8.9%) Control: 4 (7.8) <0.0001

CI: 95% confidence interval; ODI: Oswestry Disability Index; PKP: percutaneous kyphoplasty; PVP: percutaneous vertebroplasty; RCT: randomized controlled trials; RMDQ: Roland-Morris Disability Questionnaire; VAS: visual analogue scale; SD: standard deviation.

¹The RMDQ results from the Klazen paper are based on estimates due to the graphical presentation of the results, rather than the reporting of the numerical values.

² The results from the Yang paper are based on estimates due to the graphical presentation of the results; numerical results not reported.

Table 13. Relevance Study Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Follow-Up ^e
Klazen et al (2010) ^{32,}				3. None reported	
Yi et al (2014) ^{33,}	4. Selection criteria for PVP or PKP unclear, some patients had > fracture				
Leali et al (2010) ^{34,}	1.Limited to post-menopausal women				1,2 Follow-up period limited to < 6 months
Yang et al (2015) ^{35,}	4. Study population limited to > 70 years of age at single spine center				

PVP: percutaneous vertebroplasty; PKP: percutaneous kyphoplasty.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

^e Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

Table 14. Study Design and Conduct Limitations

Study	Allocation ^a	Blinding ^b	Selective Reporting ^c	Follow-Up ^d	Power ^e	Statistical ^f
Klazen et al (2010) ^{32,}		1,2. No masking.				
Yi et al(2014) ^{33,}						
Leali et al (2010) ^{34,}		1,2,3, unclear if masking occurred	2. Outcomes beyond 48 hours post-surgery not reported.			
Yang et al (2015) ^{35,}		1,2,3 No masking				3. Results reported only in graphic form

The study limitations stated in this table are those notable in the current review; this is not a comprehensive limitations assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Follow-Up key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Intervention is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Intervention is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

Section Summary: Percutaneous Vertebroplasty for Vertebral Compression Fractures of Less Than 6 Weeks Old

In a sham-controlled randomized trial, where no anesthetic was injected into the periosteum, there was a significant benefit of vertebroplasty in patients who had severe pain of fewer than 6 weeks in duration following vertebral fracture at the thoracolumbar junction. Other RCTs without sham controls have reported that vertebroplasty is associated with significant improvements in pain, earlier improvements in function, and reductions in the duration of bed rest compared with conservatively managed patients.

PERCUTANEOUS SACROPLASTY

Clinical Context and Therapy Purpose

Sacral insufficiency fractures are the consequence of stress on weakened bone and often cause low back pain in the elderly population.¹ Osteoporosis is the most common risk factor for sacral insufficiency fractures. Lourie (1982) described spontaneous fracture of the sacrum in patients with osteoporosis as presenting as lower back and buttock pain with or without referred pain in the legs.³⁶ Although common, sacral insufficiency fractures can escape detection due to low provider suspicion and poor sensitivity on plain radiographs, slowing the application of appropriate intervention.

The purpose of sacroplasty is to provide a treatment option that is an alternative to or an improvement on existing therapies, such as conservative management, in patients with sacral insufficiency fractures.

The question addressed in this evidence review is: Does sacroplasty improve the net health outcome in individuals with sacral insufficiency fractures?

The following PICO was used to select literature to inform this review.

Patients

The relevant population of interest is individuals with sacral insufficiency fractures. Sacral insufficiency fractures are a stress fracture, resulting from a regular stress applied to a bone with reduced elasticity. Often, these fractures are associated with underlying metabolic bone disease condition like osteoporosis. Examples of risk factors include corticosteroid therapy use, female sex, pelvic radiation, rheumatoid arthritis, and hyperparathyroidism.

Interventions

The therapy being considered is sacroplasty, a minimally invasive procedure for treating pathological fractures of the sacral vertebral body or sacral ala. The procedure involves percutaneous insertion of one or more bone needles into the sacrum and injection of bone cement under fluoroscopy and/or computed tomography visual guidance. This intervention is provided by an interventional radiologist typically in an outpatient setting.

Comparators

Comparators of interest include conservative management. Conservative management includes physical therapy, analgesics, narcotics, and hormone treatments. Examples of conservative management for sacral insufficiency fractures are varied and can include bed rest and pain medication to early physical therapy. Sacral insufficiency fractures are managed by orthopedists and physical therapists.

Outcomes

The general outcomes of interest are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. Possible negative outcomes include complications with sedation, cement leakage into the presacral space, spinal canal, sacral foramen, or sacroiliac joint, and possible spinal compression due to extravasation of cement. At least one year of follow-up is desirable to adequately evaluate outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the principles described in the first indication.

Sacroplasty is an evolving technique achieved using numerous methods (short-axis, long-axis, balloon-assisted short-axis, iliosacral screws). No randomized trials of sacroplasty were identified. Frey et al (2008) conducted the largest prospective observational cohort study, assessing 52 consecutive patients undergoing sacroplasty for sacral insufficiency fractures using the short-axis technique.³⁷ Patients had a mean age of 75.9 years, mean duration of symptoms of 34.5 days (range, 4-89 days), and mean visual analogue scale score of 8.1 at baseline. Improvements in visual analogue scale scores were measured at 30 minutes and 2, 4, 12, 24, and 52 weeks postprocedure. At each interval, statistically significant improvements over baseline were observed and maintained through 52 weeks.

Kortman et al (2013) reported on the largest series, a retrospective multicenter analysis.³⁸ They evaluated 204 patients with painful sacral insufficiency fractures and 39 patients with symptomatic sacral lesions treated with the short-axis or long-axis technique. One hundred sixty-nine patients had bilateral sacral insufficiency fractures, and 65 patients had additional fractures of the axial skeleton. Visual analogue scale scores improved from 9.2 before treatment to 1.9 after treatment in patients with sacral insufficiency fractures and from 9.0 to 2.6 in patients with sacral lesions. There was one case of radicular pain due to extravasation of cement requiring surgical decompression.

Frey et al (2017) reported on patients treated with percutaneous sacroplasty, particularly the long-term efficacy of sacroplasty versus nonsurgical management.³⁹ This prospective, observational cohort study spanned 10 years and comprised 240 patients with sacral insufficiency fractures. Thirty-four patients were treated with nonsurgical methods, and 210 patients were treated with sacroplasty. Pain, as measured by visual analogue scale, was recorded before treatment and at several follow-ups. Mean pretreatment visual analogue scale for the sacroplasty group was 8.29;

for the nonsurgical treatment group, it was 7.47. Both forms of treatment resulted in significant visual analogue scale improvement from pretreatment to the 2-year follow-up ($p < 0.001$). However, the sacroplasty treatment group experienced significant visual analogue scale score improvement consistently at many of the follow-up points (pretreatment to post [$p < 0.001$]; posttreatment through 2 weeks [$p > 0.001$]; 12 weeks through 24 weeks [$p = 0.014$]; 24 weeks through 1 year [$p = 0.002$]). Meanwhile, the group with nonsurgical treatment only experienced one significant pain improvement score, which was at the 2-week follow-up posttreatment ($p = 0.002$). One major limitation of this study was that the nonsurgical treatment group was not followed up at the 10-year mark whereas the sacroplasty group did receive follow-up.

There are several retrospective reviews with roughly 50 patients per publication. One reported by Dougherty et al (2014) described a series of 57 patients treated with sacroplasty for sacral insufficiency fractures.⁴⁰ The short- or the long-axis approach was dictated by the length and type of the fracture and patient anatomy. Follow-up data at 2.5 weeks were available for 45 (79%) patients, and the outcome measures were inconsistent. For example, activity pain scores were collected from 13 patients, and rest pain scores were collected from 29 patients. Of the 45 patients with outcomes data, 37 (82%) had experienced a numeric or descriptive decrease from initial pain of at least 30%.

Adverse Events

There are complications related to cement leakage with sacroplasty that are not observed with vertebroplasty. Leakage of polymethylmethacrylate into the presacral space, spinal canal, sacral foramen, or sacroiliac joint may result in pelvic injection of polymethylmethacrylate, sacral nerve root or sacral spinal canal compromise, or sacroiliac joint dysfunction.⁴¹ Performing sacroplasty only on zone 1 fractures can minimize these risks.⁴²

Section Summary: Percutaneous Sacroplasty

No RCTs evaluating percutaneous sacroplasty for sacral insufficiency were identified. The available evidence includes 2 prospective cohort studies and several retrospective series. These studies have reported rapid and sustained decreases in pain following percutaneous sacroplasty. Additional reports are mostly consistent in reporting immediate improvement following the procedure. Due to the limited number of patients and the retrospective nature of the evidence base, harms associated with sacroplasty have not been adequately studied. The small numbers of treated patients leave uncertainty regarding the impact of sacroplasty on health outcomes.

Summary of Evidence

For individuals who have symptomatic osteoporotic vertebral fractures between 6 weeks and 1 year old who receive vertebroplasty, the evidence includes 2 randomized sham-controlled trials, nonblinded randomized controlled trials (RCTs) comparing vertebroplasty with conservative management, and systematic reviews of these RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. Despite the completion of multiple RCTs, including 2 with sham controls, the efficacy of vertebroplasty for painful osteoporotic compression fractures remains uncertain. Two meta-analysis studies which included the 2 sham-controlled trials have demonstrated mixed results. The 2 studies had methodologic issues, including the choice of sham procedure and the potential of the sham procedure to have a therapeutic effect by reducing pain. Questions have also been raised about the low percentage of patients screened who participated in the trial, the volume of polymethylmethacrylate injected, and the inclusion of patients with chronic pain. Overall,

conclusions about the effect of vertebroplasty remain unclear. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with symptomatic osteoporotic vertebral fractures less than 6 weeks old who receive vertebroplasty, the evidence includes a randomized sham-controlled trial and nonblinded RCTs comparing vertebroplasty with conservative management. Relevant outcomes are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. For acute fractures, conservative therapy consisting of rest, analgesics, and physical therapy is an option, and symptoms will resolve in a large percentage of patients with conservative treatment only. However, a sham-controlled randomized trial in patients who had severe pain of fewer than 6 weeks in duration found a significant benefit of vertebroplasty for the treatment of osteoporotic vertebral fracture at the thoracolumbar junction. Other RCTs without sham controls have reported that vertebroplasty is associated with significant improvements in pain and reductions in the duration of bed rest. Given the high morbidity associated with extended bed rest in older adults, this procedure is considered to have a significant health benefit. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals with sacral insufficiency fractures who receive sacroplasty, the evidence includes 2 prospective cohort studies and a case series. Relevant outcomes are symptoms, functional outcomes, quality of life, hospitalizations, medication use, and treatment-related morbidity. No RCTs have been reported. The available evidence includes a prospective cohort study and a retrospective series of 243 patients. These studies have reported rapid and sustained decreases in pain following percutaneous sacroplasty. Additional literature has mostly reported immediate improvements following the procedure. However, due to the small size of the evidence base, the harms associated with sacroplasty have not been adequately studied. The evidence is insufficient to determine the effects of the technology on health outcomes.

Vertebroplasty has been investigated as an intervention to provide mechanical support and symptomatic relief in patients with an osteoporotic vertebral compression fracture and in those with osteolytic lesions of the spine (i.e., multiple myeloma, metastatic malignancies). Clinical input obtained in 2008 provided uniform support for the use of vertebroplasty in painful osteoporotic fractures. Reconsideration of the available evidence and evaluation of the input led to a conclusion that, consistent results of numerous case series, including large prospective reports, the evidence was sufficient to determine that vertebroplasty is a reasonable treatment option in patients with vertebral fractures who have failed to respond to conservative treatment (at least 6 weeks with analgesics, physical therapy, and rest). It is also clinically reasonable to consider the evidence supporting the clinical benefit of vertebroplasty in the osteoporotic vertebral fracture to support its use in osteolytic lesions of the spine (e.g., multiple myeloma, metastatic malignancies).

SUPPLEMENTAL INFORMATION

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2014 Input

In response to requests, input was received from 2 physician specialty societies and 3 academic medical centers while this policy was under review in 2014. Input was sought on the treatment of acute vertebral fractures when there is severe pain that has led to hospitalization or persists at a level that prevents ambulation, and on the treatment of traumatic fractures that have remained symptomatic after 6 weeks of conservative treatment. Input on these issues was mixed.

2008 Input

In response to requests, input was received from 5 physician specialty societies and 2 academic medical centers while this policy was under review in 2008. Unsolicited input was received from a sixth physician specialty society. All reviewers disagreed with the proposed policy and provided references in support of the use of vertebroplasty.

Practice Guidelines and Position Statements

American College of Radiology

In 2018, the American College of Radiology (ACR) revised its Appropriateness Criteria for the use of percutaneous vertebral augmentation in the management of vertebral compression fractures.⁴³ Table 15 shows the appropriateness categories for each variant.

Table 15. ACR Appropriateness Criteria for the Use of Percutaneous Vertebral Augmentation for the Management of Vertebral Compression Fractures

Variants	Appropriateness Category
"New symptomatic compression fracture identified on radiographs or CT. No known malignancy."	May Be Appropriate
"Osteoporotic compression fracture, with or without edema on MRI and no 'red flags.' With or without spinal deformity, worsening symptoms, or pulmonary dysfunction."	Usually Appropriate
"Asymptomatic pathologic spinal fracture with or without edema on MRI."	May Be Appropriate
"Pathologic spinal fracture with severe and worsening pain."	Usually Appropriate
"Pathologic spinal fracture with spinal deformity or pulmonary dysfunction."	Usually Appropriate

CT: computed tomography; MRI: magnetic resonance imaging; ACR: American College of Radiology.

American College of Radiology, et al

In 2014, the American College of Radiology (ACR) and 7 other medical specialty associations, including the Society for Interventional Radiology, updated a 2012 joint position statement on percutaneous vertebral augmentation.⁴⁴ The statement indicated that "percutaneous vertebral augmentation with the use of vertebroplasty or kyphoplasty is a safe, efficacious, and durable procedure in appropriate patients with symptomatic osteoporotic and neoplastic fractures, when performed in accordance with published standards...only when nonoperative medical therapy has not provided adequate pain relief or pain is significantly altering the patient's quality of life⁴⁴."

Society for Interventional Radiology

In a 2014 quality improvement guideline from the Society for Interventional Radiology, failure of medical therapy was defined as follows⁴⁵:

1. "For a patient rendered nonambulatory as a result of pain from a weakened or fractured vertebral body, pain persisting at a level that prevents ambulation despite 24 hours of analgesic therapy;
2. For a patient with sufficient pain from a weakened or fractured vertebral body that physical therapy is intolerable, pain persisting at that level despite 24 hours of analgesic therapy; or
3. For any patient with a weakened or fractured vertebral body, unacceptable side effects such as excessive sedation, confusion, or constipation as a result of the analgesic therapy necessary to reduce pain to a tolerable level."

American Academy of Orthopaedic Surgeons

In 2011, the American Academy of Orthopaedic Surgeons (AAOS) published practice guidelines on the treatment of osteoporotic spinal compression fractures.⁴⁶ The AAOS approved "a Strong recommendation against the use of vertebroplasty for patients who present with an acute osteoporotic spinal compression fracture and are neurologically intact."

National Institute for Health and Care Excellence

In 2003, the National Institute for Health and Care Excellence (NICE) concluded in its guidance on percutaneous vertebroplasty that the current evidence on the safety and efficacy of vertebroplasty for vertebral compression fractures appeared "adequate to support the use of this procedure" to "provide pain relief for people with severe painful osteoporosis with loss of height and/or compression fractures of the vertebral body...."⁴⁷ The guidance also recommended that the procedure be limited to patients whose pain is refractory to more conservative treatment. A 2013 NICE guidance indicated that percutaneous vertebroplasty and percutaneous balloon kyphoplasty "are recommended as options for treating osteoporotic vertebral compression fractures" in persons having "severe, ongoing pain after a recent, unhealed vertebral fracture despite optimal pain management" and whose "pain has been confirmed to be at the level of the fracture by physical examination and imaging."⁴⁸

In 2008, NICE issued guidance on the diagnosis and management of adults with metastatic spinal cord compression.⁴⁹ This guidance indicated that vertebroplasty or kyphoplasty should be considered for "patients who have vertebral metastases and no evidence of metastatic spinal cord compression or spinal instability if they have: mechanical pain resistant to conventional pain management, or vertebral body collapse."

U.S. Preventive Services Task Force Recommendations

Not applicable.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 16.

Table 16. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT02902250	The Comparative Study About the Effect of Vertebral Body Decompression Procedure and Conservative Treatment for Benign Vertebral Compression Fracture - Prospective Randomized Control Study	80	Apr 2020
NCT02489825	Pilot Study: Does Preventive Adjacent Level Cement Augmentation Positively Affect Reoperation Rates After Osteoporotic Vertebral Compression Fractures?	100	June 2019

NCT: national clinical trial.

CODING

The following codes for treatment and procedures applicable to this policy are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

CPT/HCPCS

- 22510 Percutaneous vertebroplasty (bone biopsy included when performed), 1 vertebral body, unilateral or bilateral injection, inclusive of all imaging guidance; cervicothoracic
- 22511 Percutaneous vertebroplasty (bone biopsy included when performed), 1 vertebral body, unilateral or bilateral injection, inclusive of all imaging guidance; lumbosacral
- 22512 Percutaneous vertebroplasty (bone biopsy included when performed), 1 vertebral body, unilateral or bilateral injection, inclusive of all imaging guidance; each additional cervicothoracic or lumbosacral vertebral body (List separately in addition to code for primary procedure)
- 0200T Percutaneous sacral augmentation (sacroplasty), unilateral injection(s), including the use of a balloon or mechanical device, when used, 1 or more needles, includes imaging guidance and bone biopsy, when performed
- 0201T Percutaneous sacral augmentation (sacroplasty), bilateral injections, including the use of a balloon or mechanical device, when used, 2 or more needles, includes imaging guidance and bone biopsy, when performed

ICD-10 Diagnoses

- C41.2 Malignant neoplasm of vertebral column
- C79.51 Secondary malignant neoplasm of bone
- C79.52 Secondary malignant neoplasm of bone marrow
- C90.00 Multiple myeloma not having achieved remission
- C90.01 Multiple myeloma in remission
- D18.09 Hemangioma of other sites
- D47.Z9 Other specified neoplasms of uncertain behavior of lymphoid, hematopoietic and related tissue
- M48.50XA Collapsed vertebra, not elsewhere classified, site unspecified, initial encounter for fracture

- M48.51XA Collapsed vertebra, not elsewhere classified, occipito-atlanto-axial region, initial encounter for fracture
- M48.52XA Collapsed vertebra, not elsewhere classified, cervical region, initial encounter for fracture
- M48.53XA Collapsed vertebra, not elsewhere classified, cervicothoracic region, initial encounter for fracture
- M48.54XA Collapsed vertebra, not elsewhere classified, thoracic region, initial encounter for fracture
- M48.55XA Collapsed vertebra, not elsewhere classified, thoracolumbar region, initial encounter for fracture
- M48.56XA Collapsed vertebra, not elsewhere classified, lumbar region, initial encounter for fracture
- M48.57XA Collapsed vertebra, not elsewhere classified, lumbosacral region, initial encounter for fracture
- M48.58XA Collapsed vertebra, not elsewhere classified, sacral and sacrococcygeal region, initial encounter for fracture
- M80.08XA Age-related osteoporosis with current pathological fracture, vertebra(e), initial encounter for fracture
- M80.88XA Other osteoporosis with current pathological fracture, vertebra(e), initial encounter for fracture
- M81.0 Age-related osteoporosis without current pathological fracture
- M81.8 Other osteoporosis without current pathological fracture
- M84.48XA Pathological fracture, other site, initial encounter for fracture
- M84.58XA Pathological fracture in neoplastic disease, vertebrae, initial encounter for fracture
- M84.68XA Pathological fracture in other disease, other site, initial encounter for fracture

REVISIONS

04-21-2005	Added "or kyphoplasty" to policy #C.
12-14-2005	In "Policy" section, #C., added 'and cervical percutaneous vertebroplasty and kyphoplasty' based on Radiology Liaison Committee recommendations from 02-12-2002.
	In "Coding" CPT/HCPCS section, added CPT codes 22523, 22524, and 22525, and added "or vertebral augmentation including cavity creation" to CPT code 76012 to reflect changes in CPT book.
	In "Coding" CPT/HCPCS section, deleted HCPCS codes S2360 and S2361 because 'cervical' is considered E/I by the Radiology Liaison Committee 02-12-2002.
12-21-2006	In "Coding", Covered Diagnosis section, added Percutaneous vertebroplasty or Kyphoplasty – CPT Codes – 22520, 22521, 22522, 22523, 22524, 22525, 76012, 76013, S2362, S2363 to the current listing of diagnosis codes.
07-27-2006 effective 10-01-2006	Deleted S2362 and S2363, the codes were deleted from HCPCS 4-1-06.
10-31-2006 effective 01-01-2007	In "Coding", CPT/HCPCS deleted CPT codes 76012 and 76013 and added CPT codes 72291 and 72292 due to the 2007 CPT changes.
07-23-2009	Removed percutaneous vertebroplasty and kyphoplasty policy language from the policy entitled: Minimally Invasive Procedures for Spine Pain creating a free-standing policy entitled: Percutaneous Vertebroplasty, Kyphoplasty and Sacroplasty.
	Description section: Updated description to reflect discussion of percutaneous vertebroplasty, kyphoplasty and sacroplasty
	Policy section:

	<p>Revised policy language from:</p> <p>C. Percutaneous vertebroplasty or kyphoplasty is considered medically necessary after failure of standard medical therapy in patients when any of the following criteria is met. Medical conditions not listed and cervical percutaneous vertebroplasty and kyphoplasty will be denied experimental/investigational.</p> <ol style="list-style-type: none"> 1. Osteolytic vertebral metastasis or myeloma with severe back pain related to destruction of the vertebral body not involving the major part of the cortical bone, and chemotherapy and radiation therapy have failed to relieve symptoms; or 2. Vertebral hemangiomas with aggressive clinical signs (severe pain or nerve compression) and/or aggressive radiological signs, and radiation therapy has failed to relieve symptoms; or 3. Osteoporotic vertebral collapse with persistent debilitating pain that has not responded to accepted standard medical therapy as documented in the medical records. Standard medical therapy may include initial bed rest with progressive activity, analgesics, physical therapy, bracing and exercises to correct postural deformity and increase muscle tone, salmon calcitonin, bisphosphonates and calcium supplementation; or 4. Painful vertebral eosinophilic granuloma with spinal instability. <p>To:</p> <p>Percutaneous vertebroplasty and kyphoplasty may be considered medically necessary for the treatment of:</p> <p>severe pain due to osteolytic lesions of the spine related to multiple myeloma or metastatic malignancies vertebral hemangiomas with pain, nerve compression or aggressive radiologic signs, and radiation therapy has failed to relieve symptoms painful vertebral eosinophilic granuloma vertebral compression fracture with persistent debilitating pain</p> <p>Sacroplasty may be considered medically necessary for the treatment of sacral insufficiency fractures that have failed to respond to conservative treatment.</p> <p>Percutaneous vertebroplasty, kyphoplasty and sacroplasty are considered experimental / investigational for all other indications.</p>
	<p>Rationale section: Added Rationale section.</p>
	<p>Coding section: Added CPT/HCPCS Codes: 0200T, 0201T, S2360, S2361. Deleted ICD-9 Code: 213.2. Added ICD-9 Codes: 203.01, 238.6.</p>
<p>01-01-2012</p>	<p>In the Coding section: Revised CPT nomenclature for the following codes: 22520, 22521, 22522</p>
<p>10-04-2013</p>	<p>Added Medical Policy and Coding Disclaimers. Description section updated. In the Policy section:</p> <ul style="list-style-type: none"> • Revised medical policy language from the following: Percutaneous vertebroplasty and kyphoplasty may be considered medically necessary for the treatment of: <ol style="list-style-type: none"> A. severe pain due to osteolytic lesions of the spine related to multiple myeloma or metastatic malignancies B. vertebral hemangiomas with pain, nerve compression or aggressive radiologic signs, and radiation therapy has failed to relieve symptoms C. painful vertebral eosinophilic granuloma D. osteoporotic vertebral compression fracture with persistent debilitating pain

	<p>Sacroplasty may be considered medically necessary for the treatment of sacral insufficiency fractures that have failed to respond to conservative treatment. Percutaneous vertebroplasty, kyphoplasty and sacroplasty are considered experimental / investigational for all other indications.</p> <p>Rationale section updated.</p> <p>In Coding section:</p> <ul style="list-style-type: none"> Added ICD-10 Diagnosis (<i>Effective October 1, 2014</i>) <p>Reference section updated.</p>
12-31-2013	<p>In Policy section:</p> <ul style="list-style-type: none"> In Item I, E, added "/bone scan" to read "The treatment of MRI / bone scan documented acute osteoporotic vertebral..."
01-01-2015	<p>In Coding section:</p> <ul style="list-style-type: none"> Added CPT Codes: 22510, 22511, 22512, 22513, 22514, 22515 (Effective January 1, 2015) Deleted CPT Codes: 22520, 22521, 22522, 22523, 22524, 22525, 72291, 72292 (Effective January 1, 2015) Revised CPT Codes: 0200T, 0201T (Effective January 1, 2015)
10-21-2015	<p>Policy title changed from "Percutaneous Vertebroplasty, Kyphoplasty and Sacroplasty". A new medical policy was created with information on kyphoplasty titled "Percutaneous Balloon Kyphoplasty and Mechanical Vertebral Augmentation".</p> <p>Updated Description section.</p> <p>In Policy section:</p> <ul style="list-style-type: none"> In Item A, removed "or kyphoplasty" to read "Percutaneous vertebroplasty may be considered medically necessary for:" In Item B, removed "or kyphoplasty" to read, "Percutaneous vertebroplasty is considered experimental / investigational for all other indications, including use in acute vertebral fractures due to trauma." <p>Updated Rationale section.</p> <p>In Coding section:</p> <ul style="list-style-type: none"> Removed CPT codes 22513, 22514, and 22515. <p>Updated References section.</p>
10-21-2015	<p>Published 10-20-2015, effective 10-21-2015.</p> <p>In Policy section:</p> <ul style="list-style-type: none"> In Item B, removed "or kyphoplasty" as indicated above to read, "Percutaneous vertebroplasty is considered experimental / investigational for all other indications, including in acute vertebral fractures due to trauma." (Not included as intended with previous 10-21-2015 publication.)
01-01-2016	<p>In Coding section:</p> <ul style="list-style-type: none"> Removed HCPCS codes S2360 and S2361.
01-19-2018	<p>Updated Description section.</p> <p>In Policy section:</p> <ul style="list-style-type: none"> Removed previous Items A 2 and 3, "2. Vertebral hemangiomas with pain, nerve compression or aggressive radiologic signs, and radiation therapy has failed to relieve symptoms; OR 3. Painful vertebral eosinophilic granuloma; OR" In previous Item A 4 (now Item A 2), removed "MRI documented acute," "compression," "with persistent debilitating pain," "with graduated activity, back bracing," "and calcitonin," "or these treatments are contraindicated," and added "symptomatic" to read, "The treatment of symptomatic osteoporotic vertebral fractures that have failed to respond to conservative treatment (e.g., rest, analgesics, physical therapy) for at least 6 weeks;" In previous Item A 5 (now Item A 3), removed "MRI/bone scan documented," "acute," "compression," "with persistent debilitating pain requiring," "admission and parenteral narcotics for treatment" and added "symptomatic," "that are less than 6 weeks in

	duration that have led to," "or persists at a level that prevents ambulation" to read, "The treatment of symptomatic osteoporotic vertebral fractures that are less than 6 weeks in duration that have led to hospitalization or persists at a level that prevents ambulation."
	Updated Rationale section.
	In Coding section: <ul style="list-style-type: none"> ▪ Removed ICD-9 codes.
	Updated References section.
05-23-2018	Updated Description section.
	Updated Rationale section.
	Updated References section.
08-01-2018	Updated Rationale section.
05-21-2019	Updated Description section.
	Updated Rationale section.
	Updated References section.
05-23-2021	Updated Description section.
	In Policy section: <ul style="list-style-type: none"> • Added "osteoporosis or" to Item B to read "Percutaneous vertebroplasty is considered experimental / investigational for all other indications, including use in acute vertebral fractures due to osteoporosis or trauma."
	Updated Rationale section.
	Updated References section.

REFERENCES

1. Gotis-Graham I, McGuigan L, Diamond T, et al. Sacral insufficiency fractures in the elderly. J Bone Joint Surg Br. Nov 1994;76(6):882-886. PMID 7983111
2. Lin J, Lachmann E, Nagler W. Sacral insufficiency fractures: a report of two cases and a review of the literature. J Womens Health Gen Based Med. Sep 2001;10(7):699-705. PMID 11571100
3. Bae H, Hatten HP, Jr., Linovitz R, et al. A prospective randomized FDA-IDE trial comparing Cortoss with PMMA for vertebroplasty: a comparative effectiveness research study with 24-month follow-up. Spine (Phila Pa 1976). Apr 1 2012;37(7):544-550. PMID 21738093
4. Dehdashti AR, Martin JB, Jean B, et al. PMMA cementoplasty in symptomatic metastatic lesions of the S1 vertebral body. Cardiovasc Intervent Radiol. May-Jun 2000;23(3):235-237. PMID 10821903
5. Marcy PY, Palussiere J, Descamps B, et al. Percutaneous cementoplasty for pelvic bone metastasis. Support Care Cancer. Nov 2000;8(6):500-503. PMID 11094996
6. Aretxabala I, Fraiz E, Perez-Ruiz F, et al. Sacral insufficiency fractures. High association with pubic rami fractures. Clin Rheumatol. Oct 2000;19(5):399-401. PMID 11055834
7. Leroux JL, Denat B, Thomas E, et al. Sacral insufficiency fractures presenting as acute low-back pain. Biomechanical aspects. Spine (Phila Pa 1976). Dec 1993;18(16):2502-2506. PMID 8303454
8. Newhouse KE, el-Khoury GY, Buckwalter JA. Occult sacral fractures in osteopenic patients. J Bone Joint Surg Am. Dec 1992;74(10):1472-1477. PMID 1364816
9. Stratford PW, Binkley J, Solomon P, et al. Defining the minimum level of detectable change for the Roland-Morris questionnaire. Phys Ther. 1996 Apr;76(4):359-65. PMID: 8606899
10. Katz J, Melzack R. Measurement of pain. Surg Clin North Am. 1999 Apr;79(2):231-52. PMID: 10352653
11. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous Vertebroplasty. TEC Assessments. 2000;Volume 15:Tab 21.

12. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty for vertebral fractures caused by osteoporosis, malignancy, or hemangioma. TEC Assessments. 2004;Volume 19:Tab 13.
13. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty for vertebral fractures caused by osteoporosis or malignancy. TEC Assessments. 2005;Volume 20:Tab 6.
14. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty or kyphoplasty for vertebral fractures caused by osteoporosis or malignancy. TEC Assessments. 2008;Volume 23:Tab 5.
15. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty or kyphoplasty for vertebral fractures caused by osteoporosis. TEC Assessments. 2009;Volume 24:Tab 7.
16. Blue Cross and Blue Shield Technology Evaluation Center (TEC). Percutaneous vertebroplasty for vertebral fractures caused by osteoporosis. TEC Assessments. 2010;Volume 25:Tab 9.
17. Buchbinder R, Johnston RV, Rischin KJ, et al. Percutaneous vertebroplasty for osteoporotic vertebral compression fracture. *Cochrane Database Syst Rev*. 2018 Apr 4;4:CD006349. PMID: 29618171
18. Staples MP, Kallmes DF, Comstock BA, et al. Effectiveness of vertebroplasty using individual patient data from two randomised placebo controlled trials: meta-analysis. *BMJ*. Jul 12 2011;343:d3952. PMID 21750078
19. Xie L, Zhao ZG, Zhang SJ, et al. Percutaneous vertebroplasty versus conservative treatment for osteoporotic vertebral compression fractures: An updated meta-analysis of prospective randomized controlled trials. *Int J Surg*. Nov 2017;47:25-32. PMID 28939236
20. Buchbinder R, Osborne RH, Ebeling PR, et al. A randomized trial of vertebroplasty for painful osteoporotic vertebral fractures. *N Engl J Med*. Aug 6 2009;361(6):557-568. PMID 19657121
21. Kallmes DF, Comstock BA, Heagerty PJ, et al. A randomized trial of vertebroplasty for osteoporotic spinal fractures. *N Engl J Med*. Aug 6 2009;361(6):569-579. PMID 19657122
22. Kroon F, Staples M, Ebeling PR, et al. Two-year results of a randomized placebo-controlled trial of vertebroplasty for acute osteoporotic vertebral fractures. *J Bone Miner Res*. Jun 2014;29(6):1346-1355. PMID 24967454
23. Ostelo RW, Deyo RA, Stratford P, et al. Interpreting change scores for pain and functional status in low back pain: towards international consensus regarding minimal important change. *Spine (Phila Pa 1976)*. Jan 1 2008;33(1):90-94. PMID 18165753
24. Comstock BA, Sitlani CM, Jarvik JG, et al. Investigational vertebroplasty safety and efficacy trial (INVEST): patient-reported outcomes through 1 year. *Radiology*. Oct 2013;269(1):224-231. PMID 23696683
25. Firanesco CE, de Vries J, Lodder P, et al. Vertebroplasty versus sham procedure for painful acute osteoporotic vertebral compression fractures (VERTOS IV): randomised sham controlled clinical trial. *BMJ*. 2018 May 9;361:k1551. PMID: 29743284
26. Chen D, An ZQ, Song S, et al. Percutaneous vertebroplasty compared with conservative treatment in patients with chronic painful osteoporotic spinal fractures. *J Clin Neurosci*. Mar 2014;21(3):473-477. PMID 24315046
27. Farrokhi MR, Alibai E, Maghami Z. Randomized controlled trial of percutaneous vertebroplasty versus optimal medical management for the relief of pain and disability in acute osteoporotic vertebral compression fractures. *J Neurosurg Spine*. May 2011;14(5):561-569. PMID 21375382
28. Edidin AA, Ong KL, Lau E, et al. Mortality risk for operated and nonoperated vertebral fracture patients in the Medicare population. *J Bone Miner Res*. Jul 2011;26(7):1617-1626. PMID 21308780

29. Edidin AA, Ong KL, Lau E, et al. Morbidity and mortality after vertebral fractures: comparison of vertebral augmentation and nonoperative management in the Medicare population. *Spine (Phila Pa 1976)*. Aug 01 2015;40(15):1228-1241. PMID 26020845
30. Lin JH, Chien LN, Tsai WL, et al. Early vertebroplasty associated with a lower risk of mortality and respiratory failure in aged patients with painful vertebral compression fractures: a population-based cohort study in Taiwan. *Spine J*. Sep 2017;17(9):1310-1318. PMID 28483705
31. Clark W, Bird P, Gonski P, et al. Safety and efficacy of vertebroplasty for acute painful osteoporotic fractures (VAPOUR): a multicentre, randomised, double-blind, placebo-controlled trial. *Lancet*. Oct 01 2016;388(10052):1408-1416. PMID 27544377
32. Klazen CA, Lohle PN, de Vries J, et al. Vertebroplasty versus conservative treatment in acute osteoporotic vertebral compression fractures (Vertos II): an open-label randomised trial. *Lancet*. Sep 25 2010;376(9746):1085-1092. PMID 20701962
33. Yi X, Lu H, Tian F, et al. Recompression in new levels after percutaneous vertebroplasty and kyphoplasty compared with conservative treatment. *Arch Orthop Trauma Surg*. Jan 2014;134(1):21-30. PMID 24287674
34. Leali PT, Solla F, Maestretti G, et al. Safety and efficacy of vertebroplasty in the treatment of osteoporotic vertebral compression fractures: a prospective multicenter international randomized controlled study. *Clin Cases Miner Bone Metab*. Sep-Dec 2016;13(3):234-236. PMID 28228788
35. Yang EZ, Xu JG, Huang GZ, et al. Percutaneous Vertebroplasty versus conservative treatment in aged patients with acute osteoporotic vertebral compression fractures: a prospective randomized controlled clinical study. *Spine (Phila Pa 1976)*. Apr 2016;41(8):653-660. PMID 26630417
36. Lourie H. Spontaneous osteoporotic fracture of the sacrum. An unrecognized syndrome of the elderly. *JAMA*. Aug 13 1982;248(6):715-717. PMID 7097924
37. Frey ME, Depalma MJ, Cifu DX, et al. Percutaneous sacroplasty for osteoporotic sacral insufficiency fractures: a prospective, multicenter, observational pilot study. *Spine J*. Mar-Apr 2008;8(2):367-373. PMID 17981097
38. Kortman K, Ortiz O, Miller T, et al. Multicenter study to assess the efficacy and safety of sacroplasty in patients with osteoporotic sacral insufficiency fractures or pathologic sacral lesions. *J Neurointerv Surg*. Sep 1 2013;5(5):461-466. PMID 22684691
39. Frey ME, Warner C, Thomas SM, et al. Sacroplasty: a ten-year analysis of prospective patients treated with percutaneous sacroplasty: literature review and technical considerations. *Pain Physician*. Nov 2017;20(7):E1063- E1072. PMID 29149151
40. Dougherty RW, McDonald JS, Cho YW, et al. Percutaneous sacroplasty using CT guidance for pain palliation in sacral insufficiency fractures. *J Neurointerv Surg*. Jan 2014;6(1):57-60. PMID 23345629
41. Zaman FM, Frey M, Slipman CW. Sacral stress fractures. *Curr Sports Med Rep*. Feb 2006;5(1):37-43. PMID 16483515
42. Denis F, Davis S, Comfort T. Sacral fractures: an important problem. Retrospective analysis of 236 cases. *Clin Orthop Relat Res*. Feb 1988;227:67-81. PMID 3338224
43. Shah LM, Jennings JW, Kirsch CFE et al. ACR Appropriateness Criteria^(R) Management of Vertebral Compression Fractures. *J Am Coll Radiol*. 2018 Nov;15(11S). PMID 30392604
44. Barr JD, Jensen ME, Hirsch JA, et al. Position statement on percutaneous vertebral augmentation: a consensus statement developed by the Society of Interventional Radiology (SIR), American Association of Neurological Surgeons (AANS) and the Congress of Neurological Surgeons (CNS), American College of Radiology (ACR), American Society of Neuroradiology (ASNR), American Society of Spine Radiology (ASSR), Canadian Interventional Radiology

- Association (CIRA), and the Society of NeuroInterventional Surgery (SNIS). *J Vasc Interv Radiol*. 2014 Feb;25(2):171-81. PMID: 24325929.
45. Baerlocher MO, Saad WE, Dariushnia S, et al. Quality improvement guidelines for percutaneous vertebroplasty. *J Vasc Interv Radiol*. Feb 2014;25(2):165-170. PMID 24238815
 46. McGuire R. AAOS Clinical Practice Guideline: the Treatment of Symptomatic Osteoporotic Spinal Compression Fractures. *J Am Acad Orthop Surg*. 2011 Mar;19(3). PMID 21368100
 47. National Institute for Health and Care Excellence (NICE). Percutaneous vertebroplasty [IPG12]. 2003; <https://www.nice.org.uk/guidance/ipg12>. Accessed March 18, 2020.
 48. National Institute for Health and Care Excellence (NICE). Percutaneous vertebroplasty and percutaneous balloon kyphoplasty for treating osteoporotic vertebral compression fractures [TA279]. 2013; <https://www.nice.org.uk/guidance/ta279>. Accessed March 18, 2020.
 49. National Institute for Health and Care Excellence (NICE). Metastatic spinal cord compression in adults: risk assessment, diagnosis and management [CG75]. 2008; <https://www.nice.org.uk/guidance/cg75/chapter/1-guidance>. Accessed March 18, 2020.

Other References

1. Blue Cross and Blue Shield of Kansas National Consultant, Practicing Board Certified Orthopedic Surgeon (241), September 2008.
2. Blue Cross and Blue Shield of Kansas National Consultant, Practicing Board Certified Interventional Radiologist (470), December 2008.
3. Blue Cross and Blue Shield of Kansas Family Practice Liaison Committee CB, May 2009.
4. Blue Cross and Blue Shield of Kansas Orthopedic Liaison Committee CB, May 2009.
5. Blue Cross and Blue Shield of Kansas Radiology Liaison Committee CB, May 2009.
6. Blue Cross and Blue Shield of Kansas Radiology Liaison Committee, February 2013; July 2017; January 2018.
7. Blue Cross and Blue Shield of Kansas Orthopedic Liaison Committee, February 2013; August 2017.
8. Blue Cross and Blue Shield of Kansas Family Practice Liaison Committee, July 2013.