

Medical Policy



Title: Radiofrequency Ablation (RFA) of Miscellaneous Solid Tumors Excluding Liver Tumors

Related Policies:	<ul style="list-style-type: none"> ▪ <i>Radiofrequency Ablation of Primary or Metastatic Liver Tumors</i> ▪ <i>Cryoablation of Tumors Located in the Kidney, Lung, Breast, Pancreas, or Bone</i> ▪ <i>Stereotactic Radiosurgery and Stereotactic Body Radiotherapy</i>
-------------------	---

Professional / Institutional
Original Effective Date: January 1, 2006 / September 22, 2008
Latest Review Date: October 28, 2025
Current Effective Date: December 7, 2018

State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact Blue Cross and Blue Shield of Kansas Customer Service.

The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.

The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.

If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"> • With painful osteolytic bone metastases who have failed or are poor candidates for standard treatments 	Interventions of interest are: <ul style="list-style-type: none"> • Radiofrequency ablation 	Comparators of interest are: <ul style="list-style-type: none"> • Medical management • Radiotherapy 	Relevant outcomes include: <ul style="list-style-type: none"> • Symptoms • Change in disease status • Quality of life • Medication use • Treatment-related morbidity

Populations	Interventions	Comparators	Outcomes
Individuals: • With painful osteoid osteomas	Interventions of interest are: • Radiofrequency ablation	Comparators of interest are: • Medical management • Surgical excision • Core drill excision • Laser photocoagulation	Relevant outcomes include: • Symptoms • Change in disease status • Quality of life • Medication use • Treatment-related morbidity
Individuals: • With localized renal cell carcinoma no more than 4 cm in size	Interventions of interest are: • Radiofrequency ablation	Comparators of interest are: • Surgical excision	Relevant outcomes include: • Overall survival • Change in disease status • Quality of life • Treatment-related morbidity
Individuals: • With inoperable primary pulmonary tumors or nonpulmonary tumors metastatic to the lung	Interventions of interest are: • Radiofrequency ablation	Comparators of interest are: • Radiotherapy	Relevant outcomes include: • Overall survival • Change in disease status • Quality of life • Treatment-related morbidity
Individuals: • With breast tumors	Interventions of interest are: • Radiofrequency ablation	Comparators of interest are: • Radiotherapy • Surgical excision	Relevant outcomes include: • Overall survival • Change in disease status • Quality of life • Treatment-related morbidity
Individuals: • With benign thyroid tumors	Interventions of interest are: • Radiofrequency ablation	Comparators of interest are: • Radiotherapy • Surgical excision	Relevant outcomes include: • Symptoms • Change in disease status • Quality of life • Medication use • Treatment-related morbidity
Individuals: • With miscellaneous solid tumors (e.g., head and neck, thyroid cancer, pancreas)	Interventions of interest are: • Radiofrequency ablation	Comparators of interest are: • Radiotherapy • Surgical excision	Relevant outcomes include: • Overall survival • Change in disease status • Quality of life • Treatment-related morbidity

DESCRIPTION

In radiofrequency ablation (RFA), a probe is inserted into the center of a tumor; then, prong-shaped, non-insulated electrodes are projected into the tumor. Next, heat is generated locally by an alternating, high-frequency current that travels through the electrodes. The localized heat treats the tissue adjacent to the probe, resulting in a 3 cm to 5.5 cm sphere of dead tissue. The cells killed by RFA are not removed but are gradually replaced by fibrosis and scar tissue. If there is a local recurrence, it occurs at the edge and can sometimes be retreated. RFA may be performed percutaneously, laparoscopically, or as an open procedure.

OBJECTIVE

The objective of this evidence review is to determine whether the use of radiofrequency ablation improves the net health outcome in individuals with a range of tumors (including osteolytic bone metastases, osteoid osteomas, renal cell carcinoma, lung cancer, breast tumors, thyroid tumors, and others).

BACKGROUND**Radiofrequency Ablation**

Radiofrequency ablation (RFA) was initially developed to treat inoperable tumors of the liver. Recently, studies have reported on the use of RFA to treat other tumors. For some of these, RFA is being investigated as an alternative to surgery for operable tumors. Well-established local or systemic treatment alternatives are available for each of these malignancies. The hypothesized advantages of RFA for these cancers include improved local control and those common to any minimally invasive procedure (eg, preserving normal organ tissue, decreasing morbidity, decreasing length of hospitalization).

Goals of RFA may include (1) controlling local tumor growth and preventing recurrence; (2) palliating symptoms; and (3) extending survival duration for patients with certain tumors. The effective volume of RFA depends on the frequency and duration of applied current, local tissue characteristics, and probe configuration (eg, single vs. multiple tips). RFA can be performed as an open surgical procedure, laparoscopically or percutaneously, with ultrasound or computed tomography guidance.

Potential complications associated with RFA include those caused by heat damage to normal tissue adjacent to the tumor (eg, intestinal damage during RFA of kidney), structural damage along the probe track (eg, pneumothorax as a consequence of procedures on the lung), and secondary tumors (if cells seed during probe removal).

REGULATORY STATUS

The U.S. Food and Drug Administration (FDA) issued a statement in September 2008, concerning the regulatory status of RFA. The FDA has cleared RFA devices for the general indication of soft tissue cutting, coagulation, and ablation by thermal coagulation necrosis. Under this general indication, RFA can be used to ablate tumors, including lung tumors. Some RFA devices have

been cleared for additional specific treatment indications, including partial or complete ablation of nonresectable liver lesions and palliation of pain associated with metastatic lesions involving bone. The FDA has not cleared any RFA devices for the specific treatment indication of partial or complete ablation of lung tumors, citing lack of sufficient clinical data to establish safety and effectiveness for this purpose. The FDA has received reports of death and serious injuries associated with the use of RFA devices in the treatment of lung tumors.

POLICY**Osteolytic bone metastases**

- A. Radiofrequency ablation may be considered **medically necessary** to palliate pain in individuals with osteolytic bone metastases who have failed or are poor candidates for standard treatments such as radiation or opioids.

Osteoid osteomas

- B. Radiofrequency ablation may be considered **medically necessary** to treat osteoid osteomas that cannot be managed successfully with medical treatment.

Renal cell carcinoma

- C. Radiofrequency ablation may be considered **medically necessary** to treat localized renal cell carcinoma that is no more than 4 cm in size when either of the following criteria is met:
1. In order to preserve kidney function; **OR**
 2. The patient is not considered a surgical candidate.

Non-small-cell lung cancer

- D. Radiofrequency ablation may be considered **medically necessary** to treat an isolated peripheral non-small-cell lung cancer lesion that is no more than 3 cm in size when the following criteria are met:
1. Surgical resection or radiation treatment with curative intent is considered appropriate based on stage of disease; however, medical co-morbidity renders the individual unfit for those interventions; **AND**
 2. Tumor is located at least 1 cm from the trachea, main bronchi, esophagus, aorta, aortic arch branches, pulmonary artery, and the heart.

Nonpulmonary tumor(s) metastatic to the lung

- E. Radiofrequency ablation may be considered **medically necessary** to treat malignant nonpulmonary tumor(s) metastatic to the lung that are no more than 3 cm in size when the following criteria are met:
1. In order to preserve lung function when surgical resection or radiation treatment is likely to substantially worsen pulmonary status **OR** the individual is not considered a surgical candidate; **AND**
 2. There is no evidence of extrapulmonary metastases; **AND**
 3. The tumor is located at least 1 cm from the trachea, main bronchi, esophagus, aorta, aortic arch branches, pulmonary artery, and the heart. (See the Policy Guidelines Section for additional criteria)

- F. Radiofrequency ablation is considered **experimental / investigational** as a technique for ablation of:
1. Breast tumors;
 2. Lung cancer not meeting the criteria above;
 3. Renal cell cancer not meeting the criteria above;
 4. Osteoid osteomas that can be managed with medical treatment;
 5. Painful bony metastases as initial treatment; **AND**
 6. All other tumors outside the liver including, but not limited to, the head and neck, thyroid, adrenal gland, ovary, and pelvic/abdominal metastases of unspecified origin.

POLICY GUIDELINES

- A. The following additional criteria have been developed by clinical judgment / consensus and existing guidelines for the use of RFA in metastatic tumors to the lung and include:
1. No more than 3 tumors per lung should be ablated;
 2. Tumors should be amenable to complete ablation; **AND**
 3. Twelve months should elapse before a repeat ablation is considered.

Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

RATIONALE

This evidence review was created using searches of the PubMed database. The most recent literature update was performed through August 5, 2025.

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life (QOL), and ability to function - including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can

generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

OSTEOLYTIC BONE METASTASES

Clinical Context and Therapy Purpose

The purpose of radiofrequency ablation (RFA) in individuals who have painful osteolytic bone metastases who have failed or are poor candidates for standard treatments is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with painful osteolytic bone metastases who have failed or are poor candidates for standard treatments. After lung and liver, bone is the third most common metastatic site and is relatively frequent among individuals with primary malignancies of the breast, prostate, and lung. Bone metastases often cause osteolysis (bone breakdown), resulting in pain, fractures, decreased mobility, and reduced QOL.

Interventions

The therapy being considered is RFA.

Comparators

The following therapies and practices are options to manage painful osteolytic bone metastases: medical management (eg, chemotherapy) and radiotherapy. External-beam radiotherapy often is the initial palliative therapy for osteolytic bone metastases. However, pain from bone metastases is refractory to radiotherapy in 20% to 30% of individuals, while recurrent pain at previously irradiated sites may be ineligible for additional radiation due to risks of normal tissue damage. Other alternatives include hormonal therapy, radiopharmaceuticals (eg, strontium 89), and bisphosphonates. Less often, surgery or chemotherapy may be used for palliation, and intractable pain may require opioid medications.

Outcomes

The general outcomes of interest are overall survival (OS), reduction in pain and medication use, fractures, functional outcomes, and QOL.

Patients would be followed for several years given the impact of bone metastases on bone remodeling.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;

- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

Cohort Studies

Levy et al (2020) conducted a global, multicenter, nonrandomized, prospective postmarketing study to evaluate the effectiveness of RFA in patients with painful osteolytic bone metastases.¹ Between October 2017 and March 2019, 134 ablations were performed in 100 patients (68% vs. 32% of the cohort had a single vs. multiple sites treated, respectively). The most common tumor location was thoracic (44%) followed by lumbar (33%). Patient outcomes including pain, pain interference, and QOL were collected. Forty percent of the cohort did not participate through the 6-month follow-up, with 2 additional discontinuations after 6 months. The most common reason for discontinuation was death (30 patients), which were all classified as related to the underlying malignancy. The primary endpoint evaluated was pain improvement, from baseline to 3 months. At baseline, the mean score for worst pain (measured by Brief Pain Inventory) for the entire cohort was 8.2. After RFA, worst pain significantly improved, with mean scores decreasing to 5.6, 4.7, 3.9, 3.7, and 3.5 at 3 days, 1 week, 1 month, 3 months, and 6 months, respectively ($p < .0001$ for all visits). Immediate improvement in pain (≥ 2 -point change in worst pain at the treatment site[s] 3 days after RFA) was achieved by 59% of patients. Four adverse events were reported, of which 2 resulted in hospitalization for pneumonia and respiratory failure, respectively.

Case Series

Goetz et al (2004) reported on an international study conducted at 9 centers in which 43 patients with painful osteolytic bone metastases were treated palliatively with RFA.² The study's primary outcome measure was the Brief Pain Inventory-Short Form, a validated scale from 0 (no pain) to 10 (worst pain imaginable). Patient eligibility required baseline values of 4 or more from 2 or fewer painful sites. Thirty-nine (91%) of the patients had previously received opioids to control pain from the lesion(s) treated with RFA, and 32 (74%) had prior radiotherapy to the same lesion. The mean pain score at baseline was 7.9 (range, 4 to 10). At 4, 12, and 24 weeks after RFA, average pain scores decreased to 4.5, 3.0, and 1.4, respectively (all $p < .001$). Forty-one (95%) patients achieved clinically significant reductions in pain scores, prospectively defined as a decrease of 2 units from baseline. Investigators also reported statistically significant ($p = .01$) decreases in opioid use at weeks 8 (by 59%) and 12 (by 54%).

An earlier case series by Gronemeyer et al (2002) showed that palliative RFA provided significant pain relief in 9 (90%) of 10 patients with unresectable, osteolytic spine metastases who had no other treatment options.³ Pain was reduced by an average of 74%; back pain-related disability was reduced by an average of 27%. Neurologic function was preserved in 9 patients and improved in the other. In another small case series, Kojima et al (2006) assessed 24 patients with painful metastatic bone tumors who experienced pain-alleviating effects with RFA, which is consistent with other evidence.⁴

Section Summary: Osteolytic Bone Metastases

A prospective cohort study and case series have shown clinically significant reductions in pain relief (defined as a decrease of 2 units from baseline on the Brief Pain Inventory scale) or reductions in opioid use following treatment with RFA of osteolytic pain metastases in patients with no or limited treatment options. A multicenter, prospective study reported significant reductions in pain through the 6-month follow-up period, with 59% of patients achieving immediate improvement in pain within 3 days of RFA.

OSTEOID OSTEOMAS**Clinical Context and Therapy Purpose**

The purpose of RFA in individuals who have painful osteoid osteomas is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant populations of interest are individuals with painful osteoid osteomas.

Osteomas are the most common benign bone tumor, comprising 10% to 20% of benign and 2% to 3% of all bone tumors. They are typically seen in children and young adults, with most diagnosed in patients between 5 and 20 years of age. Osteomas are most common in the lower extremity (usually the long bones, mainly the femur) and less common in the spine. These tumors typically have a characteristic clinical presentation and radiologic appearance, with pain that is usually continuous and worse at night and commonly relieved by aspirin or other nonsteroidal anti-inflammatory drugs (NSAIDs). The natural history of the osteoid osteoma varies based on location, and although they rarely exceed 1.5 cm in diameter, may produce bone widening and deformation, limb length inequality, or angular deviations when near a growth plate. When located in the spine, these lesions may lead to painful scoliosis or torticollis. Sometimes they heal spontaneously after 3 to 7 years.

Interventions

The therapy being considered is RFA.

RFA of osteoid osteoma is done with a needle puncture, so no incision or sutures are needed; further, patients may immediately walk on the treated extremity and return to daily activities when the anesthetic effect wears off. The risk of recurrence with RFA of an osteoma is 5% to 10%, and recurrent tumors can be retreated with RFA. In general, RFA is not performed in many spinal osteomas because of possible thermal-related nerve damage.

Comparators

The following therapies and practices are options to treat osteoid osteomas: medical management, surgical excision, core drill excision, and laser photocoagulation.

Treatment options include medical management with NSAIDs, surgical excision (wide/en bloc excision or curetting), or the use of computed tomography (CT) or magnetic resonance imaging(MRI)-guided minimally invasive procedures including core drill excision, laser

photocoagulation, or RFA. For many years, complete surgical excision was the classic treatment of osteomas, usually performed in patients with pain despite medical management. However, a substantial incision may be necessary, with the removal of a considerable amount of bone (especially in the neck of the femur). This increases the need for bone grafting plus internal fixation (which often necessitates a second procedure to remove the metalwork). Other possible risks include avascular necrosis of the femoral head and postoperative pathologic fracture. In addition, surgical excision leads to a lengthier convalescence and postoperative immobilization. Anatomically inaccessible tumors may not be completely resectable and may recur.

Outcomes

The general outcomes of interest are reductions in pain and medication use, normal bone development, and postsurgical adverse events.

Patients would be followed through adolescence to ensure normal skeletal development.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

Systematic Reviews

Tordjman et al (2020) conducted a systematic review on CT-guided RFA for osteoid osteomas.⁵ The review included 69 studies (43 retrospective and 12 prospective studies; rest of study designs were not identifiable) comprising 3023 patients. The weighted overall failure rate was 8.3% for the entire cohort. When studies were analyzed by time period conducted, the failure rate was significantly lower in studies conducted between 2011 and 2019 compared to those conducted between 2002 and 2010 (7% vs. 14%, $p=.004$). The complication rate for the entire cohort was 3%, with skin burns (0.7%) and infections (0.5%) as the most commonly reported.

Lanza et al (2014) reported on a systematic review of various ablative techniques for osteoid osteomas.⁶ Included in the review were 23 articles on RFA, 3 on interstitial laser ablation, and 1 with a combination of ablation techniques, totaling 27 articles (N=1772). The mean technical success was 100% and clinical success, defined as being pain-free, ranged from 94% to 98%, depending on the length of follow-up. Complications occurred in 2% of patients and included skin or muscle burn in 9 patients, 4 infections, nerve lesions or tool breakage in 3 patients each, delayed skin healing, hematoma, and failure to reach target temperature in 2 patients each, and fracture, pulmonary aspiration, thrombophlebitis, and cardiac arrest in 1 patient each. Eighty-six patients had tumor recurrence.

Retrospective Studies

In their retrospective study of the efficacy and complications of CT-guided RFA of spinal osteoid osteoma, Albisinni et al (2017) concluded that CT-guided RFA is as effective as first-line therapy for the disease.⁷ After RFA, clinical symptoms were evaluated at 3, 6, and 12 months, with a final evaluation at the end of the study. Results showed complete regression of osteoid osteoma symptoms in 57 (93.4%) of 61 ($p=.001$) patients observed between 2002 and 2012. Study limitations included the retrospective design and focus on a single treatment.

Lassalle et al (2017) conducted a single-center, retrospective analysis of long-term outcomes for CT-guided RFA in 126 patients with suspected osteoid osteoma.⁸ The study was conducted from 2008 to 2015. Phone evaluations were performed. The overall success rate was 94.3% among the 88 patients who participated in the follow-up calls. The study was limited by its retrospective design, imprecision of patients' memory over follow-up, the lack of clinical and imaging follow-up, and an inability to perform multivariate statistical analysis of factors associated with treatment failure.

Rimondi et al (2012) reported on a retrospective study of 557 patients treated with CT-guided RFA as primary treatment for nonspinal osteoid osteomas.⁹ All patients were followed for a mean of 3.5 years (range, 0.5 to 9 years). Pain relief occurred in all 557 patients within the first week after RFA and continued in 533 (96%) patients who remained asymptomatic through their last follow-up. Pain recurrence occurred in 24 (4%) patients. Complications occurred in 5 patients and included thrombophlebitis, skin burn, broken electrode, and 2 procedures in which the RFA generator failed to reach maximum temperature.

Sahin et al (2019) conducted a single-center, retrospective study that evaluated clinical pain symptoms to demonstrate the rapid relief of pain symptoms after CT-guided RFA for osteoid osteomas.¹⁰ A total of 116 patients were included, and the efficacy success rate in the study was 98%. All patients reported immediate pain relief following the procedure, with scores of 0 or 1 on a 10-point visual analog pain scale within 24 hours. The mean duration of follow-up was 23 months, and pain relapse was reported in 2 of 108 patients available for follow-up. Seven minor complications were reported after the procedure with superficial skin burns as the most common complication ($n=4$).

Case Series

An observational study by Knudsen et al (2015) evaluated long-term clinical outcomes after CT-guided RFA in patients diagnosed with osteoid osteoma located in the upper and lower extremities.¹¹ The study population included 52 patients with a typical clinical history and radiologically confirmed osteoid osteoma who received CT-guided RFA treatment from 1998 to 2014 at a Danish university hospital. The clinical outcome was evaluated based on patient-reported outcome measures and medical record review. The response rate was 52 (87%) of 60. After 1 RFA treatment, 46 (88%) of 52 patients experienced pain relief, and 51 (98%) of 52 patients had pain relief after repeat RFA. One patient underwent open resection after RFA. No major complications were reported; 4 patients reported minor complications including small skin burn, minor skin infection, and hypoesthesia at the needle entry point. In all, 50 (96%) of 52 patients were reported to be "very satisfied" with the RFA treatment.

Rosenthal et al (2003) reported their experience over an 11-year period with 271 RFA procedures for osteoid osteomas in 263 patients.¹² The short-term outcome was evaluated to detect procedure-related problems; by this definition, all procedures were considered technically successful. Long-term clinical success data (defined as being free of pain without additional procedures) were available in 126 patients, with complete clinical success observed in 89%. For procedures performed as the initial treatment, the success rate was 91%.

Section Summary: Osteoid Osteomas

Numerous retrospective studies and case series, and systematic reviews of observational data have evaluated RFA for the treatment of painful osteoid osteomas. In a systematic review of thermal ablation techniques, clinical success (pain-free) was achieved in 94% to 98% of patients. Results have indicated that most patients (89% to 96%) remained pain-free at longer-term follow-up. Another systematic review reported similar success rates noting an average 8.3% failure rate among patients receiving CT-guided RFA.

LOCALIZED RENAL CELL CARCINOMA

Clinical Context and Therapy Purpose

Radical nephrectomy remains the principal treatment of renal cell carcinoma (RCC); however, partial nephrectomy (PN) or nephron-sparing surgery has been shown to be as effective as radical nephrectomy, with comparable long-term recurrence-free survival rates, in a select group of individuals. Alternative therapy such as RFA is of interest in individuals with small renal tumors when preservation of renal function is necessary (eg, in patients with marginal renal function, a solitary kidney, bilateral tumors) and in individuals with comorbidities that would render them unfit for surgery. Another consideration would be in individuals at high risk of developing additional renal cancers (eg, von Hippel-Lindau disease).

The purpose of RFA in individuals who have localized RCC no more than 4 cm in size is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with localized RCC no more than 4 cm in size. Small renal masses, defined as 4 cm or less, are common findings on diagnostic imaging of the abdomen pelvis. Some of these masses are assessed to be suspicious for malignancy or have been identified by biopsy as a localized RCC. Tumors can be further categorized according to international tumor, nodes, metastasis (TNM) staging where cT1a is a clinically diagnosed tumor ≤ 4 cm that is confined to the kidney without any nodal involvement.

Interventions

The therapy being considered is RFA.

Comparators

The following practice is currently being used to treat localized RCC: surgical excision; either total nephrectomy or PN.

Outcomes

The general outcomes of interest are recurrence rates and a reduction in rates of renal failure.

Individuals should be followed for at least 10 years to monitor for tumor recurrence.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

Systematic Reviews

In a systematic review and meta-analysis, Huang et al (2025) compared stereotactic body radiotherapy (SBRT) with RFA, microwave ablation, and cryoablation outcomes for primary localized renal cell carcinoma.¹³ The review identified 133 studies that included 8,910 total patients, of whom 612 received SBRT, 2,503 received RFA, 2,069 received microwave ablation, and 3,726 received cryoablation. Overall 1-, 2-, and 5-year local control rates were highest for SBRT (99%, 97%, and 95%) compared with RFA (96%, 95%, 92%), microwave ablation (97%, 95%, 86%), and cryoablation (95%, 94%, 90%). Subgroup analysis by tumor size showed comparable outcomes across modalities for tumors <4 cm, but for tumors ≥4 cm, SBRT achieved the highest local control at all time points. Regarding survival, cancer-specific survival at 1 year was 100% and consistent across all treatment groups. By 5 years, slight variations emerged, with SBRT showing the lowest rate at 95% (and RFA the highest at 100%). The heterogeneity across studies was moderate to high for most treatment modalities, which limits the strength of the conclusions.

In a systematic review and meta-analysis, Yanagisawa et al (2022) compared differential clinical outcomes of patients treated with PN versus those treated with ablation techniques, including RFA, cryoablation, and microwave ablation, for cT1b and cT1a renal tumors.¹⁴ They identified 27 studies with 13,996 total patients who received either PN or ablation for treatment of their tumors. Investigators found that in both cT1a and cT1b renal tumors, there were no differences in the percent decline of estimated glomerular filtration rates (eGFR) or in the overall complication rates between PN and ablation therapy. There was also no difference in cancer mortality rates between PN and ablation in patients with either cT1a or cT1b tumors. However, compared to ablation, PN was associated with a lower risk of local recurrence in patients with either tumor type. There was significant heterogeneity across studies, which limits conclusions.

In their systematic review and meta-analysis, Uhlig et al (2019) compared oncologic, perioperative, and functional outcomes for PN with outcomes for various ablative techniques, including RFA and others, for small renal masses (mean diameter=2.53 to 2.84 cm).¹⁵ They identified 47 moderate-quality studies, mostly retrospective, published from 2005 to 2017, with a

total of 24,077 patients. Of these patients, 15,238 received PN and 1877 received RFA. The network meta-analysis used PN as the reference point. The overall results indicated that PN had better OS and local control over ablative techniques, but it was not significantly better for cancer-related mortality. In addition, ablation had fewer complications and better renal function outcomes. Across the studies included, patients treated by PN tended to be younger with less comorbidities compared with patients receiving thermal ablation—a consideration when assessing the outcomes for survival and local control.

In a systematic review and meta-analysis, Katsanos et al (2014) reviewed 1 RCT and 5 cohort studies (N=587) assessing thermal ablation (RFA or microwave) or nephrectomy for small renal tumors (size, 2.5 cm).¹⁶ The local recurrence rate was 3.6% in both groups (relative risk, 0.92; 95% confidence interval [CI], 0.4 to 2.14; p=.79). Disease-free survival was also similar in both groups up to 5 years (hazard ratio [HR], 1.04; 95% CI, 0.48 to 2.24; p=.92). However, the overall complication rate was significantly lower in the patients undergoing ablation (7.4%) versus nephrectomy (11.1%; pooled relative risk, 0.55; 95% CI, 0.31 to 0.97; p=.04).

El Dib et al (2012) conducted a meta-analysis evaluating RFA and cryoablation for small renal masses.¹⁷ Selected were 11 RFA case series (426 patients) and 20 cryoablation case series (457 patients) published through January 2011. The mean tumor size was 2.7 cm (range, 2 to 4.3 cm) in the RFA group and 2.5 cm (range, 2 to 4.2 cm) in the cryoablation group. Mean follow-up times for the RFA and cryoablation groups were 18.1 and 17.9 months, respectively. Clinical efficacy, defined as cancer-specific survival rate, radiographic success, no evidence of local tumor progression, or distant metastases, did not differ significantly between groups. The pooled proportion of clinical efficacy for RFA was 90% (95% CI, 86% to 93%) and 89% (95% CI, 83% to 94%) for cryoablation.

Table 1 summarizes the characteristics of all systematic reviews. Table 2 contains the results of the largest and most recent of the reviews (Uhlig et al [2019] and Yanagisawa et al [2022]).

Table 1. Characteristics of Meta-Analyses Assessing Radiofrequency Ablation for Renal Masses

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Huang et al (2025) ¹³	2000-2024	133	Patients who had received RFA, SBRT, CRA, or MWA for primary and localised renal cell carcinoma	8,910 (n/a)	Prospective, observational, RCTs	Median ranged from 24.5 to 28.7 months
Yanagisawa (2022) ¹⁴	2005-2021	27	Patients who underwent PN, RFA, CRA, or MWA for small renal tumors (cT1a or cT1b).	13,996 (18 to 8818)	Prospective, retrospective, 1 RCT	14 months to 6 years
Uhlig (2019) ¹⁵	2006-2017	47	Patients who had received PN, RFA, CRA, or MWA for small renal masses.	24,077 (18 to 1803)	Prospective, retrospective, 1 RCT	3 to 82 months
Katsanos (2014) ¹⁶	2007-2012	6	Patients with small renal tumors receiving RFA or nephrectomy.	587 (69 to 150)	1 RCT, 5 cohort	Up to 6 years
El Dib (2012) ¹⁷	2000-2008	31	Patients who had received RFA or CRA for renal tumors, regardless of size.	957 (n/a)	Case series	7 to 45.7 months

CRA: cryoablation; MWA: microwave ablation; n/a: data not available; PN: partial nephrectomy; RCT: randomized controlled trial; RFA: radiofrequency ablation; SBRT: stereotactic body radiotherapy.

The results table below does not include Katsanos et al (2014) because of complete study overlap with Uhlig et al (2019). El Dib et al (2012) and Huang et al (2025) are not included because the comparator in the studies selected were various ablative strategies, not surgery. Table A1 in the Appendix compares the trials included in these systematic reviews and meta-analyses.

Table 2. Results of Select Meta-Analyses Assessing Radiofrequency Ablation for Renal Masses

Study	Cancer-Specific Mortality		Local Recurrence		Complications		Renal Function Decline, MD in eGFR	
	<i>RFA (IRR)</i>	<i>PN (IRR)</i>	<i>RFA (IRR)</i>	<i>PN (IRR)</i>	<i>RFA (OR)</i>	<i>PN (OR)</i>	<i>RFA</i>	<i>PN</i>
Uhlig (2019) ¹⁵ ,	2.03	1.00	1.79	1.00	0.89	1.00	6.49	0.00
95% CI	0.81 to 5.08		1.16 to 2.76		0.59 to 1.33		2.87 to 10.10	
	<i>RFA (events)</i>	<i>PN (events)</i>	<i>RFA (events)</i>	<i>PN (events)</i>	<i>RFA (events)</i>	<i>PN (events)</i>	<i>RFA (total decline)</i>	<i>PN (total decline)</i>
Yanagisawa (2022) ¹⁴ ,	cT1a: 27 cT1b: 8	cT1a: 113 cT1b: 18	cT1a: 64 cT1b: 32	cT1a: 59 cT1b: 34	cT1a: 126 cT1b: 50	cT1a: 204 cT1b: 62	cT1a: 176 cT1b: 154	cT1a: 217 cT1b: 184
Pooled RR (95% CI)	cT1a: 0.87 (0.57 to 1.31) cT1b: 0.80 (0.32 to 1.98)		cT1a: 0.43 (0.28 to 0.66) cT1b: 0.41 (0.23 to 0.75)		cT1a: 1.34 (0.90 to 2.00) cT1b: 1.08 (0.76 to 1.53)		cT1a: MD, 2.42 (-0.06 to 4.89) cT1b: MD, 0.73 (-3.76 to 5.23)	
<i>I</i> ² (p-value)	cT1a: 0% (.62) cT1b: 0% (.76)		cT1a: 20% (.23) cT1b: 30% (.20)		cT1a: 63% (.003) cT1b: 22% (.26)		cT1a: 83% (.0004) cT1b: 0% (.71)	

CI: confidence interval; eGFR: estimated glomerular filtration rate; IRR: incidence rate ratio; MD: mean difference; OR: odds ratio; PN: partial nephrectomy; RFA: radiofrequency ablation; RR: risk ratio.

Randomized Controlled Trials

In an RCT, Liu et al (2016) analyzed the safety and efficacy of the operative effects of percutaneous RFA in early-stage RCC versus retroperitoneoscopic radical operation of RCC.¹⁸ There were 35 women and 28 men included; race and ethnicity of participants were not described. The observation group was treated with percutaneous RFA and the control group with a radical retroperitoneoscopy. A total of 76 clinically confirmed diagnosed cases, from January 2011 to January 2013, with RCC, were randomized to the observation (n=41) or the control (n=35) groups. Operation time, blood loss during operation, length of stay, and incidence complications were lower in the control group (p<.05). Total efficacy, tumor-free survival times, and survival rates did not differ statistically between groups (p>.05); however, percutaneous RFA reduced postoperative recovery time and was associated with fewer complications. Trial limitations included small sample size and the brief duration of follow-up.

Retrospective Studies

Relevant studies not included in the above systematic reviews and meta-analyses are described below.

Marshall et al (2020) conducted a single-center, retrospective evaluation in 100 patients with 125 RCCs who received percutaneous RFA between 2004 and 2015.¹⁹ Median follow-up in the study was 62.8 months. Five-year overall, cancer-specific, and local progression-free survival were 75%, 92%, and 92%, respectively. Ten-year overall, cancer-specific, and local progression-free survival were 32%, 86%, and 92%, respectively. The rate of local tumor progression was higher in patients with tumors >4 cm compared to those with tumors ≤4 cm, but the difference was not statistically significant (6% vs. 13%, $p=.466$). The study also noted no significant changes in eGFR from baseline to 2 to 3 years post-procedure (65.2 vs. 62.1 mL/min/1.73 m²; $p=.443$). The overall complication rate in the study was 9%. Limitations of the study include its retrospective design, lack of a control group, and selection bias where patients selected for RFA over surgical resection likely had worse baseline comorbidity status, which may have negatively impacted OS rates.

Dai et al (2017) conducted a retrospective evaluation of 30 patients with 31 central renal tumors who underwent percutaneous RFA between 2005 and 2010 to assess the clinical efficacy and safety of image-guided percutaneous RFA of central RCC with adjunctive pyeloperfusion.²⁰ OS was 96.0% (95% CI, 88.4% to 100.0%) and progression-free survival at 5 years was 80.9% (95% CI, 65.8% to 95.9%). The investigators found that complications were significantly higher for tumors located within 5 mm of the renal pelvis or 0 mm of a major calyx (28.6% vs. 4.0%; $p<.05$) and major complications occurred in 5 (12.8%) of 39 RFA sessions. They concluded that image-guided percutaneous RFA combined with pyeloperfusion had satisfactory clinical efficacy in the treatment of renal tumor but may be associated with significant major complications. The retrospective design and the small sample base are limitations to this analysis.

Over 10 years, Dvorak et al (2017) retrospectively evaluated the technical success as well as mid-term and long-term efficacy and safety of RFA and microwave ablation with guided CT in 64 patients with small, non-central renal tumors.²¹ Ninety-one ablation procedures were performed on 68 tumors, 12 to 60 mm in size. Treatment was successful in 50 (73.5%) tumors; a second procedure was successful in 13 (19.1%) cases; and for the 5 largest tumors (range, 45 to 60 mm; 7.4%), a third treatment was required. Investigators concluded that percutaneous ablation is safe and effective in treating small, non-central renal tumors of the T1a group. The retrospective study design is the major limitation of this study.

A publication by Iannuccilli et al (2016) reported a mean 34.1 months follow-up (range, 1 to 131 months) of RFA with intent to cure in 203 patients with renal tumors.²² Patients referred for RFA were at high risk or had refused surgery. Smaller tumors were treated with a single electrode with a 2 or 3 cm active tip. Larger tumors were treated with a cluster electrode with 3 active tips. Patients were assessed annually for the appearance of residual tumor at the treatment site, and 26 (13%) had residual disease. Treatment effectiveness was 87% during follow-up. The likelihood of recurrence was increased for tumors 3.5 cm or larger, clear cell subtype, and treatment temperature of 70° or less. All-cause mortality increased with increasing tumor size. The median survival was 7 years for patients with tumors less than 4 cm, with 80% survival at 5

years. Major complications, including urinary stricture or urine leak, occurred in 8 (3.9%) treatments.

Section Summary: Localized Renal Cell Carcinoma

The evidence on RFA for small renal tumors (≤ 4 cm) includes an RCT, meta-analyses, retrospective and cohort studies, and case series, that have compared RFA with nephrectomy or cryoablation. A 2014 meta-analysis that included 1 RCT and 5 cohort studies found that RFA was as effective as nephrectomy for small renal tumors, with a reduction in complications. Another, more recent, meta-analysis (2019) found that PN was superior to ablative techniques (the study included RFA but also cryoablation and microwave ablation) in overall mortality and local recurrence but not in cancer-specific mortality. It also found fewer complications and improved renal function with ablation. A meta-analysis from 2022 found that PN was superior to ablation (RFA, cryoablation, and microwave ablation) in local recurrence. Overall complications, decline in renal function, and cancer-specific mortality rates did not differ between ablation and PN. A 2025 meta-analysis found that RFA achieved high local control and cancer-specific survival rates comparable to other ablative therapies, although SBRT showed superior local control for larger tumors. The correlation between tumor size and RFA efficacy has been demonstrated by a large case series with a mean 34-month follow-up; it found that residual disease and mortality increased with tumors over 4 cm. Long-term follow-up in one single-center study found that RFA resulted in similar cancer-specific survival outcomes as PN in patients with cT1a renal tumors.

PRIMARY PULMONARY AND NONPULMONARY TUMORS

Clinical Context and Therapy Purpose

Surgery is the current treatment of choice in individuals with stage I primary non-small-cell lung cancer (NSCLC; stage I includes Ia [T1N0M0] and Ib [T2N0M0]). Approximately 20% of patients present with stage I disease, although this number is expected to increase as a result of screening programs, advances in imaging modalities and widespread use of CT scans for other indications. Postsurgical recurrence rates of stage I NSCLC have been reported as between 20% and 30%, with most occurring at distant sites; locoregional recurrences occur in approximately 12%. Large differences in survival outcomes are observed after surgery in stage I disease, with 5-year OS rates ranging from 77% for small T1 tumors to 35% for large T2 tumors. Untreated, stage I NSCLC has a 5-year OS rate range from 6% to 14%.

Individuals with early-stage NSCLC who are not surgical candidates may be candidates for radiotherapy with curative intent. In 2 large, retrospective, radiotherapy series, patients with the inoperable disease treated with definitive radiotherapy achieved 5-year survival rates of 10% and 27%. In both studies, patients with T1N0 tumors had better 5-year survival rates of 60% and 32%, respectively.

Stereotactic body radiotherapy has gained more widespread use as a treatment option because it is a high-precision mode of therapy that delivers very high doses of radiation. Two- to 3-year local control rates of stage I NSCLC with stereotactic body radiotherapy have ranged from 80% to 95%. Stereotactic body radiotherapy has been investigated in individuals unfit to undergo surgery, with survival rates similar to surgical outcomes.

RFA also is being investigated in individuals with small primary lung cancers or lung metastases who are deemed medically inoperable.

The purpose of RFA in individuals who have inoperable primary pulmonary tumors or nonpulmonary tumors metastatic to the lung is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with inoperable primary pulmonary tumors or nonpulmonary tumors metastatic to the lung.

Interventions

The therapy being considered is RFA.

Comparators

The following practice is currently being used to treat primary pulmonary tumors or nonpulmonary tumors metastatic to the lung: radiotherapy.

Outcomes

The general outcomes of interest are OS, tumor recurrence, and treatment-related adverse events (eg, pneumothorax). Individuals would be followed for at least 5 years.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

Systematic Reviews

In a systematic review of RFA, surgery, and stereotactic body radiotherapy for colorectal cancer (CRC) lung metastases, Schlijper et al (2014) did not identify any randomized trials, and evidence was insufficient to draw conclusions on the comparative effectiveness of these therapies.²³

In a comparative effectiveness review conducted for the Agency for Healthcare Research and Quality, Ratko et al (2013) assessed local nonsurgical therapies for stage I NSCLC.²⁴ In this review, no comparative RFA studies were identified. Reviewers found that available evidence was insufficient to draw conclusions on the comparative effectiveness of local nonsurgical therapies for NSCLC, including RFA.

In a review of 16 studies, Bilal et al (2012) compared RFA with stereotactic ablative radiotherapy in patients with inoperable early-stage NSCLC.²⁵ Reviewers found that OS rates for RFA and stereotactic ablative radiotherapy were similar in patients at 1 year (68.2% to 95% vs. 81% to 85.7%) and 3 years (36% to 87.5% vs. 42.7% to 56%), all respectively. However, survival rates at 5 years were lower with RFA (20.1% to 27%) than with stereotactic body radiotherapy (47%). These findings were drawn from comparisons of results from uncontrolled case series and retrospective reviews.

In an evidence-based review by Chan et al (2011), 46 studies on RFA for lung tumors were evaluated, which included 2905 ablations in 1584 patients with a mean tumor size of 2.8 cm.²⁶ Twenty-four studies reported rates of local recurrence, which occurred in 282 (12.2%) cases at a mean follow-up of 13 months (range, 3 to 45 months). Primary lung cancer rates of local recurrence did not differ significantly (22.2%) from metastases (18.1%). Twenty-one studies reported mean OS rates of 59.4% at a mean follow-up of 17.7 months. The mean cancer-specific survival rate was 82.6%, at a mean follow-up of 17.4 months. The mean overall morbidity was 24.6% and most commonly included pneumothorax (28.3%), pleural effusion (14.8%), and pain (14.1%). Mortality related to the RFA procedure was 0.21%, overall.

Prospective Studies

A relevant study not included in the above systematic reviews and meta-analyses are described below.

Hasegawa et al (2020) conducted a prospective, single-arm, multicenter study to evaluate the efficacy of RFA in patients with surgically resectable CRC lung metastases measuring 3 cm or smaller.²⁷ A total of 70 patients with CRC and 100 lung metastases were enrolled. All tumors were considered technically resectable, but not all patients were clinically able to undergo surgery. A total of 85 initial RFA sessions were performed for 100 target lung metastases. The 3-year OS rate after RFA was 84%. Primary and secondary technical success rates for RFA were 96% and 100%, respectively. Over a mean follow-up of 57 ± 32 months, local tumor progression was found in 6 patients (9%) at 6 to 19 months after the initial RFA. The 3-year progression-free survival rate was 41%. Grade 2 pneumothorax occurred after 18 of the 88 RFA sessions. The study is limited by its lack of a comparator arm.

Section Summary: Primary Pulmonary and Nonpulmonary Tumors

The evidence on RFA for primary NSCLC and nonpulmonary tumors metastatic to the lung includes prospective and observational studies and systematic reviews of those studies. No RCTs identified compared treatment approaches. For surgically resectable CRC lung metastases a multicenter study found that RFA for tumors less than 3 cm can lead to a 3-year OS rate of 84%. Two-year survival has been reported to range from 41% to 75% in case series. Survival at 1 and 2 years appears to be similar, following treatment with RFA or stereotactic ablative radiotherapy in patients with inoperable lung tumors. Survival rates at 5 years were lower with RFA (20.1% to 27%) than with stereotactic ablative radiotherapy (47%), but this finding was drawn from comparisons of uncontrolled case series and retrospective reviews. Prospective comparison in an RCT would permit greater certainty for this finding, but the studies are consistent with some effect of RFA on lung tumors.

BREAST TUMORS

Clinical Context and Therapy Purpose

The treatment of small cancers of the breast has evolved from total mastectomy to more conservative treatment options such as lumpectomy, with more acceptable cosmetic outcomes and preservation of the breast. The selection of the surgical approach balances the individual's desire for breast conservation and the need for tumor-free margins in resected tissue. Minimally invasive nonsurgical techniques such as RFA are appealing if they can produce local control and survival equivalent to breast-conserving surgical alternatives. Nonsurgical ablative techniques pose difficulties such as the inability to determine tumor size, complete tumor cell death, and local recurrence. Additionally, RFA can burn the skin and cause damage to muscle, possibly limiting use in patients with tumors near the skin or chest wall.

The purpose of RFA in individuals who have breast tumors is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with breast tumors.

Interventions

The therapy being considered is RFA .

Comparators

The following practices are currently being used as treatment options for small cancers of the breast cancer: radiotherapy and surgical excision.

Outcomes

The general outcomes of interest are tumor recurrence, reduction in medication, and treatment-related adverse events.

Patients would be followed for up to 5 years.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

Systematic Reviews

Xia et al (2021) conducted a systematic review and meta-analysis of studies assessing RFA in patients with breast cancer and tumors that were 2 cm or smaller.²⁸ The primary endpoints of interest were technical success rate, complete ablation rate, and rate of complications. A total of 17 studies were identified, which accounted for 399 patients (401 lesions). Technical success rates ranged from 86.67% to 100% in the included studies; the pooled technical success rate was 99% (95% CI, 98% to 100%). After RFA, the majority of patients underwent surgical tumor excision (65.74%, 261/397). The pooled complete ablation rate was 98% (95% CI, 97% to 100%). The complication rate in the entire cohort was 6.8%; the most common complications were skin burns (2%), breast inflammation (1.5%), and infections (1%). The pooled complications rate was 2% (95% CI, 1% to 4%). Local recurrence was reported in 10 studies (232 cases); there was no local recurrence reported after a median follow-up of 27 months in these patients. The authors noted that prospective studies evaluating the use of RFA alone are needed to validate the place in therapy.

Peek et al (2017) conducted a systematic review and meta-analysis of all studies evaluating the role of ablative techniques in the treatment of breast cancer published between 1994 and 2016.²⁹ Selection criteria included at least 10 patients with breast cancer treated with RFA, high-intensity ultrasound, or cryo-, laser, or microwave ablation; 63 studies (N=1608) were identified through PubMed and PubMed library databases. Fifty studies reported complete ablation, and RFA had the highest rate of complete ablation (87.1% [491/564]) as well as the shortest treatment time (15.6 minutes). A major limitation of this systematic review was the authors' inability to perform a comparative meta-analysis due to the inclusion of only 4 RCTs and 1 retrospective analysis that compared 2 or more techniques. There was also considerable heterogeneity across included studies.

Zhao and Wu (2010) conducted a systematic review of 38 studies on ablation techniques for breast cancer treatment published from 1994 to 2009.³⁰ Nine studies focused on RFA. Reviewers included small tumors ranging in size from 0.5 to 7 cm. Tumor resection was performed immediately after ablation or up to 4 weeks after RFA. Complete coagulation necrosis rates of 76% to 100% were reported. The results suggested RFA for breast cancer tumors is feasible, but further studies with longer follow-up on survival, tumor recurrence, and cosmetic outcomes would be needed to establish clinical efficacy.

In another review, Soukup et al (2010) examined 17 studies on RFA for the treatment of breast tumors and found RFA is feasible.³¹ Even though few adverse events and complications occurred with breast RFA, incomplete tumor ablation remains a concern.

Clinical Studies

Relevant studies not included in the above systematic reviews and meta-analyses are described below.

Retrospectively, Ito et al (2018) studied the safety and efficacy of percutaneous RFA of breast carcinomas in 386 patients from 10 institutions treated with RFA between 2003 and 2009.³² Race and ethnicity of participants were not described. Patients were followed for a median of 50 months, and ipsilateral breast tumor recurrence was more frequent in patients with initial tumor sizes of 2 cm or more (10% [3/30]) than those with initial tumors 2 cm or less (2.3% [8/355]);

p=.015). Ipsilateral breast tumor recurrence rates 5 years after RFA were 97%, 94%, and 87% in patients with initial tumor sizes of 1 cm or less, 1.1 to 2.0 cm, and greater than 2 cm, respectively. The authors concluded that RFA was safe for tumors of 2 cm or less. The retrospective design and lack of data on ipsilateral breast tumor recurrence for different types of chemotherapy and endocrine therapy and analyses to ascertain whether adjuvant chemotherapy or endocrine therapy influenced outcomes are the limitations of this study.

The efficacy and safety of using ultrasound-guided RFA for multiple breast fibroadenoma as an alternative to surgical resection were retrospectively analyzed by Li et al (2016).³³ From 2014 to 2016, 65 patients with 256 nodules were treated with ultrasound-guided RFA, and complete ablation was achieved for 251 nodules (98.04%) after the first month of treatment; after the first and third months, tumor volume overall was reduced by 39.06% and 75.99%, respectively. The study reported minimal to no complications such as skin burns, hematoma, or nipple discharge. The retrospective design and short follow-up time limited the conclusions drawn from this study. Race and ethnicity of participants were also not described.

Wilson et al (2012) reported on 73 patients with invasive breast cancer who had a lumpectomy followed immediately by RFA to the lumpectomy bed.³⁴ The average breast tumor size was 1.0 cm (range, 0.2 to 2.6 cm) and follow-up averaged 51 months. Disease-free survival was 100%, 92%, and 86% at 1, 3, and 5 years, respectively. One patient had tumor recurrence within 5 cm of the lumpectomy site, and 3 patients had ipsilateral breast recurrences.

Section Summary: Breast Tumors

Systematic reviews, retrospective studies, and observational studies have reported varied and incomplete ablation rates as well as concerns about post-ablation tumor cell viability. Long-term improvements in health outcomes have not been demonstrated. Additionally, available studies have not compared RFA with conventional breast-conserving procedures. For small breast tumors, further prospective study, with long-term follow-up, is needed to determine whether RFA can provide local control and survival rates compared with conventional breast-conserving treatment.

BENIGN THYROID NODULES

Clinical Context and Therapy Purpose

Surgical resection is the primary treatment choice for medically unresponsive, symptomatic benign thyroid tumors and thyroid carcinomas. However, techniques for ablation of thyroid tumors (eg, RFA, microwave ablation) are being investigated.

The purpose of RFA in individuals who have benign thyroid tumors is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with large or symptomatic benign thyroid tumors. Individuals with a benign cytology diagnosis or those very unlikely to be malignant (eg, purely cystic nodule) should undergo surveillance with the frequency determined by the level of

suspicion for a missed malignancy.³⁵ Medical or surgical intervention is considered if the nodules are large (>4 cm), causing compressive or structural symptoms, or if there is clinical concern.

Interventions

The therapy being considered is RFA .

Comparators

The following practices are currently being used to treat large or symptomatic benign thyroid tumors in the United States: percutaneous ethanol injection (PEI) and surgical excision.

Outcomes

The general outcomes of interest are a reduction in nodule volume, hyper- and hypothyroidism, and treatment-related adverse events (eg, voice changes).

Patients would be followed for at least 5 years.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE

Systematic Reviews

Xu et al (2024) evaluated the efficacy of thermal ablation (RFA, microwave, and laser ablation) for the treatment of benign thyroid nodules.³⁶ The analysis was limited to studies that had longer-term follow-up (approximately 5 years). A total of 5 studies (N=939) with 3 studies (n=483) specific to RFA were included. A total of 137 patients had local nodule recurrence at a median follow-up of 59.25 months.

Cho et al (2020) evaluated the efficacy of thermal ablation (RFA and laser ablation) for the treatment of benign thyroid nodules.³⁷ The analysis demonstrated long-term maintenance (up to 36 months) of volume reduction. Further, RFA was found to be superior to laser ablation. The volume reduction rate for RFA at the last follow-up was 92.2%, whereas in the laser ablation group, the volume reduction rate peaked at 12 months (52.3%) and was at 43.3% at the last follow-up.

To evaluate the efficacy of RFA for the treatment of benign thyroid nodules, Chen et al (2016) conducted a systematic review and meta-analysis and found that RFA was associated with a significant decrease in nodule volume at months 1, 3, 6, 12, and last follow-up.³⁸

Fuller et al (2014) reported on a systematic review of studies on RFA for benign thyroid tumors.³⁹ After RFA, statistically significant improvements were reported in combined symptom improvement and cosmetic scores on the 0 to 6 scale (mean, -2.96; 95% CI, -2.66 to -3.25) and withdrawal from methimazole (odds ratio, 40.34; 95% CI, 7.78 to 209.09). Twelve adverse events were reported, 2 of which were considered significant but did not require hospitalization.

Table 3 includes a comparison of studies included in the systematic reviews; the analyses by Cho et al (2020) and Zu et al (2024) contain the fewest number of included studies as a minimum follow-up duration of 3 years and 5 years, respectively were required for inclusion. Table 4 summarizes the characteristics of the systematic reviews and Table 5 contains the available results for nodule size reduction and complication rates. All of the systematic reviews are limited by high heterogeneity, inclusion of mostly single-center retrospective and/or noncontrolled studies, and generalizability concerns as included studies were mainly conducted in the Republic of Korea and Italy. They are further limited by a lack of comparison to surgical excision or PEI.

Table 3. Comparison of Meta-Analyses Assessing Radiofrequency Ablation for Benign Thyroid Nodules

Study	Xu (2024)* ³⁶ ,	Cho (2020)* ³⁷ ,	Chen (2016) ³⁸ ,	Fuller (2014) ³⁹ ,
Li (2022)	●			
Bernardi (2020)	●			
Aldea Martinez (2019)		●		
Deandrea (2019)	●	●		
Jung (2018)		●		
Sim (2017)		●		
Cesareo (2015)			●	
Sung (2015)			●	
Che (2015)			●	
Ugurlu (2015)			●	
Ji Hong (2015)			●	
Valcavi (2015)			●	
Bernardi (2014)			●	

Study	Xu (2024)* ³⁶ ,	Cho (2020)* ³⁷ ,	Chen (2016) ³⁸ ,	Fuller (2014) ³⁹ ,
Turtulici (2014)			●	
Yoon (2014)			●	
Lim (2013)		●	●	
Ha (2013)			●	
Sung (2013)			●	●
Huh (2012)				●
Faggiano (2012)			●	●
Jang (2012)				●
Kim (2012)				●
Baek (2010)			●	●
Lee (2010)			●	
Spiezia (2009)			●	●
Jeong (2008)			●	
Deandrea (2008)			●	●
Kim (2006)			●	●

*Only studies addressing radiofrequency ablation are included.

Table 4. Characteristics of Meta-Analyses Assessing Radiofrequency Ablation for Benign Thyroid Nodules

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Xu (2024) ³⁶ ,	Through Feb 2023	5	Patients with a benign thyroid nodule treated with thermal ablation (RFA [3 studies], microwave [1 study] or laser [2 studies])	939 (20 to 406)	5 retrospective cohorts	Approximately 5 years

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Cho (2020) ³⁷ ,	2010-2019	12	Patients with a benign thyroid nodule treated with thermal ablation (RFA [5 studies] or laser [7 studies])	1208 (24 to 276)	2 prospective and 10 retrospective cohorts	At least 3 years
Chen (2016) ³⁸ ,	2006-2016	20	Patients with a benign thyroid nodule treated with RFA	1090 (11 to 236)	Prospective and retrospective cohorts	Varied, 6 to 49.4 months
Fuller (2014) ³⁹ ,	2006-2013	9	Patients with a benign thyroid nodule treated with RFA	284 (15 to 94)	Prospective studies (5 observational, 4 randomized trials)	Varied, 3 to 12 months

RFA: radiofrequency ablation.

Table 5. Key Results of Meta-Analyses Assessing Radiofrequency Ablation for Benign Thyroid Nodules

Study	Reduction in nodule size from baseline	Complication rate
Cho (2020) ³⁷ ,	Relative volume reduction, VRR	
Total N, nodules (patients)	695 (680)	695 (680)
Pooled effect (95% CI)	6 months: 64.5% (56.1% to 72.1%) 12 months: 76.9% (65% to 85.7%) 24 months: 80.1% (66.4% to 89.2%) 36 months: 80.3% (66% to 89.5%)	4.6%
I^2 (p)	73.7% to 95.9%	
Chen (2016) ³⁸ ,	Absolute volume reduction, SMD	
Total N, nodules (patients)	1406 (1090)	
Pooled effect (95% CI)	1 month: 0.83 (0.47 to 1.19) 3 months: 1.31 (0.76 to 1.85) 6 months: 1.25 (0.90 to 1.59) 12 months: 4.16 (2.25 to 6.07)	
I^2 (p)	90.3% to 98.7%	

Study	Reduction in nodule size from baseline	Complication rate
Fuller (2014) ³⁹ ,	Absolute volume reduction, SMD (follow-up time frame not specified)	
Total N, nodules (patients)	284 (276)	
Pooled effect (95% CI)	-9.77 mL (-13.83 to -5.72)	
I^2 (p)	98% (<.00001)	
Xu (2024) ^{*36} ,	Volume reduction rate	Regrowth
Total N, nodules (patients)	NR (939)	
Pooled effect (95% CI)	74.48% (70.05 to 78.91)	10.60% (1.50 to 19.80)

CI: confidence interval; NR: not reported; SMD: standard mean difference; VRR: volume reduction rate.

*Includes data for all patients; RFA not separately analyzed from other methods of thermal ablation.

Section Summary: Benign Thyroid Tumors

Evidence on the treatment of benign thyroid nodules includes randomized and nonrandomized trials, case series, and systematic reviews of these studies. Systematic reviews have demonstrated that RFA results in a significant reduction in thyroid nodule size with a 2020 review showing that these changes remain durable through at least 36 months and a 2024 review indicating durability up to 5 years. Complication rates are generally low but include voice changes. The data are limited by significant heterogeneity in meta-analyses, a lack of generalizability to populations outside Republic of Korea and Italy, and a lack of comparators more relevant to practice in the United States.

MISCELLANEOUS SOLID TUMORS

Clinical Context and Therapy Purpose

RFA has been investigated for use in individuals with different lesions in different anatomic sites. These anatomic sites include but are not limited to the thyroid, pancreas, and head and neck.

In individuals with head and neck cancer with recurrent disease, surgical salvage attempts are poor in terms of local control, survival, and QOL; further, these recurrent tumors are often untreatable with standard salvage therapies. Palliative chemotherapy or comfort measures may be offered. The safety and efficacy of RFA have been investigated as an option for palliative treatment in these situations.

The purpose of RFA in patients who have miscellaneous solid tumors is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with miscellaneous solid tumors (eg, head and neck, thyroid cancer, pancreas).

Interventions

The therapy being considered is RFA.

Comparators

The following practices are currently being used to treat miscellaneous solid tumors: surgical excision or other local treatments specific to the tumor type.

Outcomes

The general outcomes of interest vary by disease state but include OS, tumor recurrence, and reductions in pain.

Patient follow-up will vary by disease state.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

REVIEW OF EVIDENCE**Thyroid Cancer**

Kim et al (2015) reported on a comparative review of 73 patients with recurrent thyroid cancer smaller than 2 cm who had been treated with RFA (n=27) or repeat surgery (n=46).⁴⁰ RFA was performed in cases of patient refusal to undergo surgery or poor medical condition. Data were weighted to minimize potential confounders. The 3-year recurrence-free survival rates were similar for RFA (92.6%) and surgery (92.2%, p=.681). Posttreatment hoarseness rate did not differ between the RFA (7.3%) and surgery (9.0%) groups. Posttreatment hypocalcemia occurred only in the surgery group (11.6%).

Head and Neck Cancer

Owen et al (2011) reported on RFA for 13 patients with recurrent and/or unresectable head and neck cancer who failed curative treatment.⁴¹ Median patient survival was 127 days. While the stable disease was reported in 8 patients after RFA, and QOL scores improved, 3 deaths occurred (1 carotid hemorrhage, 2 strokes).

A case series of RFA for 14 patients with recurrent advanced head and neck malignancies was reported by Brook et al (2008).⁴² Tumor targeting and electrode deployment were successful in all cases, and 4 of 6 patients who completed QOL assessments showed improvement. Three major complications (in 27 [11%] applications) occurred 7 days to 2 weeks postprocedure. They included stroke, carotid artery rupture leading to death, and threatened carotid artery rupture with subsequent stroke. Retrospective analysis of intraprocedural CT scans revealed that the retractable electrodes were within 1 cm of the carotid artery during ablation in these cases.

A case series by Owen et al (2004) showed that palliative CT-guided RFA provided subjective improvement with regard to pain, appearance, and function in 12 patients who had recurrent and advanced head and neck malignancies and were not candidates for radiotherapy or surgery.⁴³ The procedure appeared reasonably safe and feasible for this indication.

Uterine Myomas

A prospective observational study by Rey et al (2019) assessed the effectiveness of transvaginal ultrasound-guided RFA of myomas in reducing tumor volume and eliminating menorrhagia associated with myomas.⁴⁴ The study included 205 women with symptomatic type II/III uterine submucosal or intramural cavity-distorting myomas undergoing RFA. The preoperative mean (standard deviation) volume of the myomas was 122.4 (182.5) cm³ (95% CI, 82.1 to 162.8). Mean myoma volume decreased significantly at 1 (85.2 [147.9] cm³; p=.001), 3 (67.3 [138.0] cm³; p=.001), 6 (59.3 [135.3] cm³; p=.001), and 12 months (49.6 [121.4] cm³; p=.001). At 12 months, the mean volume reduction was 60% compared with preoperative volume. All patients returned to normal menstruation at a mean follow-up of 3 months and 12 months. Of the 205 patients, 201 (98.04%) were satisfied with the procedure. The investigators conceded that a larger population with a longer follow-up is needed but their study suggests that transvaginal ultrasound-guided RFA of myomas is effective and safe for treating select patients with menorrhagia secondary to myomas.

In a large series, Yin et al (2015) evaluated the effectiveness and safety of RFA for uterine myomas in a 10-year retrospective cohort study.⁴⁵ From 2001 to 2011, a total of 1216 patients treated for uterine myomas were divided into 2 groups. Group A consisted of 476 premenopausal patients (average age, 36 years) who had an average of 1.7 myomas with an average diameter of 4.5 cm. Group B consisted of 740 menopausal patients (average age, 48 years) with an average of 2.6 myomas with an average diameter of 5.0 cm. Patients were followed for a mean of 36 months. At 1, 3, 6, 12, and 24 months after RFA, the average diameters of myomas in group A were 3.8, 3.0, 2.7, 2.4, and 2.2 cm, respectively; 48% (227/476) of patients had a residual tumor at 12 months. In group B, myoma diameters were 4.7, 3.7, 3.3, 2.3, and 2.3 cm, respectively; 59% (435/740) of patients had trace disease at 12 months. Three months after RFA treatment, myoma volumes were significantly reduced in both groups (p<.01), although group B had a higher rate of residual tumor 12 months after RFA than group A (p<.05). Clinical symptoms and health-related QOL were significantly improved after RFA in both groups. The postoperative recurrence rate of uterine myomas was significantly higher in group A at 10.7% (51/476) than in group B at 2.4% (18/740; p<.05).

Adrenal Tumors

Liu et al (2020) retrospectively evaluated the clinical outcomes of percutaneous ultrasound-guided RFA in the treatment of adrenal metastasis as compared to adrenalectomy.⁴⁶ Of the 60 patients included, 29 received RFA and 31 received adrenalectomy. The first technical success rate for RFA was 72.4%; 5 of the 8 patients had a repeat RFA and 4 of those achieved a complete response. In the adrenalectomy group, all patients achieved a R0 resection. Major complications were reported in 1 patient in the RFA group (ventricular fibrillation) and 2 patients in the adrenalectomy group (ascites, surgical site infection). The 1-, 2-, and 3-year local tumor progression rates after RFA were 17.1%, 30.9% and 44.7%, respectively, compared to 6.5%, 6.5% and 6.5% in adrenalectomy group (p=.028). There was no significant difference between

groups for mean OS (2.3 ± 0.3 years for RFA and 3.9 ± 0.6 years for adrenalectomy, $p=.057$). Limitations of the study include its retrospective design, potential selection bias on which patients received each treatment, and a high prevalence of patients with adrenal metastasis secondary to hepatocellular carcinoma, which exceeded the expected number of cases based on global prevalence rates.

Liu et al (2016) retrospectively compared laparoscopic adrenalectomy with CT-guided percutaneous RFA for the treatment of aldosterone-producing adenoma, evaluating short-term and long-term outcomes of normalized aldosterone-to-renin ratio, hypokalemia, and hypertension.¹⁸ Of 63 patients, 27 were in the laparoscopic adrenalectomy group and 36 were in the RFA group. Primary aldosteronism was seen in 33 of 36 patients treated with RFA and all 27 who had laparoscopic adrenalectomy ($p=.180$), within a median follow-up of 5 to 7 years. RFA was associated with faster recovery postprocedure, but hypertension was less frequently resolved using RFA (13/36 patients) compared with laparoscopic adrenalectomy (19/27 patients; $p=.007$). The use of posture test and CT for subtype classification of primary aldosteronism is the major limitation of the study, as well as the retrospective design.

Retrospectively, Yang et al (2016) compared the efficacy and safety of RFA with laparoscopic adrenalectomy in treating aldosterone-producing adenoma of the adrenal gland.⁴⁷ From 2009 to 2013, 25 patients diagnosed with unilateral adrenal aldosterone-producing adenoma and similar tumor size (<25 mm) were allocated to a control group ($n=18$) that underwent laparoscopic adrenalectomy and a test group ($n=7$) that underwent CT-guided percutaneous RFA. Complete tumor ablation on follow-up CT scan and normalization of serum aldosterone-to-renin were the primary outcomes compared in this study. Success in the RFA group reached 100% within 3 to 6 months, compared with 94.4% in the laparoscopic adrenalectomy group, and normalization ability was statistically equivalent in both groups. The study's retrospective design and small sample are the main limitations of this study.

Other Tumors

A single-arm, retrospective, paired-comparison study by Locklin et al (2004) evaluated the short-term efficacy of RFA in reducing pain and improving function in patients with unresectable, painful soft tissue neoplasms recalcitrant to conventional therapies.⁴⁸ Patients had tumors located in a variety of sites including chest wall, pelvis, breast, perirectal, renal, aortocaval, retroperitoneal, and superficial soft tissues. All had failed conventional methods of palliation or experienced dose-limiting adverse events from pain medication. Although not all Brief Pain Inventory scores were statistically significant, all mean scores trended down over time after ablation. Complications from RFA were minor or insignificant in all but 1 patient who had skin breakdown and infection of an ablated superficial tumor site.

Additional research has addressed the use of RFA in solid malignancies^{49,50}, and in the pancreas.^{51,52,53} A systematic review by Rombouts et al (2015) has examined studies of ablative therapies, including RFA, in patients with locally advanced pancreatic cancer.⁵⁴ No RCTs were identified, and conclusions were limited by the sparse evidence available on RFA in this setting.

Stereotactic radiofrequency thermocoagulation for epileptogenic hypothalamic hamartomas was described in a retrospective analysis by Kameyama et al (2009) who evaluated 25 patients with gelastic seizures (a rare type of seizure).⁵⁵ Other seizure types were exhibited in 22 (88.0%)

patients, precocious puberty in 8 (32.0%), behavioral disorder in 10 (40.0%), and mental disability in 14 (56.0%). Gelastic seizures resolved in all but 2 patients. Complete seizure freedom was achieved in 19 (76.0%) patients. These patients experienced resolution of all seizure types and behavioral disorder and also demonstrated intellectual improvement.

Preliminary results of endoscopic RFA of rectosigmoid tumors have been described by Vavra et al (2009).⁵⁶ Twelve patients were treated with the Endoblate RFA device, with 10 patients having surgical resection after ablation. Histology of the resected specimens showed that, on average, 82% (range, 60% to 99%) of the tumor mass was destroyed in the ablation zone.

Small case series on RFA for colorectal and rectal carcinoma have demonstrated a debulking role for RFA.^{57,58} These case series did not permit comparison with an available alternative.

Section Summary: Miscellaneous Solid Tumor

Evidence on the use of RFA to treat other types of solid tumors consists of a small number of case series, prospective studies, or retrospective comparative studies. Reporting on outcomes is limited. The evidence base does not support a conclusion on the effects of RFA for the tumor types included in this evidence review.

SUPPLEMENTAL INFORMATION

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2010 Input

In response to requests, input was received from 2 physician specialty societies (4 reviewers) and 2 academic medical centers (4 reviewers) while this policy was under review in 2010. Input was similar to that received in 2009, except support for the use of radiofrequency ablation (RFA) to treat lung tumors was declined (only 1 respondent indicated this was an option in tumors metastatic to lung). One respondent also indicated a potential use for adrenal tumors. Input supported RFA for localized renal cell carcinoma no more than 4 cm in size when preservation of kidney function is necessary and a standard surgical approach would likely substantially worsen kidney function or when the patient is not considered a surgical candidate.

2009 Input

In response to requests, input was received from 1 physician specialty society (4 reviews) and from 2 academic medical centers (3 reviews) while this policy was under review in 2009. All reviewers supported the use of RFA in the treatment of painful bone metastases that have failed standard treatment and in the treatment of osteoid osteomas. Reviewers were divided over the use of RFA for lung tumors, although several agreed that, while it may be useful in a select population of patients, it should be used in the clinical trial setting. Reviewers were also split with regard to RFA in the treatment of renal tumors, with some supporting its use in a select

population of patients. With the exception of 1 disagreement and 1 nonresponse, the reviewers agreed to the investigational statement on the use of RFA in all other tumors outside the liver that are addressed in this policy.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Chest Physicians

The American College of Chest Physicians (2025) guidelines on management of patients with early-stage non-small cell lung cancer state that "image-guided thermal ablation therapies such as radiofrequency ablation (RFA), microwave ablation (MWA), and cryoablation (CA) have also been used to treat lung tumors, particularly in patients with contraindications to surgery. For patients deemed to have "high operative risk" (i.e., those who cannot tolerate lobectomy, but are candidates for sublobar resection) stage I NSCLC, the appropriateness of potential alternatives [e.g., SBRT; ablation] to surgery is an active area of investigation".⁵⁹ They also state that "To date, no specific technique of ablation (radiofrequency, microwave, or cryoablation) has proven superior, and therefore outcomes from various modalities are most often combined to evaluate local control and overall survival."

American Head and Neck Society - Endocrine Surgery Section

An international, multidisciplinary consensus statement on RFA and related ultrasound-guided ablation technologies for the treatment of benign and malignant thyroid disease was released in 2022 through a collaboration of international professional societies, including the Endocrine Surgery Section of the American Head and Neck Society.⁶⁰ Select relevant recommendations from the guideline are listed in Table 6.

Table 6. Summary of RFA Recommendations for Treatment of Benign and Malignant Thyroid Disease*

Recommendation 1	US-guided ablation procedures may be used as a first-line alternative to surgery for patients with benign thyroid nodules contributing to compressive and/or cosmetic symptoms.
Recommendation 2	Although less efficacious than surgery or RAI in normalizing thyroid function, thermal ablation procedures can be a safe therapeutic alternative in patients with an autonomously functional thyroid nodule and contraindications to first-line techniques.
Recommendation 3a	US-guided ablation procedures may be considered in patients with suitable primary papillary microcarcinoma who are unfit for surgery or decline surgery or active surveillance
Recommendation 3b	US-guided ablation procedures may be considered in patients with suitable recurrent papillary thyroid carcinoma who are unfit for surgery or decline surgery or active surveillance
Recommendation 3c	Repeat ablation of a benign nodule can be considered for remnant nodular tissue contributing to unresolved symptomatic or cosmetic concerns

*This is not a comprehensive list of recommendations from the guideline.
RAI: radioactive iodine; RFA: radiofrequency ablation; US: Ultrasound.

American Urological Association

The American Urological Association (AUA; 2017) guideline on renal masses and localized renal cancer affirms that partial nephrectomy should be prioritized for the management of cT1a renal masses when intervention is indicated.⁶¹ Thermal ablation should be considered "as an alternate approach for the management of cT1a renal masses <3 cm in size." The guidelines were updated in 2021 and recommendations are generally consistent with what was published in the 2017 guideline.⁶² The 2021 AUA guideline explicitly states that RFA and cryoablation may be offered as options to patients who elect thermal ablation.

American Thyroid Association

The American Thyroid Association (2015) guideline on the management of thyroid nodules and differentiated thyroid cancer provides recommendations for management.³⁵ Patients with a benign cytology diagnosis or those very unlikely to be malignant (eg, purely cystic nodule) should undergo surveillance with the frequency determined by the level of suspicion for a missed malignancy. Medical or surgical intervention is considered if the nodules are large (>4 cm), causing compressive or structural symptoms, or if there is clinical concern. Recurrent cystic thyroid nodules with benign cytology should be considered for surgical removal or percutaneous ethanol injection. For differentiated thyroid cancer, "localized treatments with thermal (radiofrequency or cryo-) ablation, ethanol ablation, or chemoembolization may be beneficial in patients with a single or a few metastases and in those with metastases at high risk of local complications."

National Comprehensive Cancer Network

The National Comprehensive Cancer Network (NCCN) guidelines for the treatment of NSCLC (v.7.2025) state:⁶³ "For medically operable disease, resection is the preferred local treatment modality (other modalities include SABR [stereotactic ablative radiotherapy], thermal ablation such as radiofrequency ablation, and cryotherapy)." For patients who are not amenable to surgery, image-guided thermal ablation therapy (IGTA; includes RFA, microwave ablation, and cryoablation) may be considered. The guidance states "IGTA is an option for the management of NSCLC lesions <3 cm. Ablation for NSCLC lesions >3 cm may be associated with higher rates of local recurrence and complications."

The NCCN guidelines for thyroid carcinoma (v.1.2025) indicate that local therapies such as RFA may be considered for locoregional recurrence of thyroid carcinoma-papillary carcinoma in select patients with limited burden nodal disease. Additionally, local therapies, including RFA, can be considered in those with metastatic disease.⁶⁴

The NCCN guidelines (v.1.2026) for renal cell carcinoma indicate that " percutaneous ablation (eg, cryosurgery, radiofrequency ablation, microwave ablation) is an option for the management of clinical stage T1 renal lesions. Percutaneous ablation is an option for clinical T1b masses in select patients not eligible for surgery. Biopsy of lesions is recommended to be done prior to or at time of ablation. Percutaneous ablation may require multiple treatments to achieve the same oncologic outcomes as conventional surgery."⁶⁵

The NCCN colon cancer guidelines (v.4.2025) state that "resection is the standard approach for the local treatment of resectable metastatic disease. However, patients with liver or lung oligometastases can also be considered for tumor ablation therapy, particularly in cases that may not be optimal for resection." ⁶⁶"There is extensive evidence on the use of RFA as a reasonable treatment option for non-surgical candidates and for recurrent disease after hepatectomy with small liver metastases that can be treated with clear margins."

The NCCN guidelines for head and neck cancers (v.4.2025),⁶⁷ breast cancer (v.4.2025),⁶⁸ bone cancer (v.2.2025),⁶⁹ and pancreatic adenocarcinoma (v.2.2025) do not mention RFA.⁷⁰

National Institute for Health and Care Excellence

The NICE guidance (2004) on osteoid osteoma indicated that "current evidence on the safety and efficacy of computed tomography (CT)-guided thermocoagulation of osteoid osteoma appears adequate to support its use...."⁷¹,

Updated NICE guidance (2010) on renal cancer has indicated that "evidence on the safety and efficacy of percutaneous radiofrequency ablation (RFA) ... in the short and medium term appears adequate to support the use of this procedure provided that patients are followed up in the long term."⁷⁰,

The NICE guidance (2010) on RFA for primary and secondary lung cancers has stated: "Current evidence on the efficacy of percutaneous radiofrequency ablation (RFA) ... is adequate in terms of tumor control." ⁷²The NICE also indicated RFA might "be used in patients with small, early-stage lung cancers or small numbers of lung metastases who are unsuitable for, or prefer not to undergo, surgery. It may also have a place in multi-modality treatment of more advanced primary lung cancers." The guidance warned of serious complications (eg, pneumothorax) among lung cancer patients.

The NICE guidance (2016) on benign thyroid nodules stated: "Current evidence on the safety and efficacy of ultrasound-guided percutaneous radiofrequency ablation ... is adequate to support the use of this procedure...."⁷³,

Society of Interventional Radiology

The Society of Interventional Radiology (2020) published a position statement on the role of percutaneous ablation in renal cell carcinoma.⁷⁴ Their relevant recommendations are as follows:

- "In patients with small renal tumors (stage T1a), percutaneous thermal ablation is a safe and effective treatment with fewer complications than nephrectomy and acceptable long-term oncological and survival outcomes. (Level of Evidence: C; Strength of Recommendation: Moderate)"
- "In selected patients with suspected T1a renal cell carcinoma, percutaneous thermal ablation should be offered over active surveillance. (Level of Evidence: C; Strength of Recommendation: Moderate)"
- "In high-risk patients with T1b renal cell carcinoma who are not surgical candidates, percutaneous thermal ablation may be an appropriate treatment option; however, further research in this area is required. (Level of Evidence: D; Strength of Recommendation: Weak)"

- "Radiofrequency ablation, cryoablation, and microwave ablation are all appropriate modalities for thermal ablation, and method of ablation should be left to the discretion of the operating physician. (Level of Evidence: D; Strength of Recommendation: Weak)"

U.S. Preventive Services Task Force Recommendations

Not applicable.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 7.

Table 7. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT05189821	RFA Treatment for Papillary Thyroid Microcarcinoma Cohort	50	Nov 2027
NCT05189808	Radiofrequency Ablation for Indeterminate Bethesda III Thyroid Nodules	50	Aug 2027
<i>Unpublished</i>			
NCT04619472	A Multicenter, Single Group Target Value Clinical Study to Evaluate Safety and Effectiveness of Radiofrequency Ablation System in the Treatment of Peripheral Lung Tumors	126	Mar 2023
NCT01051037	Phase II Study Evaluating Safety and Efficacy of Stereotactic Body Radiotherapy and Radiofrequency Ablation for Medically Inoperable and Recurrent Lung Tumors Near Central Airways	17	Dec 2017 (completed)

NCT: national clinical trial.

CODING

The following codes for treatment and procedures applicable to this policy are included below for informational purposes. This may not be a comprehensive list of procedure codes applicable to this policy.

Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

The code(s) listed below are medically necessary ONLY if the procedure is performed according to the "Policy" section of this document.

CPT/HCPCS	
20982	Ablation therapy for reduction or eradication of 1 or more bone tumors (e.g., metastasis) including adjacent soft tissue when involved by tumor extension, percutaneous, including imaging guidance when performed; radiofrequency
32998	Ablation therapy for reduction or eradication of 1 or more pulmonary tumor(s) including pleura or chest wall when involved by tumor extension, percutaneous, including imaging guidance when performed, unilateral; radiofrequency
50542	Laparoscopy, surgical; ablation of renal mass lesion(s)
50592	Ablation, one or more renal tumor(s), percutaneous, unilateral, radiofrequency
60660	Percutaneous ablation of 1 or more thyroid nodule(s)
60661	Percutaneous ablation of additional lobe of thyroid nodule(s)

REVISIONS	
04-01-2011	<p>Medical Policy Language section updated.</p> <p>In the policy language:</p> <ul style="list-style-type: none"> ▪ Removed "Radiofrequency ablation is considered medically necessary as an alternative to surgical resection for debulking of primary and metastatic neoplasms." ▪ In Item #2, removed "surgery cannot be tolerated" and inserted "patient is not considered an appropriate surgical candidate" to read Radiofrequency ablation is considered medically necessary for removal of primary or metastatic malignant neoplasms when removal of the neoplasm may be curative and the patient is not considered an appropriate surgical candidate. ▪ Added item #4, "Radiofrequency ablation may be considered medically necessary to treat osteoid osteoma that cannot be managed successfully with medical treatment." <p>Rationale section updated.</p> <p>Reference section updated.</p>
12-11-2013	<p>Replaced medical policy reference, "Treatment of Liver Tumors" with the following references:</p> <p><i>"Radioembolization for Primary and Metastatic Tumors of the Liver</i> <i>Radiofrequency Ablation of Primary or Metastatic Liver Tumors</i> <i>Transcatheter Arterial Chemoembolization (TACE) to Treat Primary or Metastatic Liver Malignancies</i></p>

REVISIONS	
	<i>Cryosurgical Ablation of Primary or Metastatic Liver Tumors"</i>
	Updated Description section.
	Updated Rationale section.
	In Coding section:
	▪ Added ICD-10 Diagnosis (<i>Effective October 1, 2014</i>)
	Updated Reference section.
01-01-2015	In Coding section:
	▪ Revised CPT Code: 20982 (<i>Effective January 1, 2015</i>)
11-11-2016	Updated Description section.
	In Policy section:
	<ul style="list-style-type: none"> ▪ Removed previous Item B, "Radiofrequency ablation is considered medically necessary for removal of primary or metastatic malignant neoplasms when removal of the neoplasm may be curative and the patient is not considered an appropriate surgical candidate." ▪ Added new Item C, "Radiofrequency ablation may be considered medically necessary to treat localized renal cell carcinoma that is no more than 4 cm in size when either of the following criteria is met: 1. In order to preserve kidney function; OR 2. The patient is not considered a surgical candidate." ▪ Added new Item D, "Radiofrequency ablation may be considered medically necessary to treat an isolated peripheral non-small-cell lung cancer lesion that is no more than 3 cm in size when the following criteria are met: 1. Surgical resection or radiation treatment with curative intent is considered appropriate based on stage of disease; however, medical co-morbidity renders the individual unfit for those interventions; AND 2. Tumor is located at least 1 cm from the trachea, main bronchi, esophagus, aorta, aortic arch branches, pulmonary artery, and the heart." ▪ Removed previous Item E, "Radiofrequency ablation is considered experimental / investigational for curative treatment of primary or metastatic malignant neoplasms in surgical candidates because the effectiveness of radiofrequency tumor ablation is improving clinical outcomes has not been established." ▪ Added new Item E, "Radiofrequency ablation may be considered medically necessary to treat malignant nonpulmonary tumor(s) metastatic to the lung that are no more than 3 cm in size when the following criteria are met: 1. In order to preserve lung function when surgical resection or radiation treatment is likely to substantially worsen pulmonary status OR the patient is not considered a surgical candidate; AND 2. There is no evidence of extrapulmonary metastases; AND 3. The tumor is located at least 1 cm from the trachea, main bronchi, esophagus, aorta, aortic arch branches, pulmonary artery, and the heart. The following additional criteria have been developed by clinical judgment / consensus and existing guidelines for the use of RFA in metastatic tumors to the lung and include: 1) No more than 3 tumors per lung should be ablated; b) Tumors should be amenable to complete ablation; AND c) Twelve months should elapse before a repeat ablation is considered." ▪ Removed previous Item F, "Radiofrequency tumor ablation, excluding the liver, is considered experimental / investigational for all other indications." ▪ Added new Item F, "Radiofrequency ablation is considered experimental / investigational as a technique for ablation of: 1. Breast tumors; 2. Lung cancer not meeting the criteria above; 3. Renal cell cancer not meeting the criteria above; 4. Osteoid osteomas that can be managed with medical treatment; 5. Painful bony metastases as initial treatment; AND 6. All other tumors outside the liver including,

REVISIONS	
	but not limited to, the head and neck, thyroid, adrenal gland, ovary, and pelvic/abdominal metastases of unspecified origin."
	Updated Rationale section.
	In Coding section:
	<ul style="list-style-type: none"> ▪ Added ICD-10 codes: C64.1, C64.2, C79.51, C79.52.
	Updated References section.
10-25-2017	Updated Description section.
	Updated Rationale section.
	Updated References section.
01-01-2018	In Coding section:
	<ul style="list-style-type: none"> ▪ Revised nomenclature to CPT code: 32998. ▪ Removed ICD-9 codes.
12-07-2018	Policy published to the bcbsks.com website on 11-07-2018 with an effective date of 12-07-2018.
	Updated Description section.
	Updated Rationale section.
	In Coding section:
	<ul style="list-style-type: none"> ▪ Added ICD-10 codes: C78.01, C78.02. ▪ Removed ICD-10 codes: C33, C68.8, C68.9.
	Updated References section.
02-24-2021	Updated Description section.
	Updated Rationale section.
	Updated References section.
12-9-2022	Updated Description Section
	Updated Policy Section
	<ul style="list-style-type: none"> ▪ Added Subtitles to sections: "Osteolytic bone metastases" to section A, ▪ "Osteoid osteomas" to section B, " Renal cell carcinoma" to section C, "Non-small-cell lung cancer" to section D, "Nonpulmonary tumor(s) metastatic to the lung" to section E ▪ Added: "(See the Policy Guidelines Section for additional criteria)" to Section E3
	Updated Guideline Section
	<ul style="list-style-type: none"> ▪ Added Policy Guidelines Section: <ul style="list-style-type: none"> A. The following additional criteria have been developed by clinical judgment / consensus and existing guidelines for the use of RFA in metastatic tumors to the lung and include: <ol style="list-style-type: none"> 1. No more than 3 tumors per lung should be ablated; 2. Tumors should be amenable to complete ablation; AND 3. Twelve months should elapse before a repeat ablation is considered.
	Updated Rationale Section
	Updated Coding Section
	<ul style="list-style-type: none"> ▪ Removed 76940 ▪ Converted ICD-10 codes to ranges, to include all codes within the range
	Updated References Section
10-24-2023	Updated Description Section
	Updated Rationale Section
	Updated Coding Section
	<ul style="list-style-type: none"> ▪ Removed ICD-10 Codes
	Updated References Section

REVISIONS	
10-22-2024	Updated Description Section
	Updated Rationale Section
	Updated References Section
01-01-2025	Updated Coding Section <ul style="list-style-type: none"> ▪ Added 60660 and 60661 (eff. 01-01-2025)
10-28-2025	Updated Description Section
	Updated Rationale Section
	Updated Reference Section

REFERENCES

1. Levy J, Hopkins T, Morris J, et al. Radiofrequency Ablation for the Palliative Treatment of Bone Metastases: Outcomes from the Multicenter OsteoCool Tumor Ablation Post-Market Study (OPuS One Study) in 100 Patients. *J Vasc Interv Radiol*. Nov 2020; 31(11): 1745-1752. PMID 33129427
2. Goetz MP, Callstrom MR, Charboneau JW, et al. Percutaneous image-guided radiofrequency ablation of painful metastases involving bone: a multicenter study. *J Clin Oncol*. Jan 15 2004; 22(2): 300-6. PMID 14722039
3. Grönemeyer DH, Schirp S, Gevargez A. Image-guided radiofrequency ablation of spinal tumors: preliminary experience with an expandable array electrode. *Cancer J*. 2002; 8(1): 33-9. PMID 11898806
4. Kojima H, Tanigawa N, Kariya S, et al. Clinical assessment of percutaneous radiofrequency ablation for painful metastatic bone tumors. *Cardiovasc Intervent Radiol*. 2006; 29(6): 1022-6. PMID 16988875
5. Tordjman M, Perronne L, Madelin G, et al. CT-guided radiofrequency ablation for osteoid osteomas: a systematic review. *Eur Radiol*. Nov 2020; 30(11): 5952-5963. PMID 32518986
6. Lanza E, Thouvenin Y, Viala P, et al. Osteoid osteoma treated by percutaneous thermal ablation: when do we fail? A systematic review and guidelines for future reporting. *Cardiovasc Intervent Radiol*. Dec 2014; 37(6): 1530-9. PMID 24337349
7. Albisinni U, Facchini G, Spinnato P, et al. Spinal osteoid osteoma: efficacy and safety of radiofrequency ablation. *Skeletal Radiol*. Aug 2017; 46(8): 1087-1094. PMID 28497160
8. Lassalle L, Campagna R, Corcos G, et al. Therapeutic outcome of CT-guided radiofrequency ablation in patients with osteoid osteoma. *Skeletal Radiol*. Jul 2017; 46(7): 949-956. PMID 28429047
9. Rimondi E, Mavrogenis AF, Rossi G, et al. Radiofrequency ablation for non-spinal osteoid osteomas in 557 patients. *Eur Radiol*. Jan 2012; 22(1): 181-8. PMID 21842430
10. Sahin C, Oc Y, Ediz N, et al. The safety and the efficacy of computed tomography guided percutaneous radiofrequency ablation of osteoid osteoma. *Acta Orthop Traumatol Turc*. Sep 2019; 53(5): 360-365. PMID 31371131
11. Knudsen M, Riishede A, Lücke A, et al. Computed tomography-guided radiofrequency ablation is a safe and effective treatment of osteoid osteoma located outside the spine. *Dan Med J*. May 2015; 62(5). PMID 26050823
12. Rosenthal DI, Hornicek FJ, Torriani M, et al. Osteoid osteoma: percutaneous treatment with radiofrequency energy. *Radiology*. Oct 2003; 229(1): 171-5. PMID 12944597

13. Huang RS, Chow R, Benour A, et al. Comparative efficacy and safety of ablative therapies in the management of primary localised renal cell carcinoma: a systematic review and meta-analysis. *Lancet Oncol.* Mar 2025; 26(3): 387-398. PMID 39922208
14. Yanagisawa T, Mori K, Kawada T, et al. Differential efficacy of ablation therapy versus partial nephrectomy between clinical T1a and T1b renal tumors: A systematic review and meta-analysis. *Urol Oncol.* Jul 2022; 40(7): 315-330. PMID 35562311
15. Uhlig J, Strauss A, Rücker G, et al. Partial nephrectomy versus ablative techniques for small renal masses: a systematic review and network meta-analysis. *Eur Radiol.* Mar 2019; 29(3): 1293-1307. PMID 30255245
16. Katsanos K, Mailli L, Krokidis M, et al. Systematic review and meta-analysis of thermal ablation versus surgical nephrectomy for small renal tumours. *Cardiovasc Intervent Radiol.* Apr 2014; 37(2): 427-37. PMID 24482030
17. El Dib R, Touma NJ, Kapoor A. Cryoablation vs radiofrequency ablation for the treatment of renal cell carcinoma: a meta-analysis of case series studies. *BJU Int.* Aug 2012; 110(4): 510-6. PMID 22304329
18. Liu SY, Chu CM, Kong AP, et al. Radiofrequency ablation compared with laparoscopic adrenalectomy for aldosterone-producing adenoma. *Br J Surg.* Oct 2016; 103(11): 1476-86. PMID 27511444
19. Marshall HR, Shakeri S, Hosseiny M, et al. Long-Term Survival after Percutaneous Radiofrequency Ablation of Pathologically Proven Renal Cell Carcinoma in 100 Patients. *J Vasc Interv Radiol.* Jan 2020; 31(1): 15-24. PMID 31767409
20. Dai Y, Covarrubias D, Uppot R, et al. Image-Guided Percutaneous Radiofrequency Ablation of Central Renal Cell Carcinoma: Assessment of Clinical Efficacy and Safety in 31 Tumors. *J Vasc Interv Radiol.* Dec 2017; 28(12): 1643-1650. PMID 28673657
21. Dvorak P, Hoffmann P, Brodak M, et al. Percutaneous radiofrequency and microwave ablation in the treatment of renal tumors - 10 years of experience. *Wideochir Inne Tech Maloinwazyjne.* Dec 2017; 12(4): 394-402. PMID 29362655
22. Iannuccilli JD, Dupuy DE, Beland MD, et al. Effectiveness and safety of computed tomography-guided radiofrequency ablation of renal cancer: a 14-year single institution experience in 203 patients. *Eur Radiol.* Jun 2016; 26(6): 1656-64. PMID 26373755
23. Schlijper RC, Grutters JP, Houben R, et al. What to choose as radical local treatment for lung metastases from colo-rectal cancer: surgery or radiofrequency ablation?. *Cancer Treat Rev.* Feb 2014; 40(1): 60-7. PMID 23768754
24. Ratko TA, Vats V, Brock J, et al. Local Nonsurgical Therapies for Stage I and Symptomatic Obstructive Non- Small-Cell Lung Cancer (Comparative Effectiveness Review No. 112). Rockville, MD: Agency for Healthcare Research and Quality; 2013.
25. Bilal H, Mahmood S, Rajashanker B, et al. Is radiofrequency ablation more effective than stereotactic ablative radiotherapy in patients with early stage medically inoperable non-small cell lung cancer?. *Interact Cardiovasc Thorac Surg.* Aug 2012; 15(2): 258-65. PMID 22581864
26. Chan VO, McDermott S, Malone DE, et al. Percutaneous radiofrequency ablation of lung tumors: evaluation of the literature using evidence-based techniques. *J Thorac Imaging.* Feb 2011; 26(1): 18-26. PMID 20829720
27. Hasegawa T, Takaki H, Kodama H, et al. Three-year Survival Rate after Radiofrequency Ablation for Surgically Resectable Colorectal Lung Metastases: A Prospective Multicenter Study. *Radiology.* Mar 2020; 294(3): 686-695. PMID 31934829

28. Xia LY, Hu QL, Xu WY. Efficacy and Safety of Radiofrequency Ablation for Breast Cancer Smaller Than 2 cm: A Systematic Review and Meta-Analysis. *Front Oncol.* 2021; 11: 651646. PMID 34012918
29. Peek MCL, Ahmed M, Napoli A, et al. Minimally invasive ablative techniques in the treatment of breast cancer: a systematic review and meta-analysis. *Int J Hyperthermia.* Mar 2017; 33(2): 191-202. PMID 27575566
30. Zhao Z, Wu F. Minimally-invasive thermal ablation of early-stage breast cancer: a systemic review. *Eur J Surg Oncol.* Dec 2010; 36(12): 1149-55. PMID 20889281
31. Soukup B, Bismohun S, Reefy S, et al. The evolving role of radiofrequency ablation therapy of breast lesions. *Anticancer Res.* Sep 2010; 30(9): 3693-7. PMID 20944155
32. Ito T, Oura S, Nagamine S, et al. Radiofrequency Ablation of Breast Cancer: A Retrospective Study. *Clin Breast Cancer.* Aug 2018; 18(4): e495-e500. PMID 29079443
33. Li P, Xiao-Yin T, Cui D, et al. Evaluation of the safety and efficacy of percutaneous radiofrequency ablation for treating multiple breast fibroadenoma. *J Cancer Res Ther.* Dec 2016; 12(Supplement): C138-C142. PMID 28230006
34. Wilson M, Korourian S, Boneti C, et al. Long-term results of excision followed by radiofrequency ablation as the sole means of local therapy for breast cancer. *Ann Surg Oncol.* Oct 2012; 19(10): 3192-8. PMID 22911363
35. Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid.* Jan 2016; 26(1): 1-133. PMID 26462967
36. Xu X, Peng Y, Han G. Five-year follow-up results of thermal ablation for benign thyroid nodules: Systematic review and meta-analysis. *Am J Otolaryngol.* 2024; 45(1): 104025. PMID 37639985
37. Cho SJ, Baek JH, Chung SR, et al. Long-Term Results of Thermal Ablation of Benign Thyroid Nodules: A Systematic Review and Meta-Analysis. *Endocrinol Metab (Seoul).* Jun 2020; 35(2): 339-350. PMID 32615718
38. Chen F, Tian G, Kong D, et al. Radiofrequency ablation for treatment of benign thyroid nodules: A PRISMA-compliant systematic review and meta-analysis of outcomes. *Medicine (Baltimore).* Aug 2016; 95(34): e4659. PMID 27559968
39. Fuller CW, Nguyen SA, Lohia S, et al. Radiofrequency ablation for treatment of benign thyroid nodules: systematic review. *Laryngoscope.* Jan 2014; 124(1): 346-53. PMID 24122763
40. Kim JH, Yoo WS, Park YJ, et al. Efficacy and Safety of Radiofrequency Ablation for Treatment of Locally Recurrent Thyroid Cancers Smaller than 2 cm. *Radiology.* Sep 2015; 276(3): 909-18. PMID 25848897
41. Owen RP, Khan SA, Negassa A, et al. Radiofrequency ablation of advanced head and neck cancer. *Arch Otolaryngol Head Neck Surg.* May 2011; 137(5): 493-8. PMID 21576561
42. Brook AL, Gold MM, Miller TS, et al. CT-guided radiofrequency ablation in the palliative treatment of recurrent advanced head and neck malignancies. *J Vasc Interv Radiol.* May 2008; 19(5): 725-35. PMID 18440462
43. Owen RP, Silver CE, Ravikumar TS, et al. Techniques for radiofrequency ablation of head and neck tumors. *Arch Otolaryngol Head Neck Surg.* Jan 2004; 130(1): 52-6. PMID 14732768

44. Rey VE, Labrador R, Falcon M, et al. Transvaginal Radiofrequency Ablation of Myomas: Technique, Outcomes, and Complications. *J Laparoendosc Adv Surg Tech A*. Jan 2019; 29(1): 24-28. PMID 30198831
45. Yin G, Chen M, Yang S, et al. Treatment of uterine myomas by radiofrequency thermal ablation: a 10-year retrospective cohort study. *Reprod Sci*. May 2015; 22(5): 609-14. PMID 25355802
46. Liu B, Mo C, Wang W, et al. Treatment outcomes of percutaneous radiofrequency ablation versus adrenalectomy for adrenal metastases: a retrospective comparative study. *J Endocrinol Invest*. Sep 2020; 43(9): 1249-1257. PMID 32166699
47. Yang MH, Tyan YS, Huang YH, et al. Comparison of radiofrequency ablation versus laparoscopic adrenalectomy for benign aldosterone-producing adenoma. *Radiol Med*. Oct 2016; 121(10): 811-9. PMID 27300650
48. Locklin JK, Mannes A, Berger A, et al. Palliation of soft tissue cancer pain with radiofrequency ablation. *J Support Oncol*. 2004; 2(5): 439-45. PMID 15524075
49. Rosenthal DI. Radiofrequency treatment. *Orthop Clin North Am*. Jul 2006; 37(3): 475-84, viii. PMID 16846772
50. Liapi E, Geschwind JF. Transcatheter and ablative therapeutic approaches for solid malignancies. *J Clin Oncol*. Mar 10 2007; 25(8): 978-86. PMID 17350947
51. Spiliotis JD, Datsis AC, Michalopoulos NV, et al. Radiofrequency ablation combined with palliative surgery may prolong survival of patients with advanced cancer of the pancreas. *Langenbecks Arch Surg*. Jan 2007; 392(1): 55-60. PMID 17089173
52. Zou YP, Li WM, Zheng F, et al. Intraoperative radiofrequency ablation combined with 125 iodine seed implantation for unresectable pancreatic cancer. *World J Gastroenterol*. Oct 28 2010; 16(40): 5104-10. PMID 20976848
53. Cantore M, Girelli R, Mambrini A, et al. Combined modality treatment for patients with locally advanced pancreatic adenocarcinoma. *Br J Surg*. Aug 2012; 99(8): 1083-8. PMID 22648697
54. Rombouts SJ, Vogel JA, van Santvoort HC, et al. Systematic review of innovative ablative therapies for the treatment of locally advanced pancreatic cancer. *Br J Surg*. Feb 2015; 102(3): 182-93. PMID 25524417
55. Kameyama S, Murakami H, Masuda H, et al. Minimally invasive magnetic resonance imaging-guided stereotactic radiofrequency thermocoagulation for epileptogenic hypothalamic hamartomas. *Neurosurgery*. Sep 2009; 65(3): 438-49; discussion 449. PMID 19687687
56. Vavra P, Dostalík J, Zacharoulis D, et al. Endoscopic radiofrequency ablation in colorectal cancer: initial clinical results of a new bipolar radiofrequency ablation device. *Dis Colon Rectum*. Feb 2009; 52(2): 355-8. PMID 19279436
57. Mylona S, Karagiannis G, Patsoura S, et al. Palliative treatment of rectal carcinoma recurrence using radiofrequency ablation. *Cardiovasc Intervent Radiol*. Aug 2012; 35(4): 875-82. PMID 22167304
58. Ripley RT, Gajdos C, Reppert AE, et al. Sequential radiofrequency ablation and surgical debulking for unresectable colorectal carcinoma: thermo-surgical ablation. *J Surg Oncol*. Feb 2013; 107(2): 144-7. PMID 22927225
59. Howington J, Souter LH, Arenberg D, et al. Management of Patients With Early-Stage Non-Small Cell Lung Cancer: An American College of Chest Physicians Clinical Practice Guideline. *Chest*. Jun 23 2025. PMID 40562304

60. Orloff LA, Noel JE, Stack BC, et al. Radiofrequency ablation and related ultrasound-guided ablation technologies for treatment of benign and malignant thyroid disease: An international multidisciplinary consensus statement of the American Head and Neck Society Endocrine Surgery Section with the Asia Pacific Society of Thyroid Surgery, Associazione Medici Endocrinologi, British Association of Endocrine and Thyroid Surgeons, European Thyroid Association, Italian Society of Endocrine Surgery Units, Korean Society of Thyroid Radiology, Latin American Thyroid Society, and Thyroid Nodules Therapies Association. *Head Neck*. Mar 2022; 44(3): 633-660. PMID 34939714
61. Campbell S, Uzzo RG, Allaf ME, et al. Renal Mass and Localized Renal Cancer: AUA Guideline. *J Urol*. Sep 2017; 198(3): 520-529. PMID 28479239
62. Campbell SC, Clark PE, Chang SS, et al. Renal Mass and Localized Renal Cancer: Evaluation, Management, and Follow-Up: AUA Guideline: Part I. *J Urol*. Aug 2021; 206(2): 199-208. PMID 34115547
63. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Non-small cell lung cancer. Version 7.2025. Updated July 10 2025. https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed August 5, 2025.
64. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Thyroid Carcinoma. Version 1.2025. Updated March 27, 2025. https://www.nccn.org/professionals/physician_gls/pdf/thyroid.pdf. Accessed August 6, 2025.
65. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Kidney Cancer. Version 1.2026. Updated July 24, 2025. https://www.nccn.org/professionals/physician_gls/pdf/kidney.pdf. Accessed August 4, 2025.
66. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Colon Cancer. Version 4.2025. Updated June 27, 2025. https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf. Accessed August 3, 2025.
67. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Head and Neck Cancers. Version 4.2025. Updated June 20, 2025. https://www.nccn.org/professionals/physician_gls/pdf/head-and-neck.pdf. Accessed July 27, 2025.
68. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Breast Cancer. Version 4.2025. Updated April 17, 2025. https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf. Accessed August 2, 2025.
69. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Bone Cancer. Version 2.2025. Updated February 28, 2025. https://www.nccn.org/professionals/physician_gls/pdf/bone.pdf. Accessed August 1, 2025.
70. National Institute for Health and Care Excellence (NICE). Percutaneous radiofrequency ablation of renal cancer [IPG353]. 2010; <https://www.nice.org.uk/guidance/ipg353>. Accessed July 28, 2025.
71. National Institute for Health and Care Excellence (NICE). Computed tomography-guided thermocoagulation of osteoid osteoma [IPG53]. 2004; <https://www.nice.org.uk/guidance/ipg53>. Accessed July 30, 2025.

72. National Institute for Health and Care Excellence (NICE). Percutaneous radiofrequency ablation for primary and secondary lung cancers [IPG372]. 2010; <https://www.nice.org.uk/guidance/ipg372>. Accessed July 29, 2025.
73. National Institute for Health and Care Excellence (NICE). Ultrasound-guided percutaneous radiofrequency ablation for benign thyroid nodules [IPG562]. 2016; <https://www.nice.org.uk/guidance/IPG562>. Accessed July 27, 2025.
74. Morris CS, Baerlocher MO, Dariushnia SR, et al. Society of Interventional Radiology Position Statement on the Role of Percutaneous Ablation in Renal Cell Carcinoma: Endorsed by the Canadian Association for Interventional Radiology and the Society of Interventional Oncology. *J Vasc Interv Radiol*. Feb 2020; 31(2): 189-194.e3. PMID 31917025

OTHER REFERENCES

1. Blue Cross and Blue Shield of Kansas Radiology Liaison Committee Consent Ballot; September 2015.
2. Blue Cross and Blue Shield of Kansas Urology Liaison Committee Consent Ballot, August 2016.
3. Blue Cross and Blue Shield of Kansas Radiology Liaison Committee, February 2009; February 2010; February 2011; February 2012; April 2016, July 2017, July 2023.
4. Blue Cross and Blue Shield of Kansas Urology Liaison Committee, June 2017, June 2023.