

## Medical Policy



### Title: Transcranial Magnetic Stimulation (TMS) as a Treatment of Depression and Other Psychiatric/Neurologic Disorders

Related Policies:	▪ <i>Vagus Nerve Stimulation</i>
-------------------	----------------------------------

#### Professional / Institutional

Original Effective Date: November 1, 2011

Latest Review Date: December 9, 2025

Current Effective Date: February 25, 2025

**State and Federal mandates and health plan member contract language, including specific provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage. To verify a member's benefits, contact [Blue Cross and Blue Shield of Kansas Customer Service](#).**

**The BCBSKS Medical Policies contained herein are for informational purposes and apply only to members who have health insurance through BCBSKS or who are covered by a self-insured group plan administered by BCBSKS. Medical Policy for FEP members is subject to FEP medical policy which may differ from BCBSKS Medical Policy.**

**The medical policies do not constitute medical advice or medical care. Treating health care providers are independent contractors and are neither employees nor agents of Blue Cross and Blue Shield of Kansas and are solely responsible for diagnosis, treatment and medical advice.**

**If your patient is covered under a different Blue Cross and Blue Shield plan, please refer to the Medical Policies of that plan.**

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"><li>• With treatment-resistant depression</li></ul>	Interventions of interest are: <ul style="list-style-type: none"><li>• Repetitive Transcranial magnetic stimulation</li></ul>	Comparators of interest are: <ul style="list-style-type: none"><li>• Pharmacotherapy</li><li>• Psychological and behavioral therapy</li><li>• Electroconvulsive therapy</li></ul>	Relevant outcomes include: <ul style="list-style-type: none"><li>• Symptoms</li><li>• Functional outcomes</li><li>• Quality of life</li></ul>

Populations	Interventions	Comparators	Outcomes
Individuals: <ul style="list-style-type: none"> <li>• With psychiatric or neurologic disorders other than depression, migraine or obsessive-compulsive disorder</li> </ul>	Interventions of interest are: <ul style="list-style-type: none"> <li>• Repetitive Transcranial magnetic stimulation</li> </ul>	Comparators of interest are: <ul style="list-style-type: none"> <li>• Pharmacotherapy</li> <li>• Therapy as appropriate including either physical and occupational therapy or psychological and behavioral therapy</li> </ul>	Relevant outcomes include: <ul style="list-style-type: none"> <li>• Symptoms</li> <li>• Functional outcomes</li> <li>• Quality of life</li> </ul>
Individuals: <ul style="list-style-type: none"> <li>• With obsessive-compulsive disorder</li> </ul>	Interventions of interest are: <ul style="list-style-type: none"> <li>• Repetitive Transcranial magnetic stimulation</li> </ul>	Comparators of interest are: <ul style="list-style-type: none"> <li>• Pharmacotherapy</li> <li>• Psychological and behavioral therapy</li> </ul>	Relevant outcomes include: <ul style="list-style-type: none"> <li>• Symptoms</li> <li>• Functional outcomes</li> <li>• Quality of life</li> </ul>
Individuals: <ul style="list-style-type: none"> <li>• With migraine</li> </ul>	Interventions of interest are: <ul style="list-style-type: none"> <li>• Repetitive Transcranial magnetic stimulation</li> </ul>	Comparators of interest are: <ul style="list-style-type: none"> <li>• Pharmacotherapy</li> </ul>	Relevant outcomes include: <ul style="list-style-type: none"> <li>• Symptoms</li> <li>• Functional outcomes</li> <li>• Quality of life</li> </ul>

## DESCRIPTION

Transcranial magnetic stimulation (TMS) is a noninvasive method of delivering electrical stimulation to the brain. The technique involves the placement of a small coil over the scalp and the passing of a rapidly alternating current through the coil wire. The electrical current produces a magnetic field that passes unimpeded through the scalp and bone and stimulates neuronal function. Repetitive TMS is being evaluated for the treatment of treatment-resistant depression (TRD) and other psychiatric and neurologic disorders. A variety of TMS modalities have been developed, which differ on parameters including stimulation intensity, frequency, pattern, and site of the brain stimulation. In conventional TMS, high frequency stimulation is delivered over the left dorsolateral prefrontal cortex (DLPFC) or low frequency stimulation over the right DLPFC. In bilateral TMS, both procedures are performed in the same session. Deep TMS employs an H-coil helmet designed to encompass a broader surface area and stimulate deeper brain structures than conventional TMS. Theta burst stimulation is administered at lower intensities and shorter intervals than conventional TMS.

## OBJECTIVE

The objective of this evidence review is to evaluate whether the use of repetitive transcranial magnetic stimulation of the brain improves the net health outcome for individuals with various psychiatric or neurologic conditions.

## BACKGROUND

### Transcranial Magnetic Stimulation

Transcranial magnetic stimulation (TMS), introduced in 1985 as a new method of noninvasive stimulation of the brain, involves placement of a small coil over the scalp, passing a rapidly alternating current through the coil wire, which produces a magnetic field that passes unimpeded

through the scalp and bone, resulting in electrical stimulation of the cortex. Transcranial magnetic stimulation was initially used to investigate nerve conduction (eg, TMS over the motor cortex will produce a contralateral muscular-evoked potential). The motor threshold, which is the minimum intensity of stimulation required to induce a motor response, is empirically determined for each person by localizing the site on the scalp for optimal stimulation of a hand muscle, then gradually increasing the intensity of stimulation. Interest in the use of TMS as a treatment for depression was augmented by the development of a device that could deliver rapid, repetitive stimulation. Imaging studies had shown a decrease in the activity of the left dorsolateral prefrontal cortex in depressed patients, and early studies suggested that high-frequency (eg, 5 to 10 Hz) TMS of the left dorsolateral prefrontal cortex had antidepressant effects. In contrast to electroconvulsive therapy (ECT), TMS does not require general anesthesia and does not generally induce a convulsion. Repetitive TMS (rTMS) is also being tested as a treatment for a variety of other psychiatric and neurologic disorders.

Conventional TMS delivers repeated electromagnetic pulses to induce prolonged modulation of neural activity, typically applied over the dorsolateral prefrontal cortex. High-frequency rTMS (usually  $\geq 10$  Hz) induces an increase in neural activity whereas low-frequency TMS (usually  $\leq 1$  Hz) has the opposite effect. If both procedures are performed in the same session, the intervention is described as bilateral rTMS.

A variety of TMS modalities have been developed, which differ on parameters including stimulation intensity, frequency, pattern, and site of the brain stimulation. Deep TMS employs an H-coil helmet design to encompass a broader surface area and stimulate deeper brain structures than conventional TMS. Theta burst stimulation is administered at lower intensities and shorter intervals than conventional rTMS.

## **REGULATORY STATUS**

Devices for transcranial stimulation have been cleared for marketing by the U.S. Food and Drug Administration (FDA) for diagnostic uses (FDA Product Code: GWF). A number of devices subsequently received FDA clearance for the treatment of major depressive disorder in adults who have failed to achieve satisfactory improvement from prior antidepressant medication in the current episode. Some of these devices use deep TMS or theta burst protocols. For example, the Brainsway Deep TMS system was FDA cleared for treatment-resistant depression in 2013 based on substantial equivalence to the Neurostar TMS Therapy System, and the Horizon (Magstim) and MagVita (Tonica Elektronik) have FDA clearance for their theta burst protocols.

Indications were expanded to include treating pain associated with certain migraine headaches in 2013, and obsessive-compulsive disorder in 2018.

In 2014, eNeura Therapeutics received 510(k) marketing clearance for the SpringTMS® for the treatment of migraine headaches. The device differs from the predicate Cerena™ TMS device with the addition of an LCD screen, a use authorization feature, a lithium battery pack, and a smaller size. The stimulation parameters are unchanged. The sTMS Mini (eNeura Therapeutics) received marketing clearance by the FDA in 2016. FDA product code: OKP.

In August 2018, the Deep TMS System (Brainsway) was granted a de novo 510(k) classification by the FDA as an adjunct for the treatment of adult patients with obsessive-compulsive disorder. The new classification applies to this device and substantially equivalent devices of this generic type.

The NeoPulse, now known as NeuroStar® TMS, was granted a de novo 510(k) classification by the FDA in 2008. The de novo 510(k) review process allows novel products with moderate or low-risk profiles and without predicates, which would ordinarily require premarket approval as a class III device, to be down-classified in an expedited manner and brought to market with a special control as a class II device.

In 2014, the Cerena™ TMS device (eNeura Therapeutics) was granted a de novo 510(k) classification by the FDA for the acute treatment of pain associated with migraine headache with aura. Warnings, precautions, and contraindications include the following:

- The device is only intended for patients experiencing the onset of pain associated with a migraine headache with aura.
- The device should not be used:
  - on headaches due to underlying pathology or trauma.
  - for medication overuse headaches.
- The device has not been demonstrated as safe and/or effective:
  - when treating cluster headache or a chronic migraine headache.
  - when treating during the aura phase.
  - in relieving the associated symptoms of a migraine (photophobia, phonophobia, and nausea).
  - in pregnant women, children under the age of 18, and adults over the age of 65.

The FDA has cleared multiple rTMS systems for adjunctive treatment of major depressive disorder in adolescents aged 15 to 21, including the NeuroStar Advanced Therapy System (K231926), the Magstim Horizon (K243869), and the MagVenture (K251125) stimulators.

Table 1 lists some devices that are FDA cleared for major depressive disorder (Product Code: OBP), migraine headache pain (Product Code: OKP), and obsessive-compulsive disorder (Product Code: QCI).

**Table 1. Repetitive Transcranial Magnetic Stimulation Devices Cleared by the U.S. Food and Drug Administration for Major Depression, Migraine, or Obsessive-Compulsive Disorder**

Device	Manufacturer	Indication	FDA Clearance No.	FDA Clearance Date
MagVenture TMS Therapy System	Tonica Elektronik	Major depressive disorder and obsessive-compulsive disorder	K193006	08/09/2020
Ultimate rTMS for OCD (M-series)	Brain Ultimate, Inc.	Major depressive disorder and obsessive-compulsive disorder	K230735	09/13/2023
CloudTMS Edge for OCD	TeleEMG, LLC	Obsessive-Compulsive Disorder	K233742	12/22/2023
Savi Dual™ Migraine Therapy	ENeura	Migraine (acute and prophylactic treatment in individuals ≥12 years of age)	K230358	05/16/2023
Horizon 3.0 TMS Therapy System	Magstim	Major depressive disorder and obsessive-compulsive disorder	K222171	01/13/2023
ALTMS Magnetic Stimulation Therapy System	REMEDI Co., Ltd	Major depressive disorder	K220625	04/06/2022
Neurostar	Neuronetics	Major Depressive Disorder	K083538	12/16/2008
		Obsessive-Compulsive Disorder	K212289	05/06/2022
Brainsway Deep TMS System	Brainsway	Major Depressive Disorder	K122288	01/07/2013
		Obsessive-Compulsive Disorder	K183303	03/08/2019
Springtms Total Migraine System	Eneura	Migraine headache with aura	K140094	05/21/2014

<b>Device</b>	<b>Manufacturer</b>	<b>Indication</b>	<b>FDA Clearance No.</b>	<b>FDA Clearance Date</b>
Rapid Therapy System	Magstim	Major Depressive Disorder	K143531	05/08/2015
Magvita	Tonica Elektronik	Major Depressive Disorder	K150641	07/31/2015
Mag Vita TMS Therapy System w/Theta Burst Stimulation	Tonica Elektronik	Major Depressive Disorder	K173620	8/14/2018
Neurosoft	TeleEMG	Major Depressive Disorder	K160309	12/22/2016
Horizon	Magstim	Major Depressive Disorder	K171051	09/13/2017
Horizon TMS Therapy System (Theta Burst Protocol)	Magstim	Major Depressive Disorder	K182853	03/15/2019
Nexstim	Nexstim	Major Depressive Disorder	K171902	11/10/2017
Apollo	Mag & More	Major Depressive Disorder	K180313	05/04/2018

## POLICY

- A. Repetitive transcranial magnetic stimulation (rTMS) of the brain using an FDA-cleared device and modality, which can include but is not limited to, conventional TMS, deep TMS, and theta burst stimulation (see Policy Guidelines) may be considered **medically necessary** as a treatment of major depressive disorder when **All** of the following conditions have been met:
1. Confirmed diagnosis of severe major depressive disorder (single or recurrent) documented by standardized rating scales that reliably measure depressive symptoms;  
**AND**
  2. Any one of the following (a, b, c, or d):
    - a. Individual has tried and had an inadequate response to 2 antidepressant agents from 2 different antidepressant classes (i.e., selective serotonin reuptake inhibitors, serotonin and norepinephrine reuptake inhibitors, tricyclic antidepressants, bupropion, or mirtazapine). An adequate trial of an antidepressant is defined by **BOTH** of the following:
      - i. The trial length was at least 6 weeks at generally accepted doses; **AND**
      - ii. Individual was  $\geq 80\%$  adherent to the agent during the trial.
    - b. Inability to tolerate a therapeutic dose of medications due to distinct side effects;  
**OR**
    - c. History of response to rTMS in a previous depressive episode (at least 3 months since the prior episode);  
**OR**
    - d. Is a candidate for electroconvulsive therapy (ECT) and ECT would not be clinically superior to rTMS (e.g., in cases with psychosis, acute suicidal risk, catatonia or life-threatening inanition rTMS should NOT be utilized).
- B. Repetitive TMS for major depressive disorder that does not meet the criteria listed above is considered **experimental / investigational**.
- C. Continued treatment with rTMS of the brain as maintenance therapy is considered **experimental / investigational**.
- D. Transcranial magnetic stimulation of the brain is considered **experimental / investigational** as a treatment of all other psychiatric/neurologic disorders including, but not limited to, bipolar disorder, schizophrenia, obsessive-compulsive disorder, or migraine headaches.

## POLICY GUIDELINES

- A. Repetitive Transcranial magnetic stimulation (TMS) should be performed using a U.S. Food and Drug Administration cleared device in appropriately selected individuals over age 18 years, by health care professionals who are adequately trained and experienced in the specific techniques used.
- B. The physician is responsible for the initial mapping (once per course or delivery) and the management of the treatment. The physician is also responsible for the re-determinations.

- C. A trained technician may perform the subsequent delivery of treatment.
- D. A variety of TMS modalities have been developed, which differ on parameters including stimulation intensity, frequency, pattern, and site of the brain stimulation.
- E. In conventional TMS, high frequency stimulation is delivered over the left dorsolateral prefrontal cortex (DLPFC) or low frequency stimulation over the right DLPFC. In bilateral TMS, both procedures are performed in the same session.
- F. Theta burst stimulation is administered at lower intensities and at shorter intervals than conventional TMS.
- G. Deep TMS employs an H-coil helmet designed to encompass a broader surface area and stimulate deeper brain structures than conventional TMS.
- H. A treatment course should not exceed 5 days a week for 6 weeks (total of 30 sessions), followed by a 3-week taper of 3 TMS treatments in week one, 2 TMS treatments the next week, and 1 TMS treatment in the last week.
- I. Theta burst stimulation may be administered using an accelerated protocol. One example of an accelerated theta burst protocol is the Stanford Accelerated Intelligent Neuromodulation Therapy (SAINT) protocol, consisting of 10 daily sessions over 5 consecutive days.
- J. Contraindications to repetitive TMS include:
  - 1. Seizure disorder or any history of seizure with increased risk of future seizure; or
  - 2. Presence of acute or chronic psychotic symptoms or disorders (e.g., schizophrenia, schizophreniform or schizoaffective disorder) in the current depressive episode; or
  - 3. Neurologic conditions that include epilepsy, cerebrovascular disease, dementia, increased intracranial pressure, having a history of repetitive or severe head trauma, or with primary or secondary tumors in the central nervous system; or
  - 4. Presence of an implanted magnetic-sensitive medical device located 30 centimeters or less from the TMS magnetic coil or other implanted metal items, including but not limited to a cochlear implant, implanted cardioverter defibrillator, pacemaker, vagus nerve stimulator, or metal aneurysm clips or coils, staples, or stents.
- K. The following should be present for the administration of repetitive TMS:
  - 1. An attendant trained in basic cardiac life support and the management of complications such as seizures, as well as the use of the equipment must be present at all times; and
  - 2. Adequate resuscitation equipment including, e.g., suction and oxygen; and
  - 3. The facility must maintain awareness of response times of emergency services (either fire/ambulance or "code team"), which should be available within 5 minutes. These relationships are reviewed on at least a 1-year basis and include mock drills.

**Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.**



## **RATIONALE**

This evidence review was created using searches of the PubMed database. The most recent literature update was performed through August 6, 2025.

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

## **TREATMENT-RESISTANT DEPRESSION**

### **Clinical Context and Therapy Purpose**

The purpose of repetitive transcranial magnetic stimulation (rTMS) is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with treatment-resistant depression (TRD).

The following PICO was used to select literature to inform this review.

### ***Populations***

The relevant population of interest is individuals with TRD.

### ***Interventions***

The therapy being considered is rTMS.

### ***Comparators***

The following therapies are currently being used to treat TRD: pharmacotherapy, psychological and behavioral therapy, and electroconvulsive therapy (ECT).

## Outcomes

The general outcomes of interest are reductions in symptoms and improvements in quality of life and functional outcomes.

**Table 2. Health Outcome Measures Relevant to Treatment-Resistant Depression, Major Depressive Disorder, Suicidal Ideation, and Suicidal Behavior**

Outcome	Description	Scale	Clinically Meaningful Difference
MADRS	<ul style="list-style-type: none"> <li>Physician scored</li> <li>Rates presence and severity of depression</li> <li>Symptom domains include sadness; pessimism; inability to feel; suicidality</li> </ul>	<ul style="list-style-type: none"> <li>Contains 10 items (scored from 0 to 6) with higher scores indicating more severe depression</li> <li>No validated cut-off score but generally 0 to 6 normal (no depression); 7 to 19 mild depression; 20 to 34 moderate depression; 35 to 59 severe depression; 60 or greater very severe depression<sup>1</sup>,</li> </ul>	<ul style="list-style-type: none"> <li>No consensus to define remission. Thresholds for remission have ranged from 6 to 12 in trials.</li> <li>One literature review reported that the mean weighted MADRS score for remission was 4.0 (95% CI, 3.5-4.5) based on 10 studies.<sup>2</sup> The definition of remission was a complete absence of clinically significant symptoms of depression.</li> <li>As per FDA, for drugs that have been approved to treat MDD as monotherapy or adjunctive treatment, treatment differences were typically closer to 3 or 4 points in MADRS scores. The observed treatment differences in esketamine studies were in that range.<sup>3</sup>,</li> </ul>
HAM-D	<ul style="list-style-type: none"> <li>Physician scored</li> <li>Rates presence and severity of depression</li> <li>Used in a number of registration studies of approved oral antidepressants</li> <li>Symptom domains include sadness; pessimism; inability to feel; suicidality</li> </ul>	<ul style="list-style-type: none"> <li>There are 2 versions: 17 or 25 items; 17 items is more common</li> <li>Each item scored in a range of 0 to 2 or 0 to 4, with higher scores indicating a greater degree of depression</li> <li>Scores range from 0 to 48</li> <li>Scores as low as 17 are associated with moderate depression and those at or above 24 are associated with severe depression<sup>2</sup>,</li> </ul>	<ul style="list-style-type: none"> <li>Remission is defined as total score of 7 or less. But 2 or less has been suggested as optimal.</li> <li>Response to treatment is defined as a 50% reduction from baseline scores.</li> </ul>
SIBAT	<ul style="list-style-type: none"> <li>Contains both patient- and clinician-reported modules and can be assessed by patient or rated by the physician</li> </ul>	<ul style="list-style-type: none"> <li>CGI-SS-r: rated from 0 (normal, not at all suicidal) to 6 (among the most extremely suicidal patients)</li> <li>CGI-SR-I: rates best clinical</li> </ul>	<ul style="list-style-type: none"> <li>No literature was identified for a consensus definition for a clinically meaningful change in scores</li> </ul>

Outcome	Description	Scale	Clinically Meaningful Difference
	<ul style="list-style-type: none"> <li>Includes assessments of <ul style="list-style-type: none"> <li>Severity of Suicidality (CGI-SS-r)</li> <li>Imminent Suicide Risk (CGI-SR-I)</li> <li>Frequency of Suicidal Thinking (FoST)<sup>4</sup>,</li> </ul> </li> </ul>	<p>judgment of participant's imminent risk for suicide within the next 7 days. Scale indicates: 0 (No imminent suicide risk), 1 (Minimal imminent), 2 (Mild imminent), 3 (Moderate imminent), 4 (Marked imminent), 5 (Severely imminent), 6 (Extreme imminent).</p> <ul style="list-style-type: none"> <li>FoST: describes the clinician determined estimate of the frequency of the participant's suicidal thinking. Scored on a 6-point Likert scale: 0 (Never), 1 (Rarely), 2 (Sometimes), 3 (Often), 4 (Most of the time), 5 (All of the time).<sup>4</sup>,</li> </ul>	

CGI-SR-I: Clinical Global Impression of Imminent Suicide Risk Scale, CGI-SS-r: Clinical Global Impression of Severity of Suicidality-Revised, CI: confidence interval; FDA: U.S. Food and Drug Administration; FoST: Frequency of Suicidal Thinking, HAM-D: Hamilton Rating Scale for Depression, MADRS: Montgomery-Asberg Depression Rating Scale, MDD: major depressive disorder; SIBAT: Suicide Ideation and Behavior Assessment Tool.

## Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

## Review of Evidence

Evaluation of rTMS for TRD includes RCTs comparing rTMS with sham as well as evidence when used as a replacement for or adjunct to pharmacotherapy that has not improved depressive symptoms. In addition, evaluation of rTMS in TRD includes the use of rTMS as an alternative to ECT. However, some individuals may not elect ECT due to its requirement for general anesthesia and induction of seizures.

There has been a trend to use rTMS at increased levels of intensity, trains of pulses, total pulses per session, and the number of sessions.<sup>5</sup> Unless otherwise indicated, stimulation was set at 100% to 120% of motor threshold, clinical response was defined as an improvement of 50% or more on the Hamilton Rating Scale for Depression (HAM-D), and remission was considered to be a score of 7 or less on the HAM-D. Refer to the meta-analysis by Schutter (2009) for a summary of study characteristics and stimulation parameters used in trials conducted prior to 2008.<sup>6</sup>

## Systematic Reviews

In pediatric major depressive disorder, Gallop et al identified 28 studies (n=640 participants), including 4 double-blind, sham-controlled RCTs, 1 additional trial which compared low-frequency vs. high-frequency rTMS, 6 open-label cohort studies, and multiple case reports.<sup>7</sup> The largest RCT (n=103) found no significant difference between active and sham rTMS on clinician-rated depression severity or remission rates, while another RCT (n=100) showed that rTMS combined with a selective serotonin reuptake inhibitor (SSRI) produced significantly greater reductions in HAM-D, children's depression rating scale-revised (CDRS-R) scores, and higher response rates than SSRI alone. A smaller sham-controlled RCT (n=55) reported reductions in depression severity and fewer self-injury impulses in non-medicated adolescents who received rTMS. Open-label cohorts (n=8 to 32) generally reported significant reductions in both clinician-rated and self-reported depression severity, with some effects maintained up to 6 months. A meta-analysis was not performed due to heterogeneity in study design, comparators, and stimulation protocols.

The Health Quality Ontario (2016) published a systematic review of left dorsolateral prefrontal cortex (DLPFC) rTMS for TRD.<sup>8</sup> Reviewers included 23 RCTs (n=1156 patients) that compared rTMS with sham and 6 RCTs (n=266 patients) that compared rTMS with ECT. In 16 studies, patients received rTMS in addition to antidepressant medication. Seven studies used intensities of less than 100% motor threshold and the definition of remission in the included studies varied (from  $\leq 7$  to  $\leq 10$  on the HAM-D). Meta-analysis showed a statistically significant improvement in depression scores compared with sham, with a weighted mean difference (WMD) of 2.31 (Table 3). However, this was smaller than the prespecified clinically important difference of 3.5 points on the HAM-D, and the effect size was small (0.33; 95% confidence interval [CI], 0.17 to 0.5;  $p < .001$ ). Subgroup analysis showed a larger and clinically significant treatment effect in the rTMS studies using 20 Hz with shorter train duration compared with other rTMS techniques (WMD, 4.96; 95% CI, 1.15 to 8.76;  $p = .011$ ). Secondary analyses showed rTMS demonstrated statistically greater response rates among 20 studies (pooled relative risk [RR], 1.72) as well as statistically greater remission rates among 13 studies (pooled RR, 2.20). For the 6 trials that compared rTMS with ECT, the WMD of 5.97 was both statistically and clinically significant in favor of ECT. The RR for remission and response rates are shown in Table 3, which while favoring ECT were not statistically significant. Remission and relapse rates at the 6-month follow-up were reported in 2 studies (n=40 and n=46 subjects) comparing rTMS with ECT. While 1 study reported a slightly higher remission rate for ECT (27.3%) than for rTMS (16.7%), the other study did not find a significant difference between ECT and rTMS for mean depression scores at 3 or 6 months, but did note relapses were less frequent for ECT. Statistical comparisons were either not significant or not available, limiting the interpretation of these findings.

**Table 3. Statistical Comparisons for Depression Scores after Repetitive Transcranial Magnetic Stimulation**

Comparison	Favors	WMD (95% CI)	p	RR for Remission (95% CI)	p	RR for Response (95% CI)	p
rTMS vs. sham	rTMS	2.31 (1.19 to 3.43)	<.001	2.20 (1.44 to 3.38)	.001	1.72 (1.13 to 2.62)	.01
rTMS vs. ECT	ECT	5.97 (0.94 to 11.0)	.02	1.44 (0.64 to 3.23)	.38	1.72 (0.95 to 3.11)	.07

CI: confidence interval; ECT: electroconvulsive therapy; rTMS: repetitive transcranial magnetic stimulation; RR: relative risk; WMD: weighted mean difference.

Brunoni et al (2017) conducted a systematic review to compare different modalities of rTMS for TRD.<sup>9</sup> Bilateral, high frequency rTMS, low-frequency rTMS, and theta burst stimulation were statistically significantly more effective than sham with respect to response (odds ratio [OR], 3.39 [95% CI, 1.91 to 6.02]; OR, 3.28 [95% CI, 2.33 to 4.61]; OR, 2.48 [95% CI, 1.22 to 5.05]; OR, 2.57 [95% CI, 1.17 to 5.62], respectively). In network meta-analysis, deep TMS was not more effective than sham TMS for response (OR, 1.49; 95% CI, 0.50 to 4.47) or remission (OR, 2.45; 95% CI, 0.74 to 8.07), but this result was based on only 1 RCT.

A systematic review conducted by Voigt et al (2021) focused on theta burst stimulation of TRD.<sup>10</sup> The reviewers included 8 RCTs comparing theta burst stimulation to sham treatment and 1 comparing theta burst stimulation to conventional rTMS. As measured by the HAM-D, theta burst stimulation was superior to sham on response (RR, 2.4; 95% CI, 1.27 to 4.55;  $p=.007$ ;  $I^2=40\%$ ). There was no statistically significant difference between theta burst stimulation and conventional rTMS (RR, 1.02; 95% CI, 0.85 to 1.23;  $p=.80$ ;  $I^2=0\%$ ). There was no difference between theta burst stimulation and rTMS in the incidence of adverse events.

### Randomized Controlled Trials

The ASCERTAIN-TRD study was a 3-arm, open-label study published by Papakostas et al (2024).<sup>11</sup> In this study, patients with TRD (failed  $\geq 2$  antidepressants) were randomized to augmentation with either aripiprazole or rTMS or were switched to venlafaxine XR. The study was open-label and limited to 8 weeks duration. A total of 235 individuals completed the study. MADRS response rates were numerically improved with rTMS augmentation compared with either aripiprazole augmentation or switching to venlafaxine/duloxetine (52.2% vs 38.1% and 35.8%, respectively). MADRS remission rates were also better among rTMS-treated individuals (34.2% vs 25.3% and 24.9%, respectively). MADRS score changes were significantly improved with rTMS (-17.39;  $p=.015$ ) compared with switch (-13.22); however, augmentation with aripiprazole (-14.9;  $p=.069$ ) was not significantly better than switch.

### Theta Burst Stimulation Compared to Conventional Transcranial Magnetic Stimulation

Blumberger et al (2018) published a multicenter, randomized, noninferiority trial, Conventional Versus Theta Burst Repetitive Transcranial Magnetic Stimulation in the Treatment of Major Depressive Disorder comparing 10-Hz rTMS with intermittent theta burst stimulation (iTBS).<sup>12</sup> Between 2013 and 2016, 414 patients with TRD were enrolled and randomized to 4 to 6 weeks of rTMS ( $n=205$ ) or iTBS ( $n=209$ ). Treatment resistance was defined as the failure to

tolerate 2 or more antidepressant trials of inadequate dose and duration or no clinical response to an adequate dose of an antidepressant. Patients who failed more than 3 antidepressant trials of adequate dosage were excluded from the trials. Patients could alter their medications during this trial. Treatment with conventional rTMS (37 minutes) and iTBS (3 minutes) was delivered 5 times a week for 4 to 6 weeks. The primary outcome measure was the 17-item HAM-D, for which scores for patients treated with rTMS improved by 10.1 points and scores for patients treated with iTBS improved by 10.2 points (adjusted difference, 0.103; lower 95% CI, -1.16;  $p=.001$ ). Treatment with iTBS resulted in a higher self-rated intensity of pain (mean score, 3.8) than treatment with rTMS (mean score, 3.4;  $p=.011$ ). Headache was the most common treatment-related adverse event for both groups (rTMS, 64% [131/204]; iTBS, 65% [136/208]). Serious adverse events were noted in patients treated with rTMS (1 case of myocardial infarction) and iTBS (1 case each of agitation, worsening suicidal ideation, worsening depression); there was no significant difference in the number of adverse events in the 2 groups. The trial lacked a treatment group with a placebo.

### **Deep Transcranial Magnetic Stimulation**

The RCT leading to 510(k) clearance of the Brainsway Deep TMS System in 2013 was conducted at 20 centers across the U.S. ( $n=13$ ), Israel ( $n=4$ ), Germany ( $n=2$ ), and Canada ( $n=1$ ).<sup>13</sup> The trial included 229 patients with major depressive disorder who had not received benefits from 1 to 4 antidepressant trials or were intolerant of at least 2 antidepressant treatments. Using a per-protocol analysis, which excluded 31 patients who did not receive adequate TMS treatment and 17 patients who did not meet the inclusion and exclusion criteria, the RCT showed a significant benefit for both response rate (38.4% vs. 21.4%) and remission rate (32.6% vs. 14.6%). A modified intention-to-treat analysis (ITT), which excluded the 17 patients not meeting selection criteria, showed a significant benefit in both response rate (37% vs. 22.8%) and remission rate (30.4% vs. 15.8%). At the end of the maintenance period (16-week follow-up), the response rate remained significantly improved for deep TMS. Remission rates were not reported. The ITT analysis found no significant benefit of treatment at 4 or 16 weeks.

## **DURABILITY OF CONVENTIONAL TRANSCRANIAL MAGNETIC STIMULATION**

### **Systematic Reviews**

Kedzior et al (2015) examined the durability of the antidepressant effect of high-frequency rTMS on the left DLPFC in the absence of maintenance treatment.<sup>14</sup> Included were 16 double-blind, sham-controlled, randomized trials ( $N=495$  patients). The range of follow-up was 1 to 16 weeks, but most studies only reported follow-up to 2 weeks. The overall effect size was small with a standardized mean difference (SMD; Cohen's  $d$ ) of -0.48, and the effect sizes were lower in RCTs with 8 to 16 weeks of follow-up ( $d=-0.42$ ) than with 1 to 4 weeks of follow-up ( $d=-0.54$ ). The effect size was larger when an antidepressant medication was initiated concurrently with rTMS (5 RCTs,  $d=-.56$ ) than when patients were on a stable dose of medication (9 RCTs,  $d=-0.43$ ) or were unmedicated (2 RCTs,  $d=-0.26$ ).

### **Observational Studies**

Dunner et al (2014) reported a 1-year follow-up with maintenance therapy from a large multicenter observational study (42 sites) of rTMS for patients with TRD.<sup>15</sup> A total of 257 patients agreed to participate in the follow-up study of 307 who were initially treated with rTMS. Of them, 205 completed the 12-month follow-up, and 120 patients had met the Inventory of Depressive

Symptoms-Self Report response or remission criteria at the end of treatment. Ninety-three (36.2%) of the 257 patients who enrolled in the follow-up study received additional rTMS (mean, 16.2 sessions). Seventy-five (62.5%) of the 120 patients who met response or remission criteria at the end of the initial treatment phase (including a 2-month taper phase) continued to meet response criteria through a 1-year follow-up.

A variety of tapering schedules are being studied. For example, Richieri et al (2013) used propensity-adjusted analysis of observational data and found that patients who had rTMS tapered over 20 weeks (from 3 times per week to once a month) had a significantly reduced relapse rate than patients who had no additional treatment (37.8% vs. 81.8%).<sup>16</sup> Connolly et al (2012) reported that in the first 100 cases treated at their institution, the response rate was 50.6% and the remission rate was 24.7%.<sup>17</sup> At 6 months after the initial rTMS treatment, 26 (62%) of 42 patients who received tapered maintenance therapy (from 2 sessions per week for the first 3 weeks to monthly) maintained their response. In another study, Janicak et al (2010) evaluated patients who met criteria for a partial response during either a sham-controlled or an open-label phase of a prior study were tapered from rTMS and simultaneously started on maintenance antidepressant monotherapy.<sup>18</sup> During the 24-week follow-up, 10 of 99 patients relapsed, 38 had symptom worsening, and of these 32 (84%) had symptomatic benefit with adjunctive rTMS.

### **Section Summary: Treatment-Resistant Depression**

There are a large number of sham-controlled randomized trials and meta-analyses of these RCTs evaluating the use of rTMS for depression. Meta-analyses found improved response rates and rates of remission for conventional rTMS and theta burst stimulation compared with sham TMS. Additionally, a head-to-head trial showed noninferiority of theta burst stimulation to conventional rTMS, with no difference in the incidence of adverse events. There is some evidence that rTMS, when given in conjunction with the initiation of pharmacologic therapy, improves the response rate compared with pharmacologic therapy alone, while the effect of rTMS is less robust when it is given in combination with a stable dose of antidepressant medication. There is limited evidence to compare the effects of these treatments on cognition, although the adverse events of rTMS appear to be minimal. While the most recent meta-analyses have found that the effect of rTMS is smaller than the effect of ECT on TRD, given that rTMS does not require general anesthesia or induce seizures and some individuals may not elect ECT, the balance of incremental benefits and harms associated with rTMS may be reasonable compared with ECT.

## **MIGRAINE HEADACHE**

### **Clinical Context and Therapy Purpose**

The purpose of rTMS is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with migraine headache pain.

The following PICO was used to select literature to inform this review.

### ***Populations***

The relevant population of interest is individuals with migraine headaches.

### ***Interventions***

The therapy being considered is rTMS.

### Comparators

The following therapies are currently being used to treat migraine headache pain: pharmacotherapy (e.g., triptans, ibuprofen, combination analgesics).

### Outcomes

The general outcomes of interest are reductions in symptoms and improvements in quality of life and functional outcomes.

### Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

## REVIEW OF EVIDENCE

### Systematic Review

Saltychev et al (2022) conducted a systematic review and meta-analysis of 8 RCTs that compared rTMS to sham stimulation in patients with migraine.<sup>19</sup> All RCTs used high-frequency rTMS to the left dorsolateral prefrontal cortex, and all studies except 1 included patients with chronic migraine. All studies except 1 had a low risk of bias and the risk of publication bias was nonsignificant. Results for the frequency of migraine days per month and the intensity of migraine pain both favored rTMS; however, the authors stated that the difference in migraine pain intensity was clinically insignificant. The analysis is summarized in Tables 4 and 5.

**Table 4. Systematic Review & Meta-Analysis Characteristics**

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Saltychev et al (2022) <sup>19</sup> ,	2004-2021	8	Adults with migraine	339 (11 to 100)	RCTs	3 to 12 rTMS sessions over 3 days to 8 weeks

**Table 5. Systematic Review & Meta-Analysis Results**

Study	Migraine days per month	Migraine pain (scale of 0 to 100)
Saltychev et al (2022) <sup>19</sup> ,		
N=339	N=339	N=257
Difference (95% CI)	-8.09 (-11.4 to -4.79)	-13.56 (-21.8 to -5.32)
<i>I</i> <sup>2</sup>	<i>I</i> <sup>2</sup> =87%	<i>I</i> <sup>2</sup> =86%

CI: confidence interval



### **Randomized Controlled Trial**

A pivotal randomized, double-blind, multicenter, sham-controlled trial was performed with the Cerena TMS device to demonstrate the safety and effectiveness of a de novo application.<sup>20</sup> Enrolled in the trial were 201 patients with a history of an aura preceding more than 30% of headaches of moderate or severe severity for approximately 90% of migraine attacks. Following a month-long baseline phase to establish the frequency and severity of the migraine, patients were randomized to a treatment phase consisting of 3 treatments or 3 months, whichever occurred first. Patients were instructed to treat their migraine headache during the aura phase and to record their pain severity (0 to 3), severity of associated migraine symptoms (photophobia, phonophobia, nausea), presence of vomiting, and use of rescue medications at the time of treatment and at 1, 2, 24, and 48 hours after treatment. The primary endpoint was the proportion of patients who were pain-free 2 hours after treatment. Of the 201 patients enrolled, 164 recorded at least 1 treatment and 113 recorded at least 1 treatment when there was pain. Post hoc analysis of these 113 patients showed a benefit of the device for the primary endpoint (37.74% pain free after 2 hours for Cerena vs. 16.67% for sham;  $p=.018$ ) and for the proportion of subjects who were pain free after 24 hours (33.96% for Cerena vs. 10% for sham;  $p=.002$ ). Active treatment was not inferior to sham for the proportion of subjects free of photophobia, suggesting that the device does not worsen photophobia. However, the device was not inferior to sham for the proportion of subjects free of nausea and phonophobia.

### **Section Summary: Migraine Headache**

The available evidence on the use of TMS devices to treat migraine include a systematic review and a pivotal RCT. The systematic review found that rTMS reduced migraine pain and intensity compared to sham. The results of the pivotal trial were limited by the 46% dropout rate and post hoc analysis. According to the FDA labeling, the device has not been demonstrated as safe or effective when treating cluster headache, chronic migraine headache, or migraine headache during the aura phase. The device has not been demonstrated to be as effective in relieving the associated symptoms of migraine (photophobia, phonophobia, nausea).

## **OBSESSIVE-COMPULSIVE DISORDER**

### **Clinical Context and Therapy Purpose**

The purpose of rTMS is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with obsessive-compulsive disorder (OCD).

The following PICO was used to select literature to inform this review.

### ***Populations***

The relevant population of interest is individuals with OCD.

OCD is characterized by the inability to suppress intrusive thoughts, impulses, images, and repetitive motor responses.

### ***Interventions***

The therapy being considered is rTMS.

The use of TMS for patients with OCD is based on the observation that OCD symptoms are associated with excessive activity in certain cortical areas. Transcranial magnetic stimulation is proposed as a treatment to modulate these brain areas.

### **Comparators**

The following therapies are currently being used to treat OCD: pharmacotherapy, psychological, and behavioral therapy.

### **Outcomes**

The general outcomes of interest are reductions in symptoms and improvements in quality of life and functional outcomes.

The Yale-Brown Obsessive Compulsive Scale (YBOCS) is a clinician-rated, 10-item scale commonly used to assess the severity of symptoms in OCD.<sup>21</sup> Each item is rated from 0 (no symptoms) to 4 (extreme symptoms) (total range, 0 to 40), with separate subtotals for the severity of obsessions and compulsions.

YBOCS scores of 0 to 13 correspond to 'mild symptoms' on the Clinical Global Impression of Severity (CGI-Severity=0 to 2), 14 to 25 with 'moderate symptoms' (CGI-Severity=3), 26 to 34 with 'moderate-severe symptoms' (CGI-Severity=4) and 35 to 40 with 'severe symptoms' (CGI-Severity=5 to 6).<sup>22</sup> An improvement of  $\geq 35\%$  on the YBOCS is most predictive of treatment response.<sup>23</sup>

Follow-up over months is of interest to monitor outcomes.

### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

## **REVIEW OF EVIDENCE**

### **Systematic Reviews**

A systematic review by Trevizol et al (2016) included 15 RCTs (N=483) that compared active with sham rTMS for OCD (Tables 6 and 7).<sup>24</sup> All studies were sham-controlled and double-blind. The sample sizes in the trials ranged from 18 to 65 patients. Seven studies used low-frequency stimulation and 8 studies used high-frequency stimulation. The cortical regions varied among the studies, targeting the supplementary motor area, orbitofrontal cortex, or left, right, or bilateral DLPFC. The researchers calculated the SMD for the primary outcome (YBOCS score). Response rates were not reported.

The pooled mean difference between groups on the YBOCS was 2.94 (95% CI, 1.26 to 4.62), translating to a small to moderate effect size for active stimulation of 0.45 (95% CI, 0.20 to 0.71). Individual adverse effects were not assessed due to a lack of reporting in the primary studies, but there was no difference between groups in the dropout rate. Intervention protocols were heterogeneous across the studies, but regression analysis did not identify any treatment protocol or other variables as predictors of TMS response.

More recently, Liang et al (2021) conducted a systematic review and meta-analysis of different TMS modalities for the treatment of OCD.<sup>25</sup> Three of the 5 protocols assessed were significantly more efficacious than sham TMS, and all treatment strategies were similar to sham TMS regarding tolerability (Table 7). Transcranial magnetic stimulation was not more effective than sham TMS, but there was direct evidence from only 1 RCT for this comparison (Carmi et al, 2019, discussed in the next section).<sup>26</sup> The overall quality of the evidence was rated very low for efficacy and low for tolerability, and the reviewers concluded that high quality RCTs with low selection and performance bias are needed to further verify the efficacy of specific rTMS strategies for OCD treatment.

Perera et al (2021) conducted a systematic review and meta-analysis of rTMS in the treatment of OCD.<sup>27</sup> All RCTs in the analysis (n=26) had a low risk of bias. A random effects model was used to compare pre- and post-stimulation YBOCS scores, with effect sizes reported as Hedges' *g*. The analysis found that rTMS had a significant effect on YBOCS scores compared to sham (effect size, 0.64; 95% CI, 0.39 to 0.89; *p*<.0001). Raw mean difference in YBOCS score between treatments was 4.04 (95% CI, 2.54 to 5.54; *p*<.001). The effect size was still significant when 2 dominant trials were removed. Effect sizes with rTMS appeared to be significant until 4 weeks after treatment, and low- and high-frequency rTMS had similar efficacy to each other. The authors performed several subgroup analyses (cortical target, stimulation frequency, total pulses per session, total duration of treatment) but none of the effect sizes were significant between rTMS and sham.

**Table 6. Systematic Review of Transcranial Magnetic Stimulation in Patients with Obsessive-Compulsive Disorder: Characteristics**

Study	Dates	Trials Included	Participants	N (Range)	Design	Duration
Perera et al (2021) <sup>27</sup> ,	Up to October 2020	26	Mean age, 33 years	781	RCT, sham-controlled	1 week to 6 weeks
Liang et al (2021) <sup>25</sup> ,	Up to March 2020	22	Mean age, 34.1 years	698	RCT, sham- or active-controlled	1 week to 10 weeks
Trevizol et al (2016) <sup>24</sup> ,	Up to March 2016	15	Mean age, 31.9 (SD, 7.6) years, 44.1% women	483 (18 to 65); mean, 16.1 (SD, 8.45)	RCT, sham-controlled	1 week to 6 weeks

OCD: obsessive-compulsive disorder; RCT: randomized controlled trial; SD: standard deviation.

**Table 7 Systematic Review and Meta-Analysis: Results**

Study	YBOCS Score	Dropouts
Perera et al (2021) <sup>27,</sup>		
Total N	781	781
	Mean difference (95% CI)	
Active rTMS	4.04 (2.54 to 5.54)	NR
	$I^2=62.06\%$ ; $p<.0001$	
Liang et al (2021) <sup>25,</sup>		
	Mean Difference (95% CrI)	OR (95% CrI)
Low frequency rTMS applied over the dorsolateral prefrontal cortex	6.34 (2.12 to 10.42)	0.81 (0.08 to 8.17)
High-frequency rTMS applied over the dorsolateral prefrontal cortex	3.75 (1.04 to 6.81)	1.08 (0.37 to 3.19)
Low-frequency rTMS applied over the supplementary motor area	4.18 (0.83 to 7.62)	0.98 (0.37 to 2.67)
Low-frequency rTMS applied over the orbitofrontal cortex	4.43 (-2.57 to 11.31)	0.59 (0.06 to 5.68)
High-frequency rTMS applied over the anterior cingulate cortex/medial prefrontal cortex (deep TMS)	4.25 (-1.16 to 9.59)	1.62 (0.26 to 15.98)
Trevizol et al (2016) <sup>24,</sup>		
Total N	483	483
	SMD: 0.45 (0.20 to 0.71)	OR: 1.02 (0.76 to 1.36)
	$I^2=43\%$ ; $p=.039$	
	Mean Difference: 2.94 (1.26 to 4.62)	
	$I^2=58\%$ ; $p=.002$	

CI: confidence interval; CrI: credible interval; NR: not reported; OR: odds ratio; rTMS: repetitive transcranial magnetic stimulation; SMD: standardized mean difference; YBOCS: Yale-Brown Obsessive-Compulsive Scale.

### Randomized Controlled Trial

This section discusses in detail the sham-controlled RCT of deep TMS for OCD conducted by Carmi et al (2019).<sup>26,</sup> The trial was submitted to the FDA as part of the de novo classification request, to establish a reasonable assurance of safety and effectiveness of the device.<sup>28,</sup> Study characteristics and results are summarized in Tables 8 and 9, and limitations are shown in Tables 10 and 11. A total of 99 patients were randomized to active treatment or sham. The primary outcome was the difference between groups in the mean change from baseline to 6 weeks on

the YBOCS. Secondary outcomes included the response rate (defined as a 30% or greater improvement from baseline on the YBOCS), the Clinical Global Impression of Improvement (CGI-I), the Clinical Global Impression of Severity (CGI-S), and the Sheehan Disability Scale, a patient-reported measure of disability and impairment. Results at 10 weeks were also reported as secondary outcomes.

The primary efficacy analysis used a modified ITT analysis (n=94), excluding 5 patients who were found to not meet eligibility criteria following randomization. There was a greater decrease from baseline in the active treatment group (-6.0 points) than the sham group (-2.8 points), translating to a moderate effect size of 0.69. At 6 weeks, the response rate was 38.1% in the active treatment group compared to 11.1% in the sham group (p=.003). The FDA review provides data from the ITT analysis of the mean change in the YBOCS score (n=99). In the ITT data set, the YBOCS score decreased by -6.0 points (95% CI, -3.8 to -8.2) in the active group and by -4.1 points (95% CI, -1.9 to -6.2) in the sham group. Although the decreases were both statistically significant from baseline, the difference of 1.9 points between the treatment arms was not statistically significant (p=.0988). Results on the secondary outcomes were mixed. More patients in the active treatment group were considered improved based on the CGI-I and the CGI-S at 6 weeks, but there was no significant difference between groups on the Sheehan Disability Scale (Table 10).

**Table 8. Summary of Key Randomized Controlled Trial Characteristics - Transcranial Magnetic Stimulation for Patients with Obsessive-Compulsive Disorder**

Study; Trial	Countries	Sites	Dates	Participants	Interventions	Duration of follow-up
Carmi et al (2019) <sup>26</sup> , NCT02229903	U.S., Israel, Canada	11	2014-2017	N=99 adults ages 22 to 68 years, diagnosis of OCD as a primary disorder, receiving treatment in an outpatient setting, and having a YBOCS score >20; in maintenance treatment with a therapeutic dosage of a SRI for at least 2 months before randomization or, if they were not on an SRI, in maintenance treatment on CBT and have failed to respond adequately to at least 1 past trial	Deep TMS 6-week treatment phase (consisting of 5 weeks of daily treatments 5 days a week and 4 treatments during the 6th week)	Sham 6 weeks (primary) 10 weeks (secondary)

Study; Trial	Countries	Sites	Dates	Participants	Interventions	Duration of follow-up
				of an SRI. Exclusions: primary axis I diagnosis other than OCD, severe neurological impairment, any condition associated with an increased risk of seizures.		

CBT: cognitive behavioral therapy; OCD: obsessive-compulsive disorder; SRI: serotonin reuptake inhibitor; TMS: transcranial magnetic stimulation; YBOCS: Yale-Brown Obsessive-Compulsive Scale.

**Table 9 Summary of Key Randomized Controlled Trial Results - Transcranial Magnetic Stimulation for Patients with Obsessive-Compulsive Disorder**

Study	YBOCS (Primary Outcome)	YBOCS Response	CGI-I	CGI-S (modified)	Sheehan Disability Scale	Adverse events (all)	Dropouts
Carmi et al (2019) <sup>26</sup> ; NCT02229903	Mean change from baseline at 6 weeks	(>30% change from baseline to 6 weeks)	Moderate to very much improved from baseline at 6 weeks				
N analyzed	94	94	94	94			
TMS	-6.0 points (95% CI, 4.0 to 8.1)	38.1% (16/42)	20/41 (49%)	25/41 (61%)	-3.8 points (95% CI, -1.5 to -6.1)	73%	6/48 (12.5%)
Sham	-3.3 points (95% CI, 1.2 to 5.3)	11.1% (5/45)	9/43 (21%)	14/43 (32.6%)	-3.0 points (95% CI, -0.8 to -5.3)	69%	6/51 (12.0%)
Difference; P-value	2.8 points; p=.01 Effect size: 0.69	p=.003	p=.011	p=.022	NS (p-value not reported)	p=.639	NS (p-value not reported)

CGI-I: Clinical Global Impression of Improvement; CGI-S: Clinical Global Impression of Severity; CI: confidence interval; NS: non-significant; TMS: transcranial magnetic stimulation; YBOCS: Yale-Brown Obsessive-Compulsive Scale.

**Table 10. Study Relevance Limitations**

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Follow-Up <sup>e</sup>
Carmi et al (2019) <sup>26</sup> ; NCT02229903					1,2. 6 weeks (primary)

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

<sup>b</sup> Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

<sup>c</sup> Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

<sup>d</sup> Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

<sup>e</sup> Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

**Table 11. Study Design and Conduct Limitations**

Study	Allocation <sup>a</sup>	Blinding <sup>b</sup>	Selective Reporting <sup>c</sup>	Data Completeness <sup>d</sup>	Power <sup>e</sup>	Statistical <sup>f</sup>
Carmi et al (2019) <sup>26</sup> ; NCT02229903				6. Modified ITT analysis of 94/100 patients who were enrolled. The difference in the primary outcome was not statistically significant in the ITT data set (n=99)		

ITT: intention-to-treat.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

<sup>b</sup> Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

<sup>c</sup> Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>d</sup> Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

<sup>e</sup> Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

<sup>f</sup> Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

## Section Summary: Obsessive-Compulsive Disorder

The evidence on rTMS for OCD includes a number of small-to-moderate sized, sham-controlled, double-blind randomized trials and meta-analyses of these RCTs. A meta-analysis of 15 RCTs

(N=483 patients, range 18 to 65 patients) conducted in 2016 found a benefit of rTMS on patient-reported OCD symptom severity at time points ranging from 2 to 6 weeks, but there was substantial variability in the stimulation parameters, including the cortical region that was stimulated and the frequency of stimulation. The largest meta-analysis conducted in 2021 included 26 RCTs. Differences in pre- and posttreatment YBOCS scores were significantly improved with rTMS compared to sham, but the evidence was only sufficient to conclude that the effects lasted until 4 weeks after the last treatment. In a network meta-analysis that included both direct and indirect evidence, the authors did not find that deep TMS was more effective than sham rTMS. The RCT that was the basis of FDA clearance of deep TMS for treatment of OCD compared deep rTMS to sham in 99 patients for 6 weeks, with an additional 4 weeks of follow-up as a secondary outcome. Using a modified ITT analysis (n=94), there was a larger mean decrease from baseline (improvement) on the YBOCS score (the primary efficacy outcome) in the active treatment group (-6.0 points) than the sham group (-2.8 points), translating to a moderate effect size of 0.69. At 6 weeks, the response rate was 38.1% in the active treatment group compared to 11.1% in the sham group (p=.003), as measured by a 30% or greater increase in the YBOCS. The difference in the primary outcome measure between active and sham groups was not statistically significant in the ITT analysis. There was a benefit for rTMS on clinician-reported measures of improvement, but no significant difference between groups on patient-reported disability and impairment. Additional trials with sufficient sample size and follow-up duration are needed to confirm these results.

## **PSYCHIATRIC OR NEUROLOGIC DISORDERS OTHER THAN DEPRESSION, MIGRAINE, OR OBSESSIVE-COMPULSIVE DISORDER**

### **Clinical Context and Therapy Purpose**

The purpose of rTMS is to provide a treatment option that is an alternative to or an improvement on existing therapies in individuals with psychiatric or neurologic disorders other than depression, migraine, or OCD.

The following PICO was used to select literature to inform this review.

### ***Populations***

The relevant population of interest is individuals with psychiatric or neurologic disorders other than depression, migraine, or OCD.

### ***Interventions***

The therapy being considered is rTMS.

### ***Comparators***

The following therapies are currently being used to treat psychiatric disorders other than depression or OCD: pharmacotherapy or psychological and behavioral therapy. The following therapies are currently being used to treat neurologic disorders other than migraine: pharmacotherapy and therapy as appropriate including either physical or occupational therapy.

### ***Outcomes***

The general outcomes of interest are reductions in symptoms and improvements in quality of life and functional outcomes.



Follow-up over months is of interest to monitor outcomes.

### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

## **REVIEW OF EVIDENCE**

### **PSYCHIATRIC DISORDERS OTHER THAN DEPRESSION OR OBSESSIVE-COMPULSIVE DISORDER**

#### **BIPOLAR DISORDER**

##### **Systematic Reviews**

Konstantinou et al (2022) conducted a systematic review of 31 RCTs of rTMS for the treatment of bipolar disorder; meta-analysis was not performed.<sup>29</sup> Most included studies were in the setting of bipolar depression (n=24). Only 8 studies had a low risk of bias. Overall, rTMS seems safe and well-tolerated but efficacy results are mixed and there is no consensus about the optimal rTMS regimen. The authors noted limitations of the available literature including heterogeneity among studies, differences in sham treatments, and small sample sizes. They also stated that adequately powered sham-controlled studies are needed to verify the efficacy of rTMS in patients with bipolar disorder.

Tee et al (2020) conducted a systematic review and meta-analysis of sham-controlled RCTs of rTMS for the treatment of bipolar disorder.<sup>30</sup> Eight trials of rTMS in bipolar depression showed small but statistically significant improvements in depression scores compared to sham control (SMD, 0.302 ;  $p < .05$ ). However, most studies had a high risk of bias, which could have exaggerated the treatment effects. The effect of rTMS was inconclusive in bipolar mania due to the high heterogeneity and limited number of controlled trials.

#### **GENERALIZED ANXIETY DISORDER**

##### **Systematic Review**

Qi et al (2024) included 7 RCTs (N=298) of rTMS for the treatment of generalized anxiety disorder in a systematic review and meta-analysis.<sup>31</sup> Results showed that rTMS significantly reduced anxiety symptoms as measured by the Hamilton Anxiety Rating Scale (SMD, -1.17; 95% CI, -2.19 to -0.15). Response rates were higher in rTMS groups (RR, 0.61; 95% CI, 0.39 to 0.96), although remission did not reach statistical significance (RR, 0.73; 95% CI, 0.50 to 1.07). Headaches were more common in the intervention groups, but no differences in discontinuation were observed. The conclusions were limited by sample sizes and significant heterogeneity.

Cui et al (2019) included 21 studies (N=1481 patients) in a meta-analysis of rTMS plus drug therapy compared to drug therapy alone for the treatment of generalized anxiety disorder.<sup>32</sup> Results of the analysis showed that rTMS improved anxiety symptoms as measured by the Hamilton Anxiety Scale (SMD, -0.68 ; 95% CI, -0.89 to -0.46). The conclusions that could be drawn from the body of evidence were limited by significant heterogeneity across studies, and the authors concluded that additional high-quality studies are needed to confirm the results.

## **PANIC DISORDER**

### **Systematic Review**

A Cochrane review by Li et al (2014) identified 2 RCTs (N=40 patients) that compared low-frequency rTMS with sham rTMS over the right DLPFC.<sup>33</sup> The larger of the 2 studies was a randomized, double-blind, sham-controlled trial by Mantovani et al (2013) who assessed 21 patients with panic disorder and comorbid major depression.<sup>34</sup> The response was defined as a 40% or greater decrease on the Panic Disorder Severity Scale and a 50% or greater decrease in HAM-D scores. After 4 weeks of treatment, the response rate for panic was 50% with active rTMS and 8% with sham. The trial had a high-risk of attrition bias. The overall quality of evidence for the 2 trials was considered low, and the sample sizes were small, precluding certainty in the conclusions about the efficacy of rTMS for panic disorder.

## **POSTTRAUMATIC STRESS DISORDER**

### **Systematic Review**

Trevizol et al (2016) published a systematic review on the efficacy of low- and high-frequency rTMS for posttraumatic stress disorder.<sup>35</sup> Five sham-controlled randomized trials (N=118 patients) were included. Most trials used stimulation of the right DLPFC, though some delivered rTMS to the left DLPFC or bilaterally. Three trials used high-frequency stimulation while 1 used low-frequency stimulation and another compared high- with low-frequency stimulation; the percent motor threshold ranged from 80% to 120%. Some trials provided rTMS in combination with a scripted narrative of the traumatic event, and different posttraumatic stress disorder scales were used. In a meta-analysis, active rTMS was found to be superior to sham (SMD, 0.74; 95% CI, 0.06 to 1.42), although heterogeneity across the trials was high.

Brown et al (2024) published a Cochrane systematic review of rTMS for posttraumatic stress disorder.<sup>36</sup> The search was conducted through January 2023, and the authors identified 13 RCTs (N=577). Notably, 5 studies were conducted in the US, primarily enrolling white, male veterans. The authors found that rTMS probably makes little to no difference on posttraumatic stress disorder severity immediately following treatment (SMD, -0.14; 95% CI, -0.54 to 0.27; 3 studies, 99 participants; moderate-certainty evidence); however, there was significant heterogeneity amongst the studies.

## **SCHIZOPHRENIA**

### **Systematic Reviews**

Blyth et al (2025) included 126 studies (N=4122 participants) in a meta-analysis of rTMS safety in schizophrenia.<sup>37</sup> Results showed that active rTMS was associated with higher rates of

headache or scalp pain, dizziness or syncope, facial twitching, and nausea compared to sham ( $p < .05$ ). The prevalence of seizure was low (1.39 per 1000) and not different from sham or from the general population. No increased risk of cognitive impairment, worsening psychosis, depression, or mania was observed.

He et al (2017) published a meta-analysis of the effects of 1-Hz (low frequency) and 10-Hz (high frequency) rTMS for auditory hallucinations and negative symptoms of schizophrenia, respectively.<sup>38</sup> For 1-Hz rTMS, 13 studies were included. Compared with sham, the rTMS group showed greater improvement in auditory hallucinations (SMD, -0.29; 95% CI, -0.57 to -0.01). However, significant heterogeneity across the studies was found ( $p = .06$ ). In the 7 studies using 10-Hz rTMS, the overall effect size for improvement in negative symptoms was -0.41 (95% CI, -1.16 to -0.35); again, there was significant heterogeneity across studies ( $p < .001$ ). The review was further limited by the small number of articles included and by the lack of original data for some studies.

A Cochrane review by Dougall et al (2015) selected 41 studies (N=1473 participants).<sup>39</sup> Based on very low-quality evidence, there was a significant benefit of low- and high-frequency temporoparietal TMS compared with sham for the global state (7 RCTs) and positive symptoms (5 RCTs). For prefrontal rTMS compared with sham, the evidence on global and cognitive state was of very low-quality and equivocal. Reviewers concluded that the evidence was insufficient to support or refute the use of TMS to treat symptoms of schizophrenia and, although some evidence suggested that temporoparietal TMS might improve certain symptoms (eg, auditory hallucinations, positive symptoms of schizophrenia), the results were not sufficiently robust to provide certainty.

### Randomized Controlled Trials

Several additional small, single center RCTs of rTMS for the treatment of schizophrenia have been published since the systematic reviews described above (Tables 12 and 13).<sup>40,41,42,43,44,45,46</sup> These studies were limited by their small sample sizes, very high loss to follow-up, and inadequate duration of follow-up (Tables 14 and 15). Due to these limitations, these studies do not provide sufficient evidence to draw conclusions about the effectiveness of the technology in patients with schizophrenia.

**Table 12. Summary of Key Randomized Controlled Trial Characteristics**

Study; Trial	Countries	Sites	Dates	Participants	Interventions <sup>1</sup>		Duration of follow- up
					Active	Comparator	
Ye et al (2024) <sup>40</sup>	China	1	2016- 2019	Inpatient male veterans diagnosed with schizophrenia per DSM-V criteria who were clinically stable on antipsychotic	Neuronavigation- guided high- frequency rTMS over left DLPFC: 10 Hz (1200 pulses per session) or 20 Hz (1600 pulses per	Sham rTMS (n=28)	24 weeks

Study; Trial	Countries	Sites	Dates	Participants	Interventions <sup>1</sup>		Duration of follow- up
				therapy ≥24 weeks (N=84 randomized)	session) administered 5 times a week (Monday to Friday) for 6 weeks (n=56 total; 28 per group)		
Hua et al (2024) <sup>41,</sup>	China	1	2016-2021	Adults ages 18 to 60 diagnosed with schizophrenia per DSM-IV criteria and medication-resistant auditory verbal hallucinations who were clinically stable on antipsychotic therapy ≥2 weeks (N=66 randomized; 62 completed)	Imaging-navigated rTMS over left temporoparietal junction administered as 3 sessions per day for 14 consecutive days (n=32)	Sham rTMS (n=30)	6 weeks
Xiu et al (2024) <sup>42,</sup>	China	1	Not reported	Adults diagnosed with schizophrenia per DSM-IV criteria and persistent daily auditory verbal hallucinations despite ≥2 antipsychotics (N=120 randomized; 89 completed)	Neuronavigation-guided high-frequency rTMS over left DLPFC: 10 Hz (1,200 pulses per session) or 20 Hz (1,600 pulses per session) administered 5 times a week (Monday–Friday) for 8 weeks. (n=80; 40 in each group)	Sham rTMS (n=25)	32 weeks
Zhu et al (2021) <sup>46,</sup>	China	7	2017-2018	Inpatients ages 18 to 50 years with a diagnosis of schizophrenia per ICD-10 criteria who were	Intermittent theta burst stimulation over the cerebellum (3 pulses at 50 Hz repeated at a	Sham intermittent theta burst stimulation (n=32)	24 weeks

Study; Trial	Countries	Sites	Dates	Participants	Interventions <sup>1</sup>	Duration of follow- up
				right-handed and clinically stable for the past 3 months (N=64)	rate of 5 Hx for a total of 600 pulses administered 5 times a week (Monday to Friday) for 2 weeks (n=32)	
Guan et al (2020) <sup>43,</sup>	China	1	Not reported	Male patients ages 20 to 60 years with a DSM-IV diagnosis of schizophrenia and >5-year duration of illness.	20 Hz stimulus on left DLPFC 40 sessions, administered 5 times a week (Monday to Friday) for 8 weeks (n=28)	8 weeks
Kumar et al (2020) <sup>44,</sup>	India	1	Not reported	<p>Patients who were right-handed, clinically diagnosed as having schizophrenia as per ICD-10 criteria for at least 1 year; on stable doses of medicines (if receiving) for the last 4 weeks, but continued to have significant negative symptoms.</p> <p>Excluded patients who had received rTMS treatment in the past for a similar condition, comorbid ICD-10 Axis I diagnosis, or Axis II Personality Disorder or any other exclusion</p>	<p>Active rTMS: 20 sessions of high frequency rTMS per day (5 consecutive sessions per week for 4 weeks) at 20 Hz frequency (n=50)</p> <p>Sham rTMS (n=50)</p>	4 months

Study; Trial	Countries	Sites	Dates	Participants	Interventions <sup>1</sup>		Duration of follow- up
				criteria common to every TMS protocol.			
Zhuo et al (2019) <sup>45</sup> ,	China	1	2013-2014	<p>Adults ages 20 to 60 years with a DSM-IV diagnosis of schizophrenia; on a stable dose of antipsychotic medication for at least 1 month before study enrollment.</p> <p>Exclusions: DSM-IV-TR axis I disorder other than schizophrenia; history of epilepsy or seizure; significant or unstable neurologic disorder; cardiac pacemaker; previous brain injury or surgery; any metal clips, plates, or other metal items in the head; or substance dependency; or ECT within 3 months.</p>	Active rTMS: 20 treatment sessions on consecutive weekdays. 20 Hz rTMS applied to the left DLPFC (n=35)	Sham rTMS (n=35)	4 weeks

DLPFC: dorsolateral prefrontal cortex; DSM-IV: Diagnostic and Statistical Manual of Mental Disorders, 4th edition; ECT: electroconvulsive therapy; ICD-10: International Classification of Disease, 10th edition; rTMS: repetitive transcranial magnetic stimulation.

**Table 13. Summary of Key Randomized Controlled Trial Results**

Study	Main Results
Ye et al (2024) <sup>40</sup> ,	During the 6-week treatment, 20 Hz rTMS, but not 10 Hz, produced greater gains in immediate memory compared with the sham group ( $p < .05$ ). By 24 weeks, both 10 Hz and 20 Hz rTMS showed benefits on immediate memory and RBANS total score ( $p < .05$ ).
Hua et al (2024) <sup>41</sup> ,	At 2 weeks, rTMS significantly reduced AHRS scores compared with sham (MD, 5.96; $p < .001$ ), and this effect was sustained through week 6 (MD, 7.89; $p < .001$ ). Individuals in the rTMS group also reported significantly better response rates, PANSS scores, and HAMD scores at both the 2-week and 6-week assessments ( $p < .05$ ).
Xiu et al (2024) <sup>42</sup> ,	Significant differences were observed in favor of the 20Hz and 10Hz groups for immediate memory, delayed memory, language, and RBANS total score compared to the sham control at 6-month follow-up ( $p < .05$ ).
Zhu et al (2021) <sup>46</sup> ,	At 2, 6, 12, and 24 weeks after the end of treatment, PANSS negative symptom scores were significantly lower in the rTMS group compared to the sham group ( $p < .05$ ). The effect of treatment on positive symptoms and PANSS total scores was not significant.
Guan et al (2020) <sup>43</sup> ,	At 2 weeks, 4 weeks, and 6 weeks, no significant differences in PANSS total score and sub scores between the sham and treatment groups. Immediate memory performance was higher in the rTMS group compared with the sham group at week 8 after covarying for education, age, and dose of drug. The improvement in immediate memory score was correlated with a decrease in the excitement factor score.
Kumar et al (2020) <sup>44</sup> ,	Total SANS score was reduced significantly after the intervention in both the active ( $60.6 \pm 11.75$ to $43.9 \pm 12.67$ ; $p < .01$ ) and sham ( $61.5 \pm 13.69$ to $50.5 \pm 14.11$ ; $p < .01$ ) groups. Post-intervention scores were significantly lower among the subjects who received active rTMS as compared to those who received sham.
Zhuo et al (2019) <sup>45</sup> ,	Significant decrease in negative symptoms but no significant improvement in cognition.

AHRS: Auditory Hallucination Rating Scale; HAMD: Hamilton Depression Rating Scale; MD: mean difference; PANSS: Positive and Negative Syndrome Scale; rTMS: repetitive transcranial magnetic stimulation; RBANS: Repeatable Battery for the Assessment of Neuropsychological Status; SANS: Scale for Assessing Negative Symptoms in Schizophrenia.

**Table 14. Study Relevance Limitations**

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Follow-Up <sup>e</sup>
Ye et al (2024) <sup>40</sup> ,	4. Included inpatient men only				
Hua et al (2024) <sup>41</sup> ,					1. 6 weeks not sufficient to show durability of effects

<b>Study</b>	<b>Population<sup>a</sup></b>	<b>Intervention<sup>b</sup></b>	<b>Comparator<sup>c</sup></b>	<b>Outcomes<sup>d</sup></b>	<b>Follow-Up<sup>e</sup></b>
Xiu et al (2024) <sup>42,</sup>					
Zhu et al (2021) <sup>46,</sup>	4. Included inpatients only				
Guan et al (2020) <sup>43,</sup>	4. Included men only				1. 8 weeks not sufficient to show durability of effects
Kumar et al (2020) <sup>44,</sup>					
Zhuo et al (2019) <sup>45,</sup>					1. 4 weeks not sufficient to show durability of effects

The evidence limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4. Enrolled populations do not reflect relevant diversity; 5. Other.

<sup>b</sup> Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4. Not the intervention of interest (e.g., proposed as an adjunct but not tested as such); 5. Other.

<sup>c</sup> Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively; 5. Other.

<sup>d</sup> Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Incomplete reporting of harms; 4. Not establish and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported; 7. Other.

<sup>e</sup> Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms; 3. Other.

**Table 15. Study Design and Conduct Limitations**

<b>Study</b>	<b>Allocation<sup>a</sup></b>	<b>Blinding<sup>b</sup></b>	<b>Selective Reporting<sup>c</sup></b>	<b>Data Completeness<sup>d</sup></b>	<b>Power<sup>e</sup></b>	<b>Statistical<sup>f</sup></b>
Ye et al (2024) <sup>40,</sup>				1. 25% of patients were lost to follow-up by 24 weeks	1. power calculation not reported	
Hua et al (2024) <sup>41,</sup>				1. 4 discontinued in active vs 12 in sham by follow-up (26%)		



Study	Allocation <sup>a</sup>	Blinding <sup>b</sup>	Selective Reporting <sup>c</sup>	Data Completeness <sup>d</sup>	Power <sup>e</sup>	Statistical <sup>f</sup>
				dropout differential)		
Xiu et al (2024) <sup>42,</sup>				1. 26% of patients were lost to follow-up by 32 weeks.	1. power calculation not reported	
Zhu et al (2021) <sup>46,</sup>					1. power calculation not reported	
Guan et al (2020) <sup>43,</sup>				1. 15/56 (26.8%) patients discontinued	1. power calculation not reported	
Kumar et al (2020) <sup>44,</sup>				1. 33% attrition (32% active and 38% sham)		
Zhuo et al (2019) <sup>45,</sup>				1. 10/70 discontinued (14.3%)	1. power calculation not reported	

The evidence limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

<sup>b</sup> Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

<sup>c</sup> Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>d</sup> Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

<sup>e</sup> Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

<sup>f</sup> Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

## SUBSTANCE USE DISORDER AND CRAVING

### Systematic Review

Jansen et al (2013) reported on results from a meta-analysis evaluating the effect of rTMS and transcranial direct current stimulation of the DLPFC on substance dependence (alcohol, nicotine, cocaine, marijuana) or food craving.<sup>47</sup> Seventeen double-blind, sham-controlled trials that used high-frequency stimulation were analyzed. Thirteen studies stimulated the left DLPFC and 7 studies stimulated the right DLPFC or both sides. Twelve of the studies gave only 1 or 2 sessions. The standardized effect size was 0.476 (95% CI, 0.316 to 0.636), indicating a medium effect size

for active stimulation over sham for a reduction in craving. However, the studies were small (range, 9 to 48 patients) and there was significant heterogeneity in selected studies. No significant differences were found in the effectiveness of rTMS versus transcranial direct current stimulation, the different substances, or the side of stimulation, although this analysis might have been biased by the number of studies for each condition.

Chang et al (2022) conducted a meta-analysis of 7 double-blind RCTs (N=462) that used rTMS to treat methamphetamine use disorder.<sup>48</sup> All studies targeted the left DLPFC and the number of sessions ranged among studies from 5 to 20. Mean craving scores at baseline ranged from 22.63 to 57.68. A random effects model showed that clinical craving scores were significantly lower with rTMS than sham treatment (SMD, 0.983; 95% CI, 0.620 to 1.345;  $p \leq .001$ ;  $I^2=67.814\%$ ). According to a subgroup analysis, intermittent theta burst stimulation had a greater effect than 10-Hz rTMS. The authors concluded that further trials with larger sample sizes are needed.

## **NEUROLOGIC DISORDERS OTHER THAN MIGRAINE**

### **AMYOTROPHIC LATERAL SCLEROSIS OR MOTOR NEURON DISEASE**

#### **Systematic Review**

A Cochrane review by Fang et al (2013) identified 3 RCTs with a total of 50 participants with amyotrophic lateral sclerosis that compared rTMS with sham TMS.<sup>49</sup> All trials were considered of poor methodologic quality. Heterogeneity prevented pooling of results, and the high rate of attrition further increased the risk of bias. Reviewers concluded that evidence was insufficient to draw conclusions about the efficacy and safety of rTMS in the treatment of amyotrophic lateral sclerosis.

### **CHRONIC PAIN**

#### **Systematic Reviews**

A Cochrane review by O'Connell et al (2018) evaluating noninvasive brain stimulation techniques was first published in 2010 and was updated in 2014<sup>50</sup>, and 2018.<sup>51</sup> The reviewers identified 42 RCTs (range, 4 to 70 participants) on TMS for chronic pain. Meta-analysis of 27 rTMS studies versus sham (N=655 participants) for pain intensity at short-term follow-up (0 to <1 week postintervention) demonstrated a small effect with heterogeneity (SMD, -0.22; 95% CI, -0.29 to -0.16, low-quality evidence). This equates to a 7% (95% CI, 5% to 9%) reduction in pain, or a 0.40 (95% CI, 0.53 to 0.32) point reduction on a 0 to 10 pain intensity scale, which did not meet the minimum clinically important difference threshold of 15% or greater. There is very low-quality evidence that single doses of high-frequency rTMS of the motor cortex may have short-term effects on chronic pain and quality of life, but multiple sources of bias exist that may have influenced the observed effects. We did not find evidence that low-frequency rTMS, rTMS applied to the DLPFC, and cranial electrotherapy stimulation are effective for reducing pain intensity in chronic pain.

Jiang et al (2022) conducted a systematic review and meta-analysis of 38 RCTs that assessed the analgesic effect of rTMS in 1338 patients with neuropathic pain.<sup>52</sup> A single rTMS session was used in 13 studies and multiple sessions were used in the remaining 25 studies. The overall risk of bias in most studies was low or uncertain. According to a random effects analysis, rTMS was

superior to sham therapy in reducing pain scores (effect size, -0.66; 95% CI, -0.87 to -0.46;  $p < .001$ ;  $I^2 = 78\%$ ). Beneficial effects of rTMS on pain were observed at 1 month ( $p < .001$ ) and 2 months ( $p = .01$ ). Low-frequency rTMS ( $\leq 1$  Hz) did not effectively reduce pain compared to higher frequency stimulation. The analysis did not find an increased risk of adverse events with rTMS compared to sham therapy. The authors concluded that larger, well-designed trials are needed to determine the long-term effect of rTMS in this setting.

## **EPILEPSY**

### **Systematic Reviews**

A Cochrane review by Chen et al (2016) included 7 RCTs on low-frequency rTMS for epilepsy, 5 of which were completed studies with published data.<sup>53</sup> The total number of participants was 230. All studies had active or placebo controls and 4 were double-blind. However, a meta-analysis could not be conducted due to heterogeneity in designs, interventions, and outcomes of the trials. Therefore, a qualitative synthesis was performed. For the outcome of seizure rate, 2 studies showed a significant reduction and 5 studies did not. Of the 4 studies evaluating the mean number of epileptic discharges, 3 studies showed a statistically significant reduction in discharges. Adverse events were uncommon and mild, involving headaches, dizziness, and tinnitus. There were no significant changes in medication use.

A more recent meta-analysis conducted by Mishra and colleagues (2020) included 7 RCTs that compared rTMS with sham or placebo controls in patients with epilepsy.<sup>54</sup> Two of the included studies showed statistically significant reductions in the seizure rate from baseline, 3 trials failed to show any statistically significant difference in seizure frequency, and 2 had unclear results due to inadequate power. In a meta-regression, when adjusted for other potential variables such as the type of coil used, stimulation frequency, and the total duration of the active intervention, seizure frequency worsened by  $2.00 \pm 0.98$  ( $p = .042$ ) for each week of lengthening of the posttreatment follow-up period. These results suggested that rTMS exerted only a short-term effect. The reviewers concluded that although the procedure may be a therapeutic alternative for patients with drug-resistant epilepsy, further RCTs using standardized protocols and with adequate sample sizes and duration are still needed.

## **FIBROMYALGIA**

### **Systematic Reviews**

Ahmed et al (2025) included 43 RCTs ( $N = 2120$ ) in a network meta-analysis evaluating noninvasive brain stimulation for fibromyalgia.<sup>55</sup> Low-frequency rTMS reduced pain intensity more than sham (SMD, -1.20; 95% CI -1.82 to -0.58), with high-frequency rTMS showing a smaller effect (SMD, -0.58; 95% CI -1.00 to -0.17). Low-frequency rTMS over the right dorsolateral prefrontal cortex (DLPFC) produced the most pronounced effects (SMD, -1.42; 95% CI -2.69 to -0.15) and maintained short-term improvements. High-frequency rTMS over the left primary motor cortex (M1) also reduced pain (SMD, -0.78; 95% CI -1.39 to -0.18). rTMS was well tolerated and achieved clinically meaningful pain reductions.

Su et al (2021) conducted a meta-analysis of 18 RCTs ( $N = 643$ ) with rTMS in patients with fibromyalgia.<sup>56</sup> Reduction in disease influence according to the Fibromyalgia Impact Questionnaire showed a significant effect of rTMS (SMD, -0.7; 95% CI, -1.173 to -0.228). The

effect of rTMS on disease influence, pain, depression, and anxiety lasted for at least 2 weeks after the last session. Older patients were most likely to experience reduced Fibromyalgia Impact Questionnaire scores. The authors concluded that larger RCTs are needed to confirm these findings.

Saltychev and Laimi (2017) published a meta-analysis of rTMS for the treatment of patients with fibromyalgia.<sup>57</sup> The meta-analysis included 7 sham-controlled, double-blind trials with a low risk of bias. Trial sample sizes ranged from 18 to 54 patients. Five studies provided high-frequency stimulation to the left primary motor cortex, and the others were to the right or left DLPFC. The number of sessions ranged from 10 to 24, and follow-up ranged from immediately after treatment to 3 months posttreatment. In the pooled analysis, pain severity decreased after the last simulation by 1.2 points (95% CI, -1.7 to -0.8 points) on a 10-point numeric rating scale, while pain severity measured at 1 week to 1 month after the last simulation decreased by 0.7 points (95% CI, -1.0 to -0.3 points). Both were statistically significant, but not considered clinically significant, based on a minimal clinically important difference of 1.5 points.

## **PARKINSON DISEASE**

### **Systematic Reviews**

A meta-analysis by Chou et al (2015) included 20 sham-controlled randomized trials (N=470 patients) evaluating Parkinson disease.<sup>58</sup> Sample sizes ranged from 8 to 102 patients. The total effect size of low- and high-frequency rTMS on Unified Parkinson's Disease Rating Scale part III score was 0.46, which is considered a small-to-medium effect size, and the mean change in the Unified Parkinson's Disease Rating Scale part III score (-6.42) was considered a clinically important difference. The greatest effect on motor symptoms was from high-frequency rTMS over the primary motor cortex (SMD, 0.77;  $p < .001$ ) and low-frequency rTMS over other frontal regions (SMD, 0.50;  $p = .008$ ). High-frequency rTMS at other frontal regions and low-frequency rTMS over the primary motor cortex did not have a statistically significant benefit. The largest trial included in the systematic review was an exploratory, multicenter, double-blind trial reported by Shirota et al (2013) who randomized 106 patients to 8 weeks of 1-Hz rTMS, 10-Hz rTMS, or sham stimulation over the supplementary motor area.<sup>59</sup> At 9 weeks, all groups showed a similar amount of improvement.

Li et al (2022) conducted a meta-analysis of 32 sham-controlled RCTs of rTMS in patients with Parkinson disease and motor dysfunction (N=1048 patients).<sup>60</sup> Motor dysfunction was assessed using the United Parkinson's Disease Rating Scale part III score. Overall, rTMS had a significant effect on motor symptoms compared to sham (SMD, 0.64; 95% CI, 0.47 to 0.80;  $p < .0001$ ;  $I^2 = 64\%$ ). High-frequency rTMS to the primary motor cortex was the most effective intervention. Significant benefit of rTMS was also demonstrated for akinesia, rigidity, and tremor.

## **STROKE RECOVERY**

### **Systematic Reviews**

A number of RCTs and systematic reviews have evaluated rTMS for recovery from stroke. For example, a Cochrane review by Hao et al (2013) included 19 RCTs (N=588 participants) evaluating the effect of low- and high-frequency TMS for improving function after stroke.<sup>61</sup> The 2 largest trials (n=183 patients) showed that rTMS was not associated with a significant

improvement in Barthel Index scores. Four trials (n=73) found no significant effect on motor function. Subgroup analyses for different stimulation frequencies or durations of illness also did not show a significant benefit of rTMS compared with sham rTMS or no treatment. Reviewers concluded that current evidence did not support the routine use of rTMS for the treatment of stroke.

A meta-analysis by Le et al (2014) assessed the effect of rTMS on the recovery of hand function and excitability of the motor cortex after stroke.<sup>62</sup> Eight RCTs (N=273 participants) were selected. The quality of the trials was rated moderate to high, although the size of the studies was small. There was variability in the time since stroke (5 days to 10 years), in the frequency of rTMS applied (1 to 25 Hz for 1 second to 25 min/d), and the stimulation sites (primary motor cortex or premotor cortex of the unaffected hemisphere). Meta-analysis found a positive effect on finger motor ability (4 studies; n=79 patients; SMD, 0.58) and hand function (3 studies; n=74 patients; SMD, -0.82), but no significant change in motor evoked potentials (n=43) or motor threshold (n=62).

A meta-analysis by Li et al (2015) included 4 RCTs on low-frequency rTMS over the right parastriangularis for patients (N=137) with aphasia after stroke.<sup>63</sup> All studies used double-blinding, but therapists were not blinded. Every trial used a different outcome measure, and sample sizes were small (range, 12 to 40 patients). Meta-analysis showed a medium effect size for naming (p=.004), a trend for a benefit on repetition (p=.08), and no significant benefit for comprehension (p=.18). Additional study in a larger number of patients would be needed to determine with greater certainty the effect of this treatment on aphasia after stroke.

Qiao et al (2022) performed a meta-analysis of RCTs that assessed the effect of rTMS in 433 patients with post-stroke dysphagia.<sup>64</sup> Twelve trials that used dysphagia severity rating scales (Dysphagia Grade and Penetration Aspiration Scale) were included. The specific controls used in each study were not specified. Study characteristics included duration of treatment of 1 to 10 days, stimulation frequency of 1 to 10 Hz, and duration of stimulation of 5 to 20 minutes. The analysis favored rTMS (SMD, -0.67; 95% CI, -0.88 to -0.45; p<.001; I<sup>2</sup>=42%). Subgroup analyses identified treatment duration >5 days and rTMS during the subacute phase after stroke as potential situations with greater clinical benefit, but there was no difference in efficacy according to stimulation frequency, location, or duration of each stimulation. The authors noted that publication bias was present and there may be limited clinical applicability of the dysphagia rating scales.

Zhang et al (2017) published a systematic review and meta-analysis evaluating the effects of rTMS on upper-limb motor function after stroke.<sup>65</sup> A search through October 2016 yielded 34 RCTs with a total of 904 participants (range, 6 to 108 participants). Pooled estimates found improvement with rTMS for both short-term (SMD, 0.43; p<.001) and long-term (SMD, 0.49; p<.001) manual dexterity. Of the 28 studies reporting on adverse events, 25 studies noted none. Mild adverse events, such as headache and increased anxiety, were reported in 3 studies. The review was limited by variation in TMS protocols across studies.

Graef et al (2016) reported a systematic review of rTMS combined with upper-limb training for improving function after stroke.<sup>66</sup> Included were 11 sham-controlled randomized trials with 199 patients that evaluated upper-limb motor and functional status and spasticity; 8 RCTs with

sufficient data were included in the meta-analysis. These studies were considered to have a low-to-moderate risk of bias. In the overall analysis, there was no benefit of rTMS on upper-limb function or spasticity (SMD, 0.03; 95% CI, -0.25 to 0.32).

Zhang et al (2024) conducted a systematic review and meta-analysis of 37 RCTs (N=1887) assessed as having a low risk of bias evaluating rTMS for motor recovery after stroke.<sup>67</sup> Pooled results showed significant improvements in upper-limb motor function following rTMS at last follow-up compared to control on the Fugl-Meyer Assessment (SMD 5.2; 95% CI, 0.5 to 10.0;  $p < .001$ ;  $I^2 = 98\%$ ), with larger effects in acute/subacute stroke and in patients with more severe baseline deficits. Among secondary measures, only the modified Rankin Scale improved (SMD = -0.5; 95% CI, -0.9 to -0.1;  $p = .013$ ;  $I^2 = 83\%$ ); Action Research Arm Test (ARAT), Box and Block Test (BBT), and Modified Ashworth Scale (MAS) showed no between-group differences. Substantial heterogeneity was present for all estimates, and trials varied widely in terms of rTMS target site, intensity, number of sessions, and pulses per day.

### **Section Summary: Psychiatric or Neurologic Disorders Other Than Depression, Migraine, or Obsessive-Compulsive Disorder**

For individuals who have psychiatric disorders other than depression or OCD (eg, panic disorder, generalized anxiety disorder, posttraumatic stress disorder, schizophrenia, substance use disorder and craving) who receive rTMS, the evidence includes numerous small RCTs and meta-analyses of these studies. The trials included in the meta-analyses are typically small and of low methodologic quality. In addition, stimulation parameters have not been established, and trial results are heterogeneous. A number of sham-controlled randomized trials and meta-analyses of these have found a medium effect size of rTMS for the reduction of substance dependence or food craving. Most studies examined acute craving after 1 or 2 rTMS sessions, and there is limited evidence on the longer-term efficacy of this treatment approach. There are no large, high-quality trials for any of these conditions demonstrating efficacy or the durability of any treatment effects.

For individuals who have neurological disorders other than migraine (eg, amyotrophic lateral sclerosis, chronic pain, epilepsy, fibromyalgia, Parkinson disease, and stroke) who receive rTMS, the evidence includes numerous small RCTs and meta-analyses of these randomized trials. The trials included in the meta-analyses are typically small and of low methodologic quality. In addition, stimulation parameters have not been established, and trial results are heterogeneous. There are no large, high-quality trials for any of these conditions demonstrating efficacy or the durability of any treatment effects.

### **SUPPLEMENTAL INFORMATION**

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

### **Clinical Input From Physician Specialty Societies and Academic Medical Centers**

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

## **2014 Input**

In response to requests, input was received from 1 physician specialty society and 3 academic medical centers while this policy was under review in 2014. Reviewers considered repetitive transcranial magnetic stimulation (rTMS) to be medically necessary for treatment-resistant depression. Input agreed with the proposed criteria for treatment of treatment-resistant depression with rTMS, as included in the policy statement.

## **Practice Guidelines and Position Statements**

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

## **American Academy of Child and Adolescent Psychiatry**

In 2013, the American Academy of Child and Adolescent Psychiatry published practice parameters on the assessment and treatment of children and adolescents with tic disorders.<sup>68</sup> The Academy did not recommend rTMS, citing the limited evidence on the safety, ethics, and long-term impact on development.

## **American Psychiatric Association**

The American Psychiatric Association (2018) published consensus recommendations on rTMS for the treatment of depression.<sup>69</sup> The guidelines state, "Multiple randomized controlled trials and published literature have supported the safety and efficacy of rTMS antidepressant therapy." The recommendations include information on the following variables: clinical environment, operator requirements, documentation, coils, cortical targets, coil positioning methods, determination of motor threshold, number of treatment sessions for acute treatment, and allowable psychotropic medications during TMS treatment.

The American Psychiatric Association's (2007, reaffirmed in 2012) guidelines on the treatment of patients with obsessive-compulsive disorder have indicated that "findings of the 4 published trials of rTMS are inconsistent, perhaps because the studies differed in design, stimulation sites, duration, and stimulation parameters. The available results and the technique's non-invasiveness and good tolerability should encourage future research, but the need for daily treatment may limit the use of TMS in practice."

## **Veteran's Affairs/Department of Defense**

The 2022 Veteran's Affairs/Department of Defense (VA/DoD) guideline for management of major depressive disorder recommends offering rTMS to patients who have experienced partial response or no response to an adequate trial of 2 or more pharmacologic treatments (strength of recommendation: weak).<sup>70</sup> Recommended options for the second treatment attempt after the initial therapy tried include switching to another antidepressant or adding augmentation therapy with a second-generation antipsychotic. The recommendation for rTMS was graded as weak due to limitations of the available literature including small study effects, high rates of discontinuation, lack of allocation concealment, and the practical limitations of the need for daily treatment and lack of widespread access to facilities that offer this therapy. The guideline also

concluded that there is limited evidence to recommend for or against theta-burst stimulation for treatment of depression.

The 2023 VA/DoD guideline for management of bipolar disorder states "for individuals with bipolar disorder who have demonstrated partial or no response to pharmacologic treatment for depressive symptoms, we suggest offering repetitive transcranial magnetic stimulation [rTMS] as an adjunctive treatment."<sup>71</sup> However, the recommendation was rated as "weak" and the confidence in the evidence was very low. For the management of PTSD, the 2023 guideline found insufficient evidence for or against rTMS.<sup>72</sup>

### **National Institute for Health and Care Excellence**

In 2015, the National Institute for Health and Care Excellence (NICE) provided revised guidance, stating that evidence on the short-term efficacy of rTMS for depression is adequate, although the clinical response is variable and some patients may not benefit.<sup>73</sup>

In 2014, the NICE provided guidance on the use of rTMS for treating and preventing migraine.<sup>74</sup> The guidance stated that evidence on the efficacy of TMS for the treatment of migraine was limited in quantity and for the prevention of migraine was limited in both quality and quantity. Evidence on its safety in the short- and medium-term was adequate, but there was uncertainty about the safety of long-term or frequent use of TMS.

In 2020, the NICE stated that rTMS has not demonstrated any major safety concerns for management of obsessive-compulsive disorder or auditory hallucinations, but evidence for both uses is lacking; therefore, NICE recommends that rTMS be used in patients with these conditions only in the context of research.<sup>75,76</sup>

### **International Neuromodulation Society/North American Neuromodulation Society**

In 2020, an expert consensus panel from the International Neuromodulation Society-North American Neuromodulation Society performed a literature review and published recommendations for transcranial magnetic stimulation in the treatment of pain and headache.<sup>77</sup> For neuropathic pain, the panel recommended transcranial magnetic stimulation to the primary motor cortex (high level evidence) or the left dorsolateral prefrontal cortex (F3 location) (at least moderate level evidence). For postoperative pain, the panel recommended that transcranial magnetic stimulation to the F3 location be only selectively offered due to at least moderate certainty that the net benefit is small. For primary headache, the panel only based 2 recommendations on moderate certainty evidence: single transcranial magnetic stimulation for acute migraine and high-frequency rTMS to the primary motor cortex for migraine prevention. For posttraumatic brain injury, high level evidence supported a recommendation for high-frequency transcranial magnetic stimulation to the primary motor cortex or the F3 location.

### **U.S. Preventive Services Task Force Recommendations**

Not applicable.

### **Ongoing and Unpublished Clinical Trials**

Some currently ongoing trials that might influence this review are listed in Table 16.



**Table 16. Summary of Key Trials**

<b>NCT No.</b>	<b>Trial Name</b>	<b>Planned Enrollment</b>	<b>Completion Date</b>
<b><i>Unpublished</i></b>			
NCT02910024	Theta-Burst-Stimulation in Early Rehabilitation of Stroke (TheSiReS)	150	Sep 2022
NCT03556722	Effectiveness and Tolerability of Repetitive Transcranial Magnetic Stimulation For Preventive Treatment Of Episodic Migraine: A Single Centre, Randomised, Double-Blind, Sham-Controlled Phase 2 Trial	76	Apr 2022
<b><i>Ongoing</i></b>			
NCT06545474	Repetitive Transcranial Magnetic Stimulation (rTMS) as an Early Intervention in Major Depression (MD) Compared to Antidepressant Selective Serotonin Reuptake Inhibitor (SSRI) Medication (Early-TMS)	106	Dec 2026
NCT06370988	Theta-Burst Stimulation for Bipolar Depression	124	May 2029
NCT06524505	Efficacy, Tolerability, and Cognitive Effects of Deep Transcranial Magnetic Stimulation for Bipolar Depression: a Double-blind, Randomized Controlled Trial	100	Oct 2024
NCT05389670	Theta-burst Repetitive Transcranial Magnetic Stimulation (TBS) of the Right Inferior Frontal Gyrus for Treatment of Nicotine Dependence	60	Apr 2026
NCT05331937	Transcranial Magnetic Stimulation (TMS) for Patients With Exposure Therapy-resistant Obsessive-compulsive Disorder (OCD): TETRO - a Multicenter Randomized Controlled Trial	250	Sep 2027
NCT05100888	Theta-burst rTMS in Schizophrenia to Ameliorate Negative and Cognitive Symptoms: a Double-blind, Randomized Clinical Trial	90	Dec 2025

NCT: national clinical trial.

## CODING

**The following codes for treatment and procedures applicable to this policy are included below for informational purposes. This may not be a comprehensive list of procedure codes applicable to this policy.**

**Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.**

**The code(s) listed below are medically necessary ONLY if the procedure is performed according to the "Policy" section of this document.**

<b>CPT/HCPCS</b>	
90867	Therapeutic repetitive transcranial magnetic stimulation (TMS) treatment; initial, including cortical mapping, motor threshold determination, delivery and management
90868	Therapeutic repetitive transcranial magnetic stimulation (TMS) treatment; subsequent deliver and management, per session
90869	Therapeutic repetitive transcranial magnetic stimulation (TMS) treatment; subsequent motor threshold re-determination with delivery and management
0889T	Personalized target development for accelerated, repetitive high-dose functional connectivity MRI-guided theta-burst stimulation derived from a structural and resting-state functional MRI, including data preparation and transmission, generation of the target, motor threshold-starting location, neuronavigation files and target report, review and interpretation
0890T	Accelerated, repetitive high-dose functional connectivity MRI-guided theta-burst stimulation, including target assessment, initial motor threshold determination, neuronavigation, delivery and management, initial treatment
0891T	Accelerated, repetitive high-dose functional connectivity MRI-guided theta-burst stimulation, including neuronavigation, delivery and management, subsequent treatment day
0892T	Accelerated, repetitive high-dose functional connectivity MRI-guided theta-burst stimulation, including neuronavigation, delivery and management, subsequent motor threshold redetermination with delivery and management, per treatment day

<b>REVISIONS</b>	
12-07-2012	Policy added to the bcbsks.com web site.
05-07-2013	Updated Description section.
	Updated Rationale section.
	Updated Reference section.
10-15-2014	Updated Description section.
	In Policy section: Removed the following, "Transcranial magnetic stimulation of the brain is considered investigational as a treatment of all other psychiatric/neurologic disorders, including but not limited to bipolar disorder, schizophrenia, obsessive-compulsive disorder, or migraine headaches."
	And added:

<b>REVISIONS</b>	
	<p>A. Repetitive transcranial magnetic stimulation (rTMS) of the brain may be considered medically necessary as a treatment of major depressive disorder when all of the following conditions (1-3) have been met:</p> <ol style="list-style-type: none"> <li>1. Confirmed diagnosis of severe major depressive disorder (single or recurrent) documented by standardized rating scales that reliably measure depressive symptoms; AND</li> <li>2. Any one of the following (a, b, c, or d): <ol style="list-style-type: none"> <li>a. Failure of 4 trials of psychopharmacologic agents including 2 different agent classes and 2 augmentation trials; OR</li> <li>b. Inability to tolerate a therapeutic dose of medications as evidenced by 4 trials of psychopharmacologic agents with distinct side effects; OR</li> <li>c. History of response to rTMS in a previous depressive episode (at least 3 months since the prior episode); OR</li> <li>d. Is a candidate for electroconvulsive therapy (ECT) and ECT would not be clinically superior to rTMS (e.g., in cases with psychosis, acute suicidal risk, catatonia or life-threatening inanition rTMS should NOT be utilized); AND</li> </ol> </li> <li>3. Failure of a trial of a psychotherapy known to be effective in the treatment of major depressive disorder of an adequate frequency and duration, without significant improvement in depressive symptoms, as documented by standardized rating scales that reliably measure depressive symptoms.</li> </ol> <p>B. rTMS for major depressive disorder that does not meet the criteria listed above is considered experimental / investigational.</p> <p>C. Continued treatment with rTMS of the brain as maintenance therapy is considered experimental / investigational.</p> <p>Added Policy Guideline section.</p> <p>In Coding section:</p> <ul style="list-style-type: none"> <li>• Added ICD-9 diagnoses: 296.23, 296.33</li> <li>• Added ICD-10 diagnoses: F32.2, F33.2</li> </ul> <p>Updated Rationale section.</p> <p>Updated References section.</p>
03-18-2015	<p>Updated Description section.</p> <p>In Policy section:</p> <ul style="list-style-type: none"> <li>▪ Added Item D: " Transcranial magnetic stimulation of the brain is considered experimental / investigational as a treatment of all other psychiatric/neurologic disorders, including but not limited to bipolar disorder, schizophrenia, obsessive-compulsive disorder, or migraine headaches."</li> </ul> <p>Updated Rationale section.</p> <p>Updated References section.</p>
02-17-2016	<p>Updated Description section.</p> <p>Updated Rationale section.</p> <p>Updated References section.</p>
08-15-2017	<p>Updated Description section.</p> <p>Updated Rationale section.</p> <p>Updated References section.</p>
11-07-2018	<p>Updated Description section.</p> <p>Updated Rationale section.</p> <p>In Coding section:</p> <ul style="list-style-type: none"> <li>▪ Removed ICD-9 codes.</li> </ul>

<b>REVISIONS</b>	
	Updated References section.
05-22-2020	Updated Description section
	In Policy section: Update Policy Guidelines <ul style="list-style-type: none"> <li>Added Items 2 and 3 to read "2. The physician is responsible for the initial mapping (once per course or delivery) and the management of the treatment. The physician is also responsible for the re-determinations. 3. A trained technician may perform the subsequent delivery of treatment." In Item 5 a removed "An attendant trained" and "as well as the use of the equipment" to read "A trained technician in administration of the treatments, and in basic cardiac life support, the management of complications such as seizures, must be present at all times;"</li> </ul>
	Updated Rationale section
	Updated References
04-16-2021	Updated Description section.
	Updated Rationale section.
	Updated References section.
12-18-2021	Updated Title to: Transcranial Magnetic Stimulation (TMS) as a Treatment of Depression and Other Psychiatric/Neurologic Disorders
	Updated Description Section
	Updated Policy Section: <ul style="list-style-type: none"> <li>Section A to read: Repetitive transcranial magnetic stimulation (rTMS) of the brain using an FDA-cleared device and modality, which can include but is not limited to, conventional TMS, deep TMS, and theta burst stimulation (see Policy Guidelines) may be considered medically necessary as a treatment of major depressive disorder when all of the following conditions (1-3) have been met:</li> </ul>
	Updated Policy Guidelines <ul style="list-style-type: none"> <li>Added lines 4, 5, 7, 9</li> </ul>
	Updated Rationale Section
	Updated Coding section: <ul style="list-style-type: none"> <li>Added ICD-10 Codes: F32.2, F32.3, F32.4, F32.5, F33.3, F33.8</li> </ul>
	Updated References Section
12-29-2022	Updated Description Section
	Updated Policy Section <ul style="list-style-type: none"> <li>Section A2a Changed to read: "Individual has tried and had an inadequate response to 2 antidepressant agents from 2 different antidepressant classes (i.e., selective serotonin reuptake inhibitors, serotonin and norepinephrine reuptake inhibitors, tricyclic antidepressants, bupropion, or mirtazapine). An adequate trial of an antidepressant is defined by BOTH of the following: <ol style="list-style-type: none"> <li>The trial length was at least 6 weeks at generally accepted doses; AND</li> <li>Individual was ≥80% adherent to the agent during the trial."</li> </ol> </li> <li>Section A2b Removed: "as evidenced by 4 trials of psychopharmacologic agents with" and added "due to"</li> </ul>
	Updated Rationale Section

<b>REVISIONS</b>	
	Updated References Section
11-17-2023	Updated Description Section
	Updated Rationale Section
	Updated Coding Section <ul style="list-style-type: none"> <li>Removed ICD-10 Codes</li> </ul>
	Updated References Section
07-01-2024	Updated Coding Section <ul style="list-style-type: none"> <li>Added codes 0889T, 0890T, 0891T and 0892T (eff. 07-01-2024)</li> </ul>
12-03-2024	Updated Description Section
	Updated Rationale Section
	Updated References Section
02-25-2025	Updated Policy Section <ul style="list-style-type: none"> <li>Removed A3: "Failure of a trial of a psychotherapy known to be effective in the treatment of major depressive disorder of an adequate frequency and duration, without significant improvement in depressive symptoms, as documented by standardized rating scales that reliably measure depressive symptoms."</li> </ul>
12-09-2025	Updated Description Section
	Updated Rationale Section
	Updated Reference Section

## REFERENCES

1. FDA Briefing Document Psychopharmacologic Drugs Advisory Committee (PDAC) and Drug Safety and Risk Management (DSaRM) Advisory Committee Meeting February 12, 2019. <https://www.fda.gov/media/121376/download>. Accessed September 26, 2025.
2. Zimmerman M, Chelminski I, Posternak M. A review of studies of the Montgomery-Asberg Depression Rating Scale in controls: implications for the definition of remission in treatment studies of depression. *Int Clin Psychopharmacol*. Jan 2004; 19(1): 1-7. PMID 15101563
3. Center for Drug Evaluation and Research Application Number: 211243Orig1s000 Summary Review  
[https://www.accessdata.fda.gov/drugsatfda\\_docs/nda/2019/211243Orig1s000SumR.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/nda/2019/211243Orig1s000SumR.pdf). Accessed August 18, 2025.
4. Alphs L, Fu D-J, Williamson D, et al. Validation and mapping of the Suicidal Ideation and Behavior Assessment Tool (SIBAT). (abstract W88) *Neuropsychopharmacology*. 2018;43:S427S428.
5. Gross M, Nakamura L, Pascual-Leone A, et al. Has repetitive transcranial magnetic stimulation (rTMS) treatment for depression improved? A systematic review and meta-analysis comparing the recent vs. the earlier rTMS studies. *Acta Psychiatr Scand*. Sep 2007; 116(3): 165-73. PMID 17655557
6. Schutter DJ. Antidepressant efficacy of high-frequency transcranial magnetic stimulation over the left dorsolateral prefrontal cortex in double-blind sham-controlled designs: a meta-analysis. *Psychol Med*. Jan 2009; 39(1): 65-75. PMID 18447962
7. Gallop L, Westwood SJ, Hemmings A, et al. Effects of repetitive transcranial magnetic stimulation in children and young people with psychiatric disorders: a systematic review. *Eur Child Adolesc Psychiatry*. Feb 2025; 34(2): 403-422. PMID 38809301

8. Sehatzadeh Sh, Tu HA, Palimaka S, et al. Repetitive Transcranial Magnetic Stimulation for Treatment-Resistant Depression: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Ont Health Technol Assess Ser.* 2016; 16(5): 1-66. PMID 27099642
9. Brunoni AR, Chaimani A, Moffa AH, et al. Repetitive Transcranial Magnetic Stimulation for the Acute Treatment of Major Depressive Episodes: A Systematic Review With Network Meta-analysis. *JAMA Psychiatry.* Feb 01 2017; 74(2): 143-152. PMID 28030740
10. Voigt JD, Leuchter AF, Carpenter LL. Theta burst stimulation for the acute treatment of major depressive disorder: A systematic review and meta-analysis. *Transl Psychiatry.* May 28 2021; 11(1): 330. PMID 34050123
11. Papakostas GI, Trivedi MH, Shelton RC, et al. Comparative effectiveness research trial for antidepressant incomplete and non-responders with treatment resistant depression (ASCERTAIN-TRD) a randomized clinical trial. *Mol Psychiatry.* Aug 2024; 29(8): 2287-2295. PMID 38454079
12. Blumberger DM, Vila-Rodriguez F, Thorpe KE, et al. Effectiveness of theta burst versus high-frequency repetitive transcranial magnetic stimulation in patients with depression (THREE-D): a randomised non-inferiority trial. *Lancet.* Apr 28 2018; 391(10131): 1683-1692. PMID 29726344
13. Food and Drug Administration. 510(k) Summary: Brainsway deep TMS System (K122288). 2013; [https://www.accessdata.fda.gov/cdrh\\_docs/pdf12/k122288.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf12/k122288.pdf). Accessed August 20, 2025.
14. Kedzior KK, Reitz SK, Azorina V, et al. Durability of the antidepressant effect of the high-frequency repetitive transcranial magnetic stimulation (rTMS) In the absence of maintenance treatment in major depression: a systematic review and meta-analysis of 16 double-blind, randomized, sham-controlled trials. *Depress Anxiety.* Mar 2015; 32(3): 193-203. PMID 25683231
15. Dunner DL, Aaronson ST, Sackeim HA, et al. A multisite, naturalistic, observational study of transcranial magnetic stimulation for patients with pharmacoresistant major depressive disorder: durability of benefit over a 1-year follow-up period. *J Clin Psychiatry.* Dec 2014; 75(12): 1394-401. PMID 25271871
16. Richieri R, Guedj E, Michel P, et al. Maintenance transcranial magnetic stimulation reduces depression relapse: a propensity-adjusted analysis. *J Affect Disord.* Oct 2013; 151(1): 129-35. PMID 23790811
17. Connolly KR, Helmer A, Cristancho MA, et al. Effectiveness of transcranial magnetic stimulation in clinical practice post-FDA approval in the United States: results observed with the first 100 consecutive cases of depression at an academic medical center. *J Clin Psychiatry.* Apr 2012; 73(4): e567-73. PMID 22579164
18. Janicak PG, Nahas Z, Lisanby SH, et al. Durability of clinical benefit with transcranial magnetic stimulation (TMS) in the treatment of pharmacoresistant major depression: assessment of relapse during a 6-month, multisite, open-label study. *Brain Stimul.* Oct 2010; 3(4): 187-99. PMID 20965447
19. Saltychev M, Juhola J. Effectiveness of High-Frequency Repetitive Transcranial Magnetic Stimulation in Migraine: A Systematic Review and Meta-analysis. *Am J Phys Med Rehabil.* Nov 01 2022; 101(11): 1001-1006. PMID 35034064
20. Food and Drug Administration. De Novo classification request for cerena transcranial magnetic stimulator (TMS) device. 2013;

- [https://www.accessdata.fda.gov/cdrh\\_docs/reviews/K130556.pdf](https://www.accessdata.fda.gov/cdrh_docs/reviews/K130556.pdf). Accessed August 20, 2025.
21. Goodman WK, Price LH, Rasmussen SA, et al. The Yale-Brown Obsessive Compulsive Scale. I. Development, use, and reliability. *Arch Gen Psychiatry*. Nov 1989; 46(11): 1006-11. PMID 2684084
  22. Storch EA, De Nadai AS, Conceição do Rosário M, et al. Defining clinical severity in adults with obsessive-compulsive disorder. *Compr Psychiatry*. Nov 2015; 63: 30-5. PMID 26555489
  23. Farris SG, McLean CP, Van Meter PE, et al. Treatment response, symptom remission, and wellness in obsessive-compulsive disorder. *J Clin Psychiatry*. Jul 2013; 74(7): 685-90. PMID 23945445
  24. Trevizol AP, Shiozawa P, Cook IA, et al. Transcranial Magnetic Stimulation for Obsessive-Compulsive Disorder: An Updated Systematic Review and Meta-analysis. *J ECT*. Dec 2016; 32(4): 262-266. PMID 27327557
  25. Liang K, Li H, Bu X, et al. Efficacy and tolerability of repetitive transcranial magnetic stimulation for the treatment of obsessive-compulsive disorder in adults: a systematic review and network meta-analysis. *Transl Psychiatry*. May 28 2021; 11(1): 332. PMID 34050130
  26. Carmi L, Tendler A, Bystritsky A, et al. Efficacy and Safety of Deep Transcranial Magnetic Stimulation for Obsessive-Compulsive Disorder: A Prospective Multicenter Randomized Double-Blind Placebo-Controlled Trial. *Am J Psychiatry*. Nov 01 2019; 176(11): 931-938. PMID 31109199
  27. Perera MPN, Mallawaarachchi S, Miljevic A, et al. Repetitive Transcranial Magnetic Stimulation for Obsessive-Compulsive Disorder: A Meta-analysis of Randomized, Sham-Controlled Trials. *Biol Psychiatry Cogn Neurosci Neuroimaging*. Oct 2021; 6(10): 947-960. PMID 33775927
  28. U.S. Food and Drug Administration. De novo classification request for Brainsway Deep Transcranial Magnetic Stimulation System. 2018; [https://www.accessdata.fda.gov/cdrh\\_docs/reviews/DEN170078.pdf](https://www.accessdata.fda.gov/cdrh_docs/reviews/DEN170078.pdf). Accessed August 20, 2025.
  29. Konstantinou G, Hui J, Ortiz A, et al. Repetitive transcranial magnetic stimulation (rTMS) in bipolar disorder: A systematic review. *Bipolar Disord*. Feb 2022; 24(1): 10-26. PMID 33949063
  30. Tee MMK, Au CH. A Systematic Review and Meta-Analysis of Randomized Sham-Controlled Trials of Repetitive Transcranial Magnetic Stimulation for Bipolar Disorder. *Psychiatr Q*. Dec 2020; 91(4): 1225-1247. PMID 32860557
  31. Qi L, Wang S, Li X, et al. Non-invasive brain stimulation in the treatment of generalized anxiety disorder: A systematic review and meta-analysis. *J Psychiatr Res*. Oct 2024; 178: 378-387. PMID 39208534
  32. Cui H, Jiang L, Wei Y, et al. Efficacy and safety of repetitive transcranial magnetic stimulation for generalised anxiety disorder: A meta-analysis. *Gen Psychiatr*. 2019; 32(5): e100051. PMID 31673675
  33. Li H, Wang J, Li C, et al. Repetitive transcranial magnetic stimulation (rTMS) for panic disorder in adults. *Cochrane Database Syst Rev*. Sep 17 2014; 2014(9): CD009083. PMID 25230088

34. Mantovani A, Aly M, Dagan Y, et al. Randomized sham controlled trial of repetitive transcranial magnetic stimulation to the dorsolateral prefrontal cortex for the treatment of panic disorder with comorbid major depression. *J Affect Disord.* Jan 10 2013; 144(1-2): 153-9. PMID 22858212
35. Trevizol AP, Barros MD, Silva PO, et al. Transcranial magnetic stimulation for posttraumatic stress disorder: an updated systematic review and meta-analysis. *Trends Psychiatry Psychother.* 2016; 38(1): 50-5. PMID 27074341
36. Brown R, Cherian K, Jones K, et al. Repetitive transcranial magnetic stimulation for post-traumatic stress disorder in adults. *Cochrane Database Syst Rev.* Aug 02 2024; 8(8): CD015040. PMID 39092744
37. Blyth SH, Cruz Bosch C, Raffoul JJ, et al. Safety of rTMS for Schizophrenia: A Systematic Review and Meta-analysis. *Schizophr Bull.* Mar 14 2025; 51(2): 392-400. PMID 39278637
38. He H, Lu J, Yang L, et al. Repetitive transcranial magnetic stimulation for treating the symptoms of schizophrenia: A PRISMA compliant meta-analysis. *Clin Neurophysiol.* May 2017; 128(5): 716-724. PMID 28315614
39. Dougall N, Maayan N, Soares-Weiser K, et al. Transcranial magnetic stimulation (TMS) for schizophrenia. *Cochrane Database Syst Rev.* Aug 20 2015; 2015(8): CD006081. PMID 26289586
40. Ye S, Guan X, Xiu M, et al. Early efficacy of rTMS intervention at week 2 predicts subsequent responses at week 24 in schizophrenia in a randomized controlled trial. *Neurotherapeutics.* Sep 2024; 21(5): e00392. PMID 38944636
41. Hua Q, Wang L, He K, et al. Repetitive Transcranial Magnetic Stimulation for Auditory Verbal Hallucinations in Schizophrenia: A Randomized Clinical Trial. *JAMA Netw Open.* Nov 04 2024; 7(11): e2444215. PMID 39527055
42. Xiu MH, Guan HY, Zhao JM, et al. Cognitive Enhancing Effect of High-Frequency Neuronavigated rTMS in Chronic Schizophrenia Patients With Predominant Negative Symptoms: A Double-Blind Controlled 32-Week Follow-up Study. *Schizophr Bull.* Sep 21 2020; 46(5): 1219-1230. PMID 32185388
43. Guan HY, Zhao JM, Wang KQ, et al. High-frequency neuronavigated rTMS effect on clinical symptoms and cognitive dysfunction: a pilot double-blind, randomized controlled study in Veterans with schizophrenia. *Transl Psychiatry.* Feb 25 2020; 10(1): 79. PMID 32098946
44. Kumar N, Vishnubhatla S, Wadhawan AN, et al. A randomized, double blind, sham-controlled trial of repetitive transcranial magnetic stimulation (rTMS) in the treatment of negative symptoms in schizophrenia. *Brain Stimul.* 2020; 13(3): 840-849. PMID 32289715
45. Zhuo K, Tang Y, Song Z, et al. Repetitive transcranial magnetic stimulation as an adjunctive treatment for negative symptoms and cognitive impairment in patients with schizophrenia: a randomized, double-blind, sham-controlled trial. *Neuropsychiatr Dis Treat.* 2019; 15: 1141-1150. PMID 31190822
46. Zhu L, Zhang W, Zhu Y, et al. Cerebellar theta burst stimulation for the treatment of negative symptoms of schizophrenia: A multicenter, double-blind, randomized controlled trial. *Psychiatry Res.* Nov 2021; 305: 114204. PMID 34587567
47. Jansen JM, Daams JG, Koeter MW, et al. Effects of non-invasive neurostimulation on craving: a meta-analysis. *Neurosci Biobehav Rev.* Dec 2013; 37(10 Pt 2): 2472-80. PMID 23916527



48. Chang CH, Liou MF, Liu CY, et al. Efficacy of Repetitive Transcranial Magnetic Stimulation in Patients With Methamphetamine Use Disorder: A Systematic Review and Meta-Analysis of Double-Blind Randomized Controlled Trials. *Front Psychiatry*. 2022; 13: 904252. PMID 35711590
49. Fang J, Zhou M, Yang M, et al. Repetitive transcranial magnetic stimulation for the treatment of amyotrophic lateral sclerosis or motor neuron disease. *Cochrane Database Syst Rev*. May 31 2013; 2013(5): CD008554. PMID 23728676
50. O'Connell NE, Wand BM, Marston L, et al. Non-invasive brain stimulation techniques for chronic pain. *Cochrane Database Syst Rev*. Apr 11 2014; (4): CD008208. PMID 24729198
51. O'Connell NE, Marston L, Spencer S, et al. Non-invasive brain stimulation techniques for chronic pain. *Cochrane Database Syst Rev*. Apr 13 2018; 4(4): CD008208. PMID 29652088
52. Jiang X, Yan W, Wan R, et al. Effects of repetitive transcranial magnetic stimulation on neuropathic pain: A systematic review and meta-analysis. *Neurosci Biobehav Rev*. Jan 2022; 132: 130-141. PMID 34826512
53. Chen R, Spencer DC, Weston J, et al. Transcranial magnetic stimulation for the treatment of epilepsy. *Cochrane Database Syst Rev*. Aug 11 2016; (8): CD011025. PMID 27513825
54. Mishra A, Maiti R, Mishra BR, et al. Effect of Repetitive Transcranial Magnetic Stimulation on Seizure Frequency and Epileptiform Discharges in Drug-Resistant Epilepsy: A Meta-Analysis. *J Clin Neurol*. Jan 2020; 16(1): 9-18. PMID 31942753
55. Ahmed I, Mustafaoglu R, Memon AR, et al. Comparative Effectiveness of Noninvasive Brain Stimulation for the Treatment of Pain, Fatigue, and Sleep Quality in Fibromyalgia. A Systematic Review With Network Meta-Analysis. *Clin J Pain*. May 01 2025; 41(5). PMID 40091857
56. Su YC, Guo YH, Hsieh PC, et al. Efficacy of Repetitive Transcranial Magnetic Stimulation in Fibromyalgia: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *J Clin Med*. Oct 12 2021; 10(20). PMID 34682790
57. Saltychev M, Laimi K. Effectiveness of repetitive transcranial magnetic stimulation in patients with fibromyalgia: a meta-analysis. *Int J Rehabil Res*. Mar 2017; 40(1): 11-18. PMID 27977465
58. Chou YH, Hickey PT, Sundman M, et al. Effects of repetitive transcranial magnetic stimulation on motor symptoms in Parkinson disease: a systematic review and meta-analysis. *JAMA Neurol*. Apr 2015; 72(4): 432-40. PMID 25686212
59. Shirota Y, Ohtsu H, Hamada M, et al. Supplementary motor area stimulation for Parkinson disease: a randomized controlled study. *Neurology*. Apr 09 2013; 80(15): 1400-5. PMID 23516319
60. Li R, He Y, Qin W, et al. Effects of Repetitive Transcranial Magnetic Stimulation on Motor Symptoms in Parkinson's Disease: A Meta-Analysis. *Neurorehabil Neural Repair*. Jul 2022; 36(7): 395-404. PMID 35616427
61. Hao Z, Wang D, Zeng Y, et al. Repetitive transcranial magnetic stimulation for improving function after stroke. *Cochrane Database Syst Rev*. May 31 2013; 2013(5): CD008862. PMID 23728683
62. Le Q, Qu Y, Tao Y, et al. Effects of repetitive transcranial magnetic stimulation on hand function recovery and excitability of the motor cortex after stroke: a meta-analysis. *Am J Phys Med Rehabil*. May 2014; 93(5): 422-30. PMID 24429509

63. Li Y, Qu Y, Yuan M, et al. Low-frequency repetitive transcranial magnetic stimulation for patients with aphasia after stroke: A meta-analysis. *J Rehabil Med*. Sep 2015; 47(8): 675-81. PMID 26181486
64. Qiao J, Ye QP, Wu ZM, et al. The Effect and Optimal Parameters of Repetitive Transcranial Magnetic Stimulation on Poststroke Dysphagia: A Meta-Analysis of Randomized Controlled Trials. *Front Neurosci*. 2022; 16: 845737. PMID 35573312
65. Zhang L, Xing G, Fan Y, et al. Short- and Long-term Effects of Repetitive Transcranial Magnetic Stimulation on Upper Limb Motor Function after Stroke: a Systematic Review and Meta-Analysis. *Clin Rehabil*. Sep 2017; 31(9): 1137-1153. PMID 28786336
66. Graef P, Dadalt MLR, Rodríguez DAMDS, et al. Transcranial magnetic stimulation combined with upper-limb training for improving function after stroke: A systematic review and meta-analysis. *J Neurol Sci*. Oct 15 2016; 369: 149-158. PMID 27653882
67. Zhang JJY, Ang J, Saffari SE, et al. Repetitive Transcranial Magnetic Stimulation for Motor Recovery After Stroke: A Systematic Review and Meta-Analysis of Randomized Controlled Trials With Low Risk of Bias. *Neuromodulation*. Jan 2025; 28(1): 16-42. PMID 39320286
68. Murphy TK, Lewin AB, Storch EA, et al. Practice parameter for the assessment and treatment of children and adolescents with tic disorders. *J Am Acad Child Adolesc Psychiatry*. Dec 2013; 52(12): 1341-59. PMID 24290467
69. McClintock SM, Reti IM, Carpenter LL, et al. Consensus Recommendations for the Clinical Application of Repetitive Transcranial Magnetic Stimulation (rTMS) in the Treatment of Depression. *J Clin Psychiatry*. 2018; 79(1). PMID 28541649
70. VA/DoD Clinical Practice Guideline. (2022). The Management of Major Depressive Disorder. Washington, DC: U.S. Government Printing Office.  
<https://www.healthquality.va.gov/guidelines/MH/mdd/VADoDMDDCPGFinal508.pdf>. Accessed August 19, 2025.
71. VA/DoD Clinical Practice Guideline. (2023). VA/DoD Clinical Practice Guideline for Management of Bipolar Disorder. Washington, DC: U.S. Government Printing Office.  
<https://www.healthquality.va.gov/guidelines/MH/bd/VA-DoD-CPG-BD-Full-CPGFinal508.pdf>. Accessed August 20, 2025.
72. VA/DoD Clinical Practice Guideline. (2023). VA/DoD Clinical Practice Guideline for Management of Posttraumatic Stress Disorder and Acute Stress Disorder. Washington, DC: U.S. Government Printing Office.  
<https://www.healthquality.va.gov/guidelines/MH/ptsd/VA-DoD-CPG-PTSD-Full-CPGAug242023.pdf>. Accessed August 18, 2025.
73. National Institute for Health and Care Excellence (NICE). Repetitive transcranial magnetic stimulation for depression [IPG542]. 2015; <https://www.nice.org.uk/guidance/ipg542>. Accessed August 17, 2025.
74. National Institute for Health and Care Excellence (NICE). Transcranial magnetic stimulation for treating and preventing migraine [IPG477]. 2014; <https://www.nice.org.uk/guidance/ipg477>. Accessed August 20, 2025.
75. National Institute for Health and Care Excellence (NICE). Transcranial magnetic stimulation for auditory hallucinations [IPG680]. 2020; <https://www.nice.org.uk/guidance/ipg680/chapter/1-Recommendations>. Accessed August 18, 2025.

76. National Institute for Health and Care Excellence (NICE). Transcranial magnetic stimulation for obsessive-compulsive disorder [IPG676]. 2020; <https://www.nice.org.uk/guidance/ipg676>. Accessed August 19, 2025.
77. Leung A, Shirvalkar P, Chen R, et al. Transcranial Magnetic Stimulation for Pain, Headache, and Comorbid Depression: INS-NANS Expert Consensus Panel Review and Recommendation. *Neuromodulation*. Apr 2020; 23(3): 267-290. PMID 32212288

## **OTHER REFERENCES**

1. Blue Cross and Blue Shield of Kansas Behavioral Health Liaison Committee, June 2012; June 2014; September 2014 (CB); June 2015; August 2022; February 2023, February 2024, February 2025.
2. BCBSKS Medical Consultant, Practicing Board Certified Psychiatrist, November 2012.